



UNICA

UNIVERSITÀ
DEGLI STUDI
DI CAGLIARI



Università di Cagliari

UNICA IRIS Institutional Research Information System

This is the Author's [*accepted*] manuscript version of the following contribution:

Fattore L, Pisanu A, Concas L, Casula C, Siddi C, Pisu MG, **Serra M**, Concas A, Porcu P. Behavioral characterization of co-exposure to cannabinoids and hormonal contraceptives in female rats. Prog Neuropsychopharmacol Biol Psychiatry, 129: 110890, 2024.

The publisher's version is available at:

<http://dx.doi.org/10.1016/j.pnpbp.2023.110890>

When citing, please refer to the published version.

Behavioral characterization of co-exposure to cannabinoids and hormonal contraceptives in female rats

Liana Fattore^a, Augusta Pisanu^a, Luca Concas^{a*}, Claudia Casula^a, Carlotta Siddi^{a*}, Maria Giuseppina Pisu^a, Mariangela Serra^b, Alessandra Concas^b, Patrizia Porcu^a

^a*Neuroscience Institute, National Research Council of Italy (CNR), Cagliari, Italy*

^b*Department of Life and Environment Sciences, Section of Neuroscience and Anthropology, University of Cagliari, Cagliari, Italy*

**present address: Department of Biomedical Sciences, University of Cagliari, Cagliari, Italy*

Corresponding Author:

Patrizia Porcu, Ph.D.

Neuroscience Institute

National Research Council of Italy (CNR)

Cittadella Universitaria, 09042 Monserrato, Cagliari, Italy

Phone: +39-070-6754163

Email: patrizia.porcu@in.cnr.it

Abbreviations

ANOVA, analysis of variance; CB₁, cannabinoid receptor type 1; EDTA, ethylenediaminetetraacetic acid; EE, ethinyl estradiol; EPM, elevated plus maze; FST, forced swimming test; HC, hormonal contraceptive; i.v., intravenous; LA, locomotor activity; LNG, levonorgestrel; MB, marble burying; NOR, novel object recognition; OCD, obsessive-compulsive disorder; PND, post-natal day; PPI, prepulse inhibition; SAL, saline; T1, training session; T2, choice session; THC, Δ^9 -tetrahydrocannabinol; UNODC, United Nations Office on Drugs and Crime; VEH, vehicle; WIN, WIN 55,212-2

Abstract

Hormonal contraceptives are among the most widely used drugs by young healthy women to block ovulation and avoid pregnancy. They reduce the ovarian secretion of estradiol and progesterone, hormones that also modulate neuronal plasticity, cognitive functions, emotions and mood. Cannabis is the most commonly used illicit drug worldwide and its use is increasing among young women, many of which regularly take the “pill”. Despite evidence of a bidirectional interaction between the endocannabinoid system and gonadal hormones, only very few studies have examined the consequences of cannabis consumption in young females under hormonal contraceptives treatment. To fill this gap, this study evaluated the behavioral effects of co-exposure to chronic 1) hormonal contraceptives, i.e., ethinyl estradiol (EE) plus levonorgestrel (LNG), one of the synthetic estrogen-progestin combinations of hormonal contraceptives, and 2) a cannabinoid receptor agonist, i.e., WIN 55,212-2 (WIN), on motor activity, emotional state and cognitive functions in young adult female rats (8-11/experimental group). Hormonal and cannabinoid treatment started at post-natal day (PND) 52 and 56, respectively, while behavioral testing occurred between PND 84-95. The results show that chronic EE-LNG treatment, at doses (0.020 and 0.060 mg/rat, respectively) known to drastically reduce plasma progesterone levels, and the contextual exposure to WIN, at a dose (12.5 µg/kg/infusion) known to be rewarding in the rat, alters the hormonal milieu but does not cause further changes in locomotor activity compared to EE-LNG or WIN alone, and does not modify anxiety-like state (as measured by the elevated plus maze and the marble burying tests) and cognitive abilities (as measured by the novel object recognition and the prepulse inhibition tests) in young adult female rats. Although exposure to EE-LNG and WIN tends to increase the duration of immobility and to reduce the time spent swimming in the forced swimming test, there was not a significant additive effect suggestive of a depressive-like state. These findings allow deepening the current knowledge on the interaction between cannabinoid agonists and hormonal contraceptives and suggest that low, rewarding doses of cannabinoids do not significantly alter the motor and cognitive


skills and do not induce anxiety or depressive-like states in females that use hormonal contraceptives.

Keywords: hormonal contraceptives; cannabinoid agonists; ethinyl estradiol; levonorgestrel; emotional state; cognitive function

Highlights

- Hormonal contraceptives - cannabinoid agonist behavioral interactions in rats
- Hormonal contraceptives plus cannabinoids do not alter spontaneous locomotor activity
- Hormonal contraceptives plus cannabinoids do not alter emotional state
- Hormonal contraceptives plus cannabinoids do not alter cognitive abilities

Graphical abstract



1. Introduction

Steroid hormones, synthesized both in the periphery and the central nervous system, influence key brain functions through rapid modulation of neuronal excitability via membrane receptors, or by promoting gene expression via intracellular steroid receptors (McEwen, 1994; Rupprecht, 2003). These so called neurosteroids regulate mood, emotional states, the stress response and cognition, and their levels are altered in neuropsychiatric disorders, including anxiety and depression, and following administration of psychoactive substances (Belelli and Lambert, 2005; Porcu et al., 2016; Boero et al., 2020; Paul et al., 2020). Chronic treatment with ethinyl estradiol (EE) and levonorgestrel (LNG), two of the synthetic estrogens and progestins used in the hormonal contraceptive (HC) combinations (Levin et al., 2018), markedly decrease brain and plasma levels of progesterone and its neuroactive metabolite allopregnanolone in female rats (Concas et al., 2022). HCs have been used for almost 60 years by more than 150 million women worldwide, mostly for contraceptive purposes (United Nations, 2019), but also as a therapy for endometriosis, polycystic ovarian syndrome and menstrual cycle-related symptoms (Levin et al., 2018). Several studies have shown that HCs affect the central nervous system, altering the volume and structure of several brain regions with subsequent changes in brain function that may persist even after discontinuation (Petersen et al., 2023). This is particularly concerning for the increasing number of young women who start taking HCs early in adolescence, when the brain is still under development and influenced by steroid hormones (Cahill, 2018).

Administration of several drugs of abuse, including alcohol, nicotine, morphine, and Δ^9 -tetrahydrocannabinol (THC), alters brain and plasma pregnenolone, progesterone and allopregnanolone levels (Concas et al., 2006; Grobin et al., 2005; Morrow et al., 2020; Vallée et al., 2014), and such alterations are thought to strengthen the rewarding properties of these drugs. Indeed, neurosteroids display rewarding properties in behavioral paradigms of drug addiction (Finn et al., 1997; Sinnott et al., 2002) and modulate dopamine release in the rat nucleus accumbens and prefrontal cortex (Motzo et al., 1996; Rougé-Pont et al., 2002). Likewise, estradiol modulates motivation and dopamine release (Yoest et al., 2014),

enhancing the rewarding properties of alcohol (Hilderbrand and Lasek, 2018) and psychostimulants (Quigley et al., 2021), although it also decreases risky choice and promotes risk aversion in both male and female rats (Orsini et al., 2021). Thus, chronic treatment with HCs might alter reward threshold and sensitivity to drugs of abuse, a notion strengthened by substantial evidence showing that females are more vulnerable to drug abuse than males, and that endogenous and exogenous steroid hormones are likely involved in this phenomenon (Quigley et al., 2021).

According to the United Nations Office on Drugs and Crime (UNODC)'s World Drug Report 2022, marijuana is still the most commonly abused drug worldwide, particularly among young adults, and cannabis legalization in several countries appears to have accelerated daily use and related health impacts. Cannabis consumption increased during the COVID-19 pandemic, and almost 6% of the global cannabis users estimated in 2020 (209 million people) were 15-16 years old (UNODC, 2022). Worryingly, early onset of cannabis use among adolescents has been shown to be associated with cognitive impairment (Pope et al., 2003), enhanced risk for depression (Hengartner et al., 2020), and transition from recreational use to cannabis use disorder (Behrendt et al., 2009). Adolescence is a critical period for brain development, marked by the increasing levels of gonadal hormones including estrogens and progesterone. These steroids interact with the endocannabinoid system in modulating, among others, the reproductive system, motivational processes and sensitivity to exogenous cannabinoids (Gorzalka and Dang, 2012; Struik et al., 2018). Bi-directional interactions between (endo)cannabinoids and gonadal hormones could account, at least in part, for the greater sensitivity of women to the subjective effects of cannabis, which may contribute to the faster progression from first use to abuse and the greater withdrawal symptoms of women relative to men (Cooper and Haney, 2014). In support of an enhanced vulnerability of females to cannabis abuse, female rats were shown to be more sensitive to the reinforcing effects of cannabinoids (Fattore et al., 2007) and at higher risk to reinstate cannabinoid-seeking behavior after extinction (Fattore et al., 2010), with gonadal

hormones significantly affecting cannabinoid CB₁ receptor density and function in selected brain areas (Castelli et al., 2014).

Since approximately 25% of women aged 15-44 who currently use contraception reported using the pill as their preferred method (Cooper et al., 2022) and the prevalence of cannabis use is increasing among the female population (UNODC, 2022), it can be easily assumed that a high number of girls take HCs and smoke cannabis. On the basis of the clinical and preclinical evidence available in the literature, concomitant use of exogenous steroid hormones and cannabinoids are very likely to interact and affect brain and behavior (Gorzalka and Dang, 2012). Yet, the effect of smoking marijuana in girls under HC treatment is still to be determined. Similarly, no preclinical study has been performed so far to test the effect of rewarding doses of cannabinoids in young females exposed to chronic HC treatment. In this study, we used an animal model of chronic HC exposure capable to effectively blunt the hypothalamic-pituitary-ovarian axis, thus decreasing brain and plasma concentrations of neurosteroids with subsequent changes in behavior (see Porcu et al., 2019, and Concas et al., 2022 for review). We thus tested the hypothesis that co-exposure to EE-LNG, one of the estrogen-progestin combinations of the HC pills, and to WIN 55,212-2 (WIN), a synthetic cannabinoid agonist known to induce rewarding effects and sustain motivated behavior in rats, may affect motor, emotional and cognitive behavior, as well as plasma progesterone in female rats.

2. Methods

2.1 Animals

Female Sprague-Dawley rats (Charles River, Italy) aged 32-35 days were housed 4/cage and maintained on an artificial 12 h light/dark inverted cycle (light ON at 7 PM) at a constant temperature of 22±2°C, a relative humidity of 65% and with food and water available *ad libitum*. After arrival in the animal facility, rats were handled once daily for 5 minutes for 10 consecutive days before undergoing surgery for the implantation of an intravenous catheter, after which they were housed individually to avoid damage to the external part of the

catheter.

All measures were taken to minimize pain or discomfort of animals whose care and handling throughout the experimental procedures were in accordance with the European Parliament and the Council Directive of 22 September 2010 (2010/63/EU), the Italian Legislative decree n. 26, 4 March 2014, and were approved by the Italian Ministry of Health (authorization n. 969/2019-PR).

2.2 Experimental procedure. Surgery for intravenous catheter implantation (PND 42-45)

Following 10 days of acclimation, starting on post-natal day (PND) 42, animals were deeply anaesthetized with isoflurane (3%) and a permanent intravenous (i.v.) catheter was surgically implanted into the external jugular vein as previously described (Fattore et al., 2021). Briefly, under sterile conditions, a silastic catheter (L-155; Silmedic medical grade, Gessil, France) was inserted into the right jugular vein, tunnelled under the skin and exited in the midscapular region, where it was bonded to a larger diameter extension (L-205) and anchored to the back of the neck with two sutures. During the 7 days following surgery, rats were allowed to recover individually in their own home cage and received a daily i.v. infusion of gentamicin (0.16 mg/kg) followed by 0.2 ml of a heparinized (1%) sterile saline solution to flush the antibiotic through the catheter. Antibiotics and anesthetics were purchased as sterile solutions from local distributors. As shown in Figure 1, at the end of the 10-day recovery, animals started the chronic treatment with HCs.

2.3 Drugs

2.3.1 Hormonal contraceptives treatment (PND 52-97)

The hypothalamic-pituitary-ovarian axis in female rats is fully functional by PND 48, when estradiol reaches levels comparable to those in adult females (Vetter-O'Hagen and Spear, 2012). Given that HCs inhibit hypothalamic-pituitary-ovarian axis function to prevent ovulation, EE-LNG treatment started at PND 52 to avoid interfering with its physiological maturation. Rats were injected subcutaneously at the nape of the neck with either a

combination of EE (0.020 mg, Sigma-Aldrich E4876) and LNG (0.060 mg, Sigma-Aldrich N2260) (EE-LNG-treated group) or vehicle (VEH-treated group) once a day at 4 PM. The EE-LNG combination was emulsified in 1% Tween 80 before being diluted in distilled water, vortexed and immediately administered in a volume of 1 ml/rat (Follesa et al., 2002; Porcu et al., 2012). Animals were subjected to behavioral tests, or were euthanized 20 to 24 hours after the last EE-LNG daily treatment (see timeline on Figure 1).

2.3.2 Cannabinoid treatment (PND 56-98)

Daily cannabinoid treatment started from PND 56. WIN (Tocris, UK) was dissolved in one drop of Tween 80 and then diluted in heparinized (1%) 0.9% sterile saline solution and i.v. infused at a unit dose of 12.5 µg/kg/0.1 ml over the 2-h session until reaching a cumulative dose of 0.3 mg/kg (WIN-treated group). This WIN dosage regimen was selected based on previous studies showing that it increases dopamine release in the rat nucleus accumbens (Fadda et al., 2006) and corresponds to the mean amount of cannabinoid agonist daily self-administered by trained rats (Fattore et al., 2007; Fattore et al., 2010), which suggests it induces rewarding effects. A passive protocol was adopted (i) to better focus on the pharmacological effects of the co-administration of HCs and cannabinoid agonists, which requires animals to be exposed to precise dosages of these drugs (EE-LNG, WIN), and (ii) to avoid any individual between-subjects differences in drug intake by exposing animals to the same amount of cannabinoid at the same time intervals. To closely mimic the self-administration pattern of cannabinoid intake observed in our previous studies (Fattore et al., 2007; Fattore et al., 2010), rats were daily placed in self-administration boxes for 2 hours and WIN was infused at the same rate of infusion (100 µl/5 s), at the same unit dose (12.5 µg/kg/infusion) and approximately at the same time interval until reaching a cumulative i.v. dose of 0.3 mg/kg. Specifically, rats received 12 infusions during the 1st half hour of the session, 7 infusions during the 2nd half hour, 3 infusions during the 60-90 min interval and 2 infusions during the last 30-min of the session. In parallel, control rats received i.v. infusion of sterile saline (SAL-treated group) under the same experimental conditions (i.e., with the

only difference of receiving saline in the place of the cannabinoid). During each 2-h session, approximately 50% of EE-LNG- or VEH-treated animals were infused with WIN (WIN-treated group), while the remaining 50% with saline (SAL-treated group). Throughout the entire duration of the study, locomotor activity was monitored inside the operant chambers, which were equipped with a series of photocells located 3.5 cm above the cage floor. The number of photocell beam breaks was recorded and served as a measure of general horizontal locomotor activity. Rats were euthanized on PND 98, 2 hours after the WIN/SAL administration session (see timeline on Figure 1). WIN treatment took place once daily, 7 days/week, during the dark phase of the cycle (between 9.30 AM and 13.30 PM) and was performed in operant chambers (29.5×32.5×23.5 cm), each encased in a sound- and light-attenuating cubicle equipped with a ventilation fan (Med Associates, USA). In each chamber, the two levers (used by rats in self-administration studies as *modus operandi* to obtain the drug) were kept retracted during the 2-h session, and WIN (or saline) was delivered through a swivel device connected to the implanted catheter by an infusion pump controlled remotely by the experimenter through a computer with Med Associates software. The WIN solution was freshly remixed every day before use.

2.4 Experimental procedure. Behavioral testing (PND 84-95)

Behavioral tests started at PND 84, i.e., after 33 days of EE-LNG/VEH treatment and 29 days of WIN/SAL infusion. Experiments were conducted during the dark phase (between 1 PM and 4 PM) in dedicated rooms under dim light (20 lux). Rats were left undisturbed in each experimental room for 1 hour acclimatation period before starting behavioral testing. Each animal underwent the following sequence of behavioral tests at least 1 hour after the last WIN infusion: locomotor activity (LA), novel object recognition (NOR), forced swimming (FST), marble burying (MB), elevated plus maze (EPM), and prepulse inhibition (PPI), with at least 45 hours (i.e., every two days) or 69 hours (i.e., every three days) separating two consecutive test to prevent carryover effects (see timeline on Figure 1).

2.4.1 Locomotor activity

Rats were individually tested for LA under standardized environmental conditions using the Digiscan Animal Activity Analyzer (Omnitech Electronics, USA) that assessed animals' activity for 60 minutes at 10-min intervals, beginning immediately after the animal was placed in the cage (Spano et al., 2013). Each operant cage (42×30×60 cm) was equipped with two sets of 16 photocells located at right angles to each other, projecting horizontal infrared beams 2.5 cm apart and 4 cm above the cage floor, and a further set of 16 horizontal beams whose height could be adapted to the size of the animals. LA was performed in a dark room dimly illuminated by a red lamp. The following behavioral parameters were measured: 1) horizontal activity, i.e., the total number of beam interruptions that occurred in the horizontal sensors; 2) vertical activity, i.e., the total number of beam interruptions that occurred in the vertical sensors, i.e., the number of times the rat rose onto its hind legs either against the wall or freely in the air (number of rearings); 3) total distance travelled (cm), i.e. the horizontal distance travelled by the animal during the 60-min session. The arena was wiped out with H₂O₂ before each trial to prevent olfactory cues.

2.4.2 Novel object recognition

Ability to discriminate novel objects in the NOR test is a marker of non-spatial short-term memory. The NOR test was carried out in a black box (60×60 cm) according to our previous studies (Boi et al., 2022). Briefly, each rat (with no prior experience of the objects to be discriminated) was given a 10-min habituation session (habituation trial) and 3 min later placed in the box and allowed to explore two identical objects for 10 min (training session, T1). After a 1-h interval, the second trial (choice session, T2) started and lasted 3 min, with 2 objects placed in the same positions and the animal exposed to the same condition as in T1, except that one object was replaced with a new one (novel object). The objects to be discriminated were made of glass, plastic or metal and had different shapes and colors, devoid of any natural significance, and were thoroughly cleaned with H₂O₂ before each trial to avoid olfactory cues. Exploratory behavior was defined as the animal directing its nose

toward the object at a distance ≤ 2 cm and/or sniffing, biting or touching the objects with the nose. Sitting on, climbing over and/or turning around the objects were not considered exploratory behaviors. Combination (novel/old) and position (right/left) of objects were counterbalanced to compensate for lateralization. Training and test sessions were recorded with a camera and the total time spent exploring the objects during T1 and T2 and the frequency of approaches, i.e., numbers of approaching the two objects, were measured by three independent researchers blind to the experimental condition. Recognition skills were calculated using the following formula: $N/(N+F) \times 100$, where N=time spent exploring (or frequency of contacts with) the novel object, and F=time spent exploring (or frequency of contacts with) the familiar one.

2.4.3 Forced swimming test

Introduced by Porsolt et al. in 1977 for rats (Porsolt et al., 1977a) and mice (Porsolt et al., 1977b), the FST is the most commonly used test in animal studies for screening antidepressant drugs or for evaluating the effect of environmental or pharmacological manipulations on mood. Being able to induce a depressive-like state in rodents that are forced to swim without possibility of escaping, this test has been classified as one of the two only high-ranking tests in the categories of reliability, specificity, and ease of use (Cryan et al., 2002). The FST was conducted according to our previous studies (Zanda et al., 2017). Briefly, animals were individually placed in transparent Plexiglas cylinders (height: 40 cm; diameter: 20 cm) containing 20 cm of water at 25°C. After 5 minutes, they were removed from the water and thoroughly dried in a heated environment (30°C) for 30 minutes. The sessions were videotaped and the videos analyzed by three independent researchers blind to the experimental condition. Animals' behaviors were classified according to three possible and mutually exclusive categories: swimming, climbing (i.e., attempts to climb the wall of the cylinder), and immobility (i.e., floating in the water performing the minimum amount of movements with the anterior paws to maintain its head above the water surface). The rats' emotional state was determined by quantifying the time spent swimming or trying to escape

in any direction (sign of reaction) or staying completely immobile (sign of behavioral hopelessness or passivity). An increase or decrease in immobility time is considered a sign of a depressant or antidepressant-like effect, respectively.

2.4.4 Marble burying

The MB test, a cost-effective rodent model with good face validity, commonly used to quantify anxiety, repetitive or obsessive-compulsive-like behavior in rodents (Jimenez-Gomez et al., 2011), was conducted in open transparent plastic boxes (54×34.5×20 cm) with the floor covered with 5 cm of fresh sawdust as previously described (Zanda et al., 2017). In rodents, the burying of the marble is a natural defense mechanism that occurs under anxiety or conditions of stress, or fear from new or strange objects (i.e., neophobia). Twenty-four standard glass marbles (1.5 cm in diameter, arranged in six rows of four marbles each) were placed uniformly over the sawdust surface. On the test day, rats were individually placed in the cage in a marble-free area (34.5×15 cm) and their activity was videotaped for 30 min. At the end of the session, animals were gently removed from the test cage and the number of marbles partially ($\geq 67\%$) and totally ($>95\%$) buried was counted by an experimenter blind to the pharmacological treatments. Bedding was replaced and marbles were thoroughly cleaned with soapy water, followed by H_2O_2 , before each test to avoid the presence of olfactory cues.

2.4.5 Elevated plus maze

The EPM test is among the most commonly used tests to assess anxiety-like behavior in rodents; it is based on the inclination of rodents to explore novel environments and their innate avoidance for bright, unprotected, and elevated places, as the open arms of the apparatus are. The test was conducted as previously described (Porcu et al., 2012) in a black polyvinyl chloride maze equipped with two open and two closed arms (12×60 cm) that converged on a small central square (12×12 cm), which served as starting point, thus reproducing the shape of a plus sign; the apparatus was elevated 50 cm from the floor. The

rat was placed in the center of the maze apparatus (i.e., at the start point) facing an open arm, and allowed to freely explore for 5 minutes. Rats' behavior was videotaped by a camera positioned above the maze. Three independent researchers blind to the treatment group measured the time spent by each rat in the open vs. the closed arms and counted the number of entries into open and closed arms, considering the animal inside a specific arm when it had all the four paws inside that arm. The EPM was thoroughly cleaned with H₂O₂ before each test to avoid olfactory cues.

2.4.6 Prepulse inhibition

The startle response is an unconditioned reflexive response to an unexpected environmental stimulus. Prepulse inhibition (PPI, a sensorimotor gating reflex) is a phenomenon in which a weak pre-stimulus (i.e., prepulse) suppresses the response to a startling stimulus. Deficits in PPI reflect attentional dysfunctions and are common in schizophrenic patients. The PPI test is one of the few paradigms in which humans and rodents are tested in a similar fashion (Geyer and Swerdlow, 2001). The apparatus for the detection of the startle reflex consisted of 4 standard cages, each placed inside a sound-attenuated and ventilated chamber (Med Associated, USA). Startle cages were non-restrictive Plexiglas cylinders (diameter 9 cm) mounted on a piezoelectric accelerometer platform connected to an analogue-digital converter. Background noise and acoustic bursts were conveyed through two speakers, placed in proximity to the startle cage, in order to produce a variation in sound intensity within 1 dB. Both startle cages and speakers were connected to a PC dedicated to the detection and analysis of all variables of the cage. Before each session, calibration of acoustic stimuli and mechanical responses was checked. On test day, rats were individually placed in the test cage for a 5-min acclimatization period with a 70 dB white noise background that continued for the remainder of the session. Animals were then tested on 3 consecutive blocks of trials. The 1st and 3rd blocks consisted of 5 pulse-alone trials of 40 ms at 115 dB, while the 2nd block (test block) was a pseudorandom sequence of 50 trials including 12 pulse-alone trials, 30 pulse trials preceded by 73, 76 or 82 dB prepulses of 20

milliseconds (10 for each level of prepulse loudness), and 8 no-stimulus trials where the only background noise was delivered. Inter-trial intervals were selected randomly between 10 and 15 s, while the inter-stimulus intervals were set at 100 milliseconds. The percent (%) PPI was calculated based on the values relative to the second block only and using the following formula: $100 - [(mean\ startle\ amplitude\ for\ prepulse+pulse\ trials / mean\ startle\ amplitude\ for\ pulse-alone\ trials) \times 100]$, as previously described (Spano et al., 2010).

2.5 Progesterone assay

On PND 98, three days after the last behavioral test, rats were euthanized by decapitation; blood was collected from the trunk into ethylenediaminetetraacetic acid (EDTA)-coated tubes and centrifuged at 900xg for 10 min at 4°C. The resulting plasma was frozen at -20°C until use. Progesterone levels were quantified by an enzyme-linked immunosorbent assay kit (DEV9988, Demeditec Diagnostics GmbH, Germany), according to manufacturer's instructions, using a 96-well plate pre-coated with progesterone antiserum. Each sample was run in duplicate. The lowest detectable level is 0.156 ng/ml; within- and between-assay coefficients of variation ranged 5-12% and 6-14%, respectively.

2.6 Statistical analysis

A sample size of 7.75 per experimental group was estimated using G*Power software (G*Power 3.1.9.2; based on $\alpha=0.05$, power=0.80), a value consistent with the numbers used in the past for similar studies. Thus, we determined a sample size of ≥ 8 per experimental group throughout the experiment (specifically, 9 Vehicle + Saline, 11 Vehicle + WIN, 8 EE-LNG + Saline and 11 EE-LNG + WIN rats/group). Statistical analysis was performed using commercially available statistical programs (Statistica 8.0, StatSoft Inc., Tulsa, OK, USA; GraphPad Prism 8.0, GraphPad Software, San Diego, CA, USA). Normality was checked by the Kolmogorov-Smirnov test, which indicated that data met the requirement for parametric analysis. Data are presented as mean $\pm$ SEM and statistical comparisons were performed by two-way analysis of variance (ANOVA), with EE-LNG and WIN treatments as factors, except

for locomotor activity whose data were analyzed by two-way repeated measures ANOVA, with EE-LNG and WIN treatments as factors and time as repeated measure. ANOVAs were followed by Tukey's multiple comparisons *post hoc* test. A *p* value <0.05 was considered statistically significant.

3. Results

3.1 Chronic co-exposure to EE-LNG and WIN does not alter spontaneous locomotor activity

Locomotor activity decreased over time to a similar extent in all experimental groups (Figure 2). A significant effect of time was observed for horizontal activity (Fig. 2A, $F_{5,175}=99.96$; $p<0.00001$), vertical activity (Fig. 2B, $F_{5,175}=41.90$; $p<0.00001$), and total distance travelled (Fig. 2C, $F_{5,175}=99.43$; $p<0.00001$). For vertical activity, two-way ANOVA also revealed a significant effect of WIN treatment ($F_{1,35}=6.56$; $p=0.015$) and a time x WIN treatment interaction ($F_{5,175}=3.53$; $p=0.005$); likewise, for total distance travelled there was a significant time x WIN treatment interaction ($F_{5,175}=4.25$; $p=0.001$) (see Supplementary Tables 1 and 2 for a summary of the ANOVA and *post hoc* results). WIN-treated rats display a trend for a reduced vertical activity and total distance travelled, compared to vehicle+saline- and EE-LNG+saline-treated rats; however, the *post hoc* test failed to reveal significant differences among the four experimental groups.

3.2 Chronic co-exposure to EE-LNG and WIN does not induce neophobia / compulsive traits in the marble burying test

No differences among experimental groups were observed in the number of fully (>95%) buried marbles (Fig. 3A, EE-LNG treatment: $F_{1,35}=0.001$, $p=0.974$; WIN treatment: $F_{1,35}=1.22$, $p=0.276$; interaction $F_{1,35}=0.04$, $p=0.848$), partially ($\geq 67\%$) buried marbles (Fig. 3B, EE-LNG treatment: $F_{1,35}=2.30$, $p=0.138$; WIN treatment: $F_{1,35}=1.22$, $p=0.277$; interaction $F_{1,35}=0.46$, $p=0.502$), or in the total number (fully+partially) of buried marbles (Fig. 3C, EE-LNG treatment: $F_{1,35}=0.31$, $p=0.582$; WIN treatment: $F_{1,35}=1.74$, $p=0.195$; interaction

$F_{1,35}=0.16$, $p=0.691$). Overall, chronic treatment with EE-LNG and co-exposure with WIN did not affect burying behavior in female rats.

3.3 Chronic co-exposure to EE-LNG and WIN does not alter emotional behavior in the elevated plus maze and forced swimming tests

Regarding anxiety-like behavior, no differences among experimental groups were observed in the number of open arm entries (Fig. 4A, EE-LNG treatment: $F_{1,35}=0.67$, $p=0.419$; WIN treatment: $F_{1,35}=0.52$, $p=0.477$; interaction $F_{1,35}=0.45$, $p=0.507$) or time spent in the open arms (Fig. 4B, EE-LNG treatment: $F_{1,35}=0.009$, $p=0.923$; WIN treatment: $F_{1,35}=1.63$, $p=0.209$; interaction $F_{1,35}=0.52$, $p=0.477$). Likewise, no significant differences were observed in the number of closed arm entries (EE-LNG treatment: $F_{1,35}=1.25$, $p=0.271$; WIN treatment: $F_{1,35}=3.62$, $p=0.065$; interaction $F_{1,35}=0.02$, $p=0.873$) or time spent in the closed arms (EE-LNG treatment: $F_{1,35}=0.74$, $p=0.394$; WIN treatment: $F_{1,35}=0.38$, $p=0.543$; interaction $F_{1,35}=0.60$, $p=0.443$) (data not shown). Overall, chronic treatment with EE-LNG and co-exposure with WIN did not affect anxiety-like behavior in female rats.

Regarding depressive-like behavior in the FST, two-way ANOVA for immobility (Fig. 4C), revealed a significant effect of EE-LNG treatment ($F_{1,35}=5.09$, $p=0.030$) and WIN treatment ($F_{1,35}=5.59$, $p=0.024$), but no interaction ($F_{1,35}=0.42$, $p=0.522$). A trend for increased immobility was found following both EE-LNG and WIN treatments, although no significant changes vs. the respective controls were observed (Fig. 4C). For swimming (Fig. 4D), we found a significant effect of WIN treatment ($F_{1,35}=6.52$, $p=0.015$), but no effect of EE-LNG treatment ($F_{1,35}=3.16$, $p=0.084$) or their interaction ($F_{1,35}=1.31$, $p=0.260$), and no significant changes vs. the respective controls were revealed by the *post hoc* test (Fig. 4D). Finally, no change in climbing behavior (Fig. 4E) was observed among groups (EE-LNG treatment: $F_{1,35}=2.13$, $p=0.154$; WIN treatment: $F_{1,35}=0.31$, $p=0.580$; interaction $F_{1,35}=2.18$, $p=0.149$). Overall, concurrent exposure to HC and cannabinoid agonists did not induce depressive-like behavior in female rats.

3.4 Chronic co-exposure to EE-LNG and WIN does not alter cognitive performance in the novel object recognition and prepulse inhibition tests

All groups displayed similar exploration of the novel object, both for time (Fig. 5A, EE-LNG treatment: $F_{1,35}=0.06$, $p=0.806$; WIN treatment: $F_{1,35}=2.46$, $p=0.126$; interaction $F_{1,35}=2.39$, $p=0.131$) and frequency (Fig. 5B, EE-LNG treatment: $F_{1,35}=1.55$, $p=0.221$; WIN treatment: $F_{1,35}=0.16$, $p=0.695$; interaction $F_{1,35}=0.01$, $p=0.919$) of investigation, suggesting that chronic treatment with EE-LNG and co-exposure with WIN does not affect recognition memory in female rats.

Likewise, all groups displayed a similar percentage of PPI (Fig. 5C, EE-LNG treatment: $F_{1,35}=0.04$, $p=0.841$; WIN treatment: $F_{1,35}=0.75$, $p=0.392$; interaction $F_{1,35}=0.16$, $p=0.692$), suggesting that chronic treatment with EE-LNG and co-exposure with WIN does not affect sensory-motor gating abilities.

3.5 Chronic treatment with EE-LNG, alone and plus WIN, decreases plasma progesterone level

As illustrated in Figure 6, two-way ANOVA for plasma progesterone levels revealed a significant effect of EE-LNG treatment ($F_{1,35}=96.95$, $p<0.0001$), but no effect of WIN treatment ($F_{1,35}=2.75$, $p=0.106$) and no interaction ($F_{1,35}=2.54$, $p=0.119$). As expected (Follesa et al., 2002; Porcu et al., 2012), chronic EE-LNG treatment markedly decreased plasma progesterone concentrations in SAL-treated rats (-85%, $p<0.0001$). Interestingly, WIN treatment *per se* was ineffective, and it did not alter the extent of EE-LNG-induced decrease in plasma progesterone levels (-81%, $p<0.0001$).

4. Discussion

There is a complex interplay between steroid hormones and cannabinoids, with endocannabinoid signaling being able to affect ovarian function in both physiological and pathophysiological states. This study investigated for the first time whether prolonged exposure to a cannabinoid agonist, at a dose regimen known to be rewarding in rats and to

not affect spontaneous motor activity during passive administration sessions (data not shown), may interact with HCs to alter behavior in female rats. Chronic HC administration inhibited ovulation, as shown by the marked suppression in circulating progesterone levels, as expected (Follesa et al., 2002; Porcu et al., 2012). Surprisingly, WIN did not affect either basal or the HC-induced decrease in plasma levels of this hormone. To our knowledge, an effect of WIN administration on plasma progesterone levels has never been reported.

However, while acute THC administration increased plasma progesterone levels in male rats (Grobin et al., 2005), chronic administration of cannabinoid agonists reduces gonadotropin-releasing hormone secretion, ultimately suppressing gonadal progesterone (Gorzalka and Dang, 2012). Thus, WIN, at the low dose used in our experiments, may exert a less potent effect on peripheral progesterone levels.

The effect of chronic EE-LNG treatment and WIN co-exposure was examined on different domains as the prolonged use of both HCs (Concas et al., 2022) and cannabinoid receptor agonists (Howlett and Abood, 2017) are known to affect motor, cognitive and emotional domains. Contrary to our expectations, findings showed that the combination of HCs and low doses of the cannabinoid agonist does not induce significant motor, emotional or cognitive alterations.

The finding that motor activity was not significantly different in animals exposed to EE-LNG is in line with previous evidence showing that HCs do not significantly alter motor parameters in rats (Simone et al., 2015). Likewise, WIN treatment, despite a reduced vertical activity (as compared to saline-treated rats), did not alter horizontal activity and total distance travelled, in agreement with previous studies (Spano et al., 2013). Here, we demonstrated for the first time that the combination of the two pharmacological treatments also does not alter spontaneous motor activity, despite estrogen, progesterone and cannabinoid CB₁ receptors are all widely distributed in rat brain regions regulating motor systems. Importantly, notwithstanding the overall effect of WIN treatment on vertical activity, lack of significant differences in the *post hoc* tests for motor effects of the HC and cannabinoid chronic co-

exposure excludes the possibility that animals' performance in the other behavioral tests might be impaired by nonspecific motor effects.

This study is the first to evaluate the effect of chronic treatment with EE-LNG, low rewarding doses of WIN, or their combinations, on burying behavior in female rats, and findings of no significant changes suggest that a prolonged exposure to HCs and/or a mild chronic stimulation of CB₁ receptors does not trigger obsessive-compulsive-like behavior or neophobia in rats. Yet, since ovarian hormones are known to modulate compulsive behavior in the signal attenuation rat model of obsessive-compulsive disorder (OCD) (Flaisher-Grinberg et al., 2009) and to affect symptom severity in OCD patients (Vulink et al., 2006), we expected to observe a significant effect of EE-LNG in the MB test. In support of our original hypothesis, LNG was found to be effective in contrasting OCD symptoms in a young woman (Perciaccante and Perciaccante, 1993). However, low doses of EE were found to be anxiolytic in the shock-probe defensive burying, while higher doses increased freezing behavior in rats (Simone et al., 2015); moreover, LNG also did not affect burying behavior in female rats (Picazo et al., 1998), which is in line with our finding using EE-LNG, suggesting inconsistent effects of a specific HC treatment on this parameter, likely depending in part on the test used to measure burying behavior. Conversely, lack of significant effects of WIN in the MB test was not totally unexpected, since adolescent exposure to a higher dose of WIN (1.2 mg/kg/day, PND 30-43) was reported not to affect food neophobia in adult male rats (Schoch et al., 2018), while in humans cannabis use seems to ameliorate, rather than trigger, OCD symptoms (Szejko et al., 2020).

In line with data from the MB test, we did not find significant effects of EE-LNG and/or WIN in the EPM test, suggesting that they did not induce an anxiety-like state. With respect to HCs, this result was unexpected since we previously observed anxiety-like behavior in the EPM test following chronic EE-LNG treatment in female rats (Follesa et al., 2002; Porcu et al., 2012). However, those rats were only subjected to the EPM test, thus avoiding repeated manipulations and testing. Further, the different HC dose regimen might also contribute to the discrepant results in the EPM test; while higher EE and LNG doses (0.030 and 0.125

mg, respectively) were anxiogenic (Follesa et al., 2002; Porcu et al., 2012), intermediate doses (0.030 and 0.060 mg, respectively) did not alter behavior (present results), and lower doses (0.010 and 0.020 mg, respectively) induced an anxiolytic effect in the same rat strain (Simone et al., 2015), suggesting that EE and LNG may affect anxiety-like behavior in a dose-dependent manner. Regarding WIN's effect on anxiety-like behavior in rats, there is a degree of inconsistency in the literature. WIN elicited anxiety in the EPM test when acutely administered at doses higher than those used in the present study (Komaki et al., 2015), an effect that was found to be partially more pronounced in pubertal than adult rats (Klugmann et al., 2011). However, other studies reported no effect of WIN (2 mg/kg, i.p.) on anxiety-like behavior in the EPM (Cassar et al., 2022). Importantly, anxiolytic effects were detected in rats that underwent both WIN self-administration and WIN passive (experimenter-given) administration at comparable self-administered doses when tested in the locomotor activity cages, where they constantly spent less time on the box margins (and more at the center) than controls (Spano et al., 2013), which is evocative of a reduced level of anxiety. It is therefore possible that, similarly to the above-mentioned study, WIN treatment in our study might induce anxiolytic effects, although too mild to be detected in the EPM test.

In the FST we observed an increase in immobility time induced by both HCs and WIN treatments, although the *post hoc* test failed to confirm significant differences among groups; moreover, no significant interaction was present, suggesting that there was not an additive effect of the EE-LNG/WIN combination. Increased immobility was paralleled by a concomitant reduction in the time spent swimming, which was significant only for WIN-treated rats, although, once again, the *post hoc* test failed to confirm significant differences among groups. Thus, overall animals that received the concomitant HCs and cannabinoid treatments did not show a depressive-like behavior in the FST. The interpretation of FST behaviors has been challenged, suggesting that increased immobility may be indicative of a passive behavior necessary to cope with the acute swim stress (Molendijk and de Kloet, 2015). Despite the main effect of HC treatment in increasing immobility time, the *post hoc* test results failed to identify significant changes between EE-LNG-treated rats vs. the

respective vehicle-treated controls, suggesting that HCs-treated rats are not depressed, in line with lack of changes in anxiety-like behavior. This interpretation also agrees with lack of changes in anhedonia in the sucrose preference test following chronic EE-LNG treatment (Santorù et al., 2014). In line with our results, female rats exposed during adolescence (PND 35-45) to THC displayed depressive-like behavior in the FST, contrary to male rats that in the same test did not show behavioral despair (Rubino et al., 2008). Yet, no effects of either adolescent cannabis smoke or THC exposure in the FST was also reported (Bruijnzeel et al., 2019), while adulthood treatment with a more potent cannabinoid agonist (HU-210, 0.1 mg/kg), induced an antidepressant-like response in the FST (Morrish et al., 2009). Prolonged exposure to EE-LNG in combination with WIN did not significantly affect animals' performance in the NOR test, suggesting that they did not induce working memory deficits. Likewise, treatment with EE-LNG alone (i.e., plus saline) also had no effect, in line with our previous observation that chronic EE-LNG treatment did not affect recognition memory in the NOR and novel place location tests (Boi et al., 2022). Effects of EE-LNG on recognition memory in rats might be dose-dependent as lower doses (10-20 µg) were found to impair, while a higher EE dose (30 µg) improved performance in the NOR test (Simone et al., 2015). These findings are not surprising as HCs' effects on cognition vary in both women and animals depending on the cognitive domains examined, length of treatment, doses and type of HC administered (Concas et al., 2022). Similarly, chronic treatment with the cannabinoid agonist did not induce recognition memory deficits, in line with our previous observation (Spano et al., 2010), but in contrast with other studies that observed persistent object recognition deficits after chronic pubertal WIN treatment (Schneider et al., 2008). Yet, the different experimental design may explain such a discrepancy, including the higher WIN dose (1.2 mg/kg administered intraperitoneally once/daily vs. a cumulative dose of 0.3 mg/kg achieved after 24 intravenous infusions of WIN 12.5 µg/kg over a 2-h daily session), and the earlier age window for WIN treatment (PND 40-65 vs. PND 56-95). Actually, the age at which WIN exposure occurs appears particularly critical, given that when the same WIN

schedule is administered to adult rats (PND>70), chronic cannabinoid exposure does not alter behavior (Schneider and Koch, 2003).

Finally, we showed for the first time that chronic EE-LNG treatment, alone and in combination with chronic WIN, does not alter PPI in female rats, suggesting that HCs do not induce deficits in the filtering or gating of sensory and cognitive information. Our findings are in line with previous studies showing no change in PPI in rats with different levels of ovarian hormones (i.e., intact, ovariectomized, and ovariectomized rats treated with 17β -estradiol) (Vaillancourt et al., 2002) or exposed to a similar cannabinoid chronic treatment (Spano et al., 2010). However, they contrast with previous studies showing that estrogenic compounds are able to affect baseline PPI in rats (Sbisa et al., 2018). Indeed, human studies reported discrepant findings on HCs' effects on the prepulse inhibition and startle responsiveness, with a recent study describing HC-related PPI differences (Naysmith et al., 2022), while Gogos (2013) did not observe significant PPI differences between HC and non-HC users. To complicate the issue, Beck et al. (2008) reported facilitated fine motor coordination and startle responsiveness in women taking HC, Holloway et al. (2011) found that PPI was lower in young HC users depending on stimulus intensity, and Borgström et al. (2008) reported lower PPI in HC users with negative mood in comparison to users without negative mood, positing that PPI differences may result from HC-related negative affect, rather than HC use *per se*. The finding that WIN does not impair sensorimotor gating in rats is further supported by a study showing that acute, subchronic, and chronic treatment with WIN did not affect PPI in rats (Bortolato et al., 2005), although discrepant findings were also reported after prepubertal and pubertal prolonged exposure to higher WIN doses, i.e., 1.2 vs. 0.3 mg/kg (Schneider and Koch, 2003).

We are aware that the current study is not without limitations. Firstly, we cannot rule out the possibility that different estrogen-progestin combinations, doses and route of administration might elicit a different outcome. Likewise, we cannot rule out the possibility that HCs and rewarding doses of cannabinoids might alter behavior in other species, including humans or their effects might be age-dependent, being more pronounced in young adolescents, an age

window that, due to its short duration, is difficult to capture in rats subjected to long-term chronic drug treatments.

5. Conclusions

We showed that co-exposure to HCs and low but rewarding doses of cannabinoids does not significantly alter the motor and cognitive skills and the emotional state. These findings allow deepening the current knowledge on the interaction between cannabinoid agonists and HCs. Studies on the effects of HCs on behavior contribute to our understanding of their effects on the central nervous system and the implications for women's health.

Acknowledgements

We thank the CeSASst personnel at University of Cagliari for their valuable assistance with animal housing and care.

References

- Beck, K.D., McLaughlin, J., Bergen, M.T., Cominski, T.P., Moldow, R.L., Servatius, R.J., 2008. Facilitated acquisition of the classically conditioned eyeblink response in women taking oral contraceptives. *Behav. Pharmacol.* 19(8), 821-828.
<https://doi.org/10.1097/FBP.0b013e32831c3b82>
- Behrendt, S., Wittchen, H.U., Höfler, M., Lieb, R., Beesdo, K., 2009. Transitions from first substance use to substance use disorders in adolescence: is early onset associated with a rapid escalation? *Drug Alcohol Depend.* 99(1-3), 68-78.
<https://doi.org/10.1016/j.drugalcdep.2008.06.014>
- Belelli, D., Lambert, J.J., 2005. Neurosteroids: endogenous regulators of the GABA(A) receptor. *Nat. Rev. Neurosci.* 6(7), 565-575. <https://doi.org/10.1038/nrn1703>
- Boero, G., Porcu, P., Morrow, A.L., 2020. Pleiotropic actions of allopregnanolone underlie therapeutic benefits in stress-related disease. *Neurobiol. Stress.* 12, 100203.
<https://doi.org/10.1016/j.ynstr.2019.100203>
- Boi, L., Petralla, S., Monti, B., Talani, G., Sanna, E., Pisu, M.G., Calderisi, G., Maciocco, E., Serra, M., Concas, A., Porcu, P., 2022. Chronic treatment with hormonal contraceptives alters hippocampal BDNF and histone H3 post-translational modifications but not learning and memory in female rats. *Horm. Behav.* 144, 105218.
<https://doi.org/10.1016/j.yhbeh.2022.105218>
- Borgström, A., Kask, K., Gulinello, M., Odland, V., Sundström-Poromaa, I., 2008. Patients with adverse mood effects from combined oral contraceptives have lower levels of prepulse inhibition than healthy controls. *Psychoneuroendocrinology.* 33(4), 487-496.
<https://doi.org/10.1016/j.psyneuen.2008.01.007>
- Bortolato, M., Aru, G.N., Frau, R., Orrù, M., Luckey, G.C., Boi, G., Gessa, G.L., 2005. The CB receptor agonist WIN 55,212-2 fails to elicit disruption of prepulse inhibition of the

- startle in Sprague-Dawley rats. *Psychopharmacology (Berl)*. 177(3), 264-271.
<https://doi.org/10.1007/s00213-004-1941-4>
- Bruijnzeel, A.W., Knight, P., Panunzio, S., Xue, S., Bruner, M.M., Wall, S.C., Pompilus, M., Febo, M., Setlow, B., 2019. Effects in rats of adolescent exposure to cannabis smoke or THC on emotional behavior and cognitive function in adulthood. *Psychopharmacology (Berl)*. 236(9), 2773-2784. <https://doi.org/10.1007/s00213-019-05255-7>
- Cahill, L., 2018. How does hormonal contraception affect the developing human adolescent brain? *Curr. Opin. Behav. Sci.* 23, 131-135.
<https://doi.org/10.1016/j.cobeha.2018.06.015>
- Cassar, D., Radic, M., Casarrubea, M., Crunelli, V., Di Giovanni, G., 2022. The effect of cannabinoid receptor agonist WIN 55,212-2 on anxiety-like behavior and locomotion in a genetic model of absence seizures in the elevated plus-maze. *CNS Neurosci. Ther.* 28(8), 1268-1270. <https://doi.org/10.1111/cns.13848>
- Castelli, M.P., Fadda, P., Casu, A., Spano, M.S., Casti, A., Fratta, W., Fattore, L., 2014. Male and female rats differ in brain cannabinoid CB1 receptor density and function and in behavioural traits predisposing to drug addiction: effect of ovarian hormones. *Curr. Pharm. Des.* 20(13), 2100-2113. <https://doi.org/10.2174/13816128113199990430>
- Concas, A., Serra, M., Porcu, P., 2022. How hormonal contraceptives shape brain and behavior: A review of preclinical studies. *Front. Neuroendocrinol.* 66, 101017.
<https://doi.org/10.1016/j.yfrne.2022.101017>
- Concas, A., Sogliano, C., Porcu, P., Marra, C., Brundu, A., Biggio, G., 2006. Neurosteroids in nicotine and morphine dependence. *Psychopharmacology*. 186(3), 281-292.
<https://doi.org/10.1007/s00213-005-0111-7>
- Cooper, D.B., Patel, P., Mahdy, H., 2022. Oral contraceptive pills. In *StatPearls*. StatPearls Publishing, Treasure Island (FL). <https://doi.org/NBK430882>

- Cooper, Z.D., Haney, M., 2014. Investigation of sex-dependent effects of cannabis in daily cannabis smokers. *Drug Alcohol Depend.* 136, 85-91. <https://doi.org/https://doi.org/10.1016/j.drugalcdep.2013.12.013>
- Cryan, J.F., Markou, A., Lucki, I., 2002. Assessing antidepressant activity in rodents: recent developments and future needs. *Trends Pharmacol. Sci.* 23(5), 238-245. [https://doi.org/10.1016/s0165-6147\(02\)02017-5](https://doi.org/10.1016/s0165-6147(02)02017-5)
- Fadda, P., Scherma, M., Spano, M.S., Salis, P., Melis, V., Fattore, L., Fratta, W., 2006. Cannabinoid self-administration increases dopamine release in the nucleus accumbens. *Neuroreport.* 17(15), 1629-1632. <https://doi.org/10.1097/01.wnr.0000236853.40221.8e>
- Fattore, L., Fadda, P., Zanda, M.T., Fratta, W., 2021. Analysis of opioid-seeking behavior through the intravenous self-administration reinstatement model in rats. *Methods Mol. Biol.* 2201, 231-245. https://doi.org/10.1007/978-1-0716-0884-5_21
- Fattore, L., Spano, M.S., Altea, S., Angius, F., Fadda, P., Fratta, W., 2007. Cannabinoid self-administration in rats: sex differences and the influence of ovarian function. *Br. J. Pharmacol.* 152(5), 795-804. <https://doi.org/10.1038/sj.bjp.0707465>
- Fattore, L., Spano, M.S., Altea, S., Fadda, P., Fratta, W., 2010. Drug- and cue-induced reinstatement of cannabinoid-seeking behaviour in male and female rats: influence of ovarian hormones. *Br. J. Pharmacol.* 160(3), 724-735. <https://doi.org/10.1111/j.1476-5381.2010.00734.x>
- Finn, D.A., Phillips, T.J., Okorn, D.M., Chester, J.A., Cunningham, C.L., 1997. Rewarding effect of the neuroactive steroid 3 alpha-hydroxy-5 alpha-pregnan-20-one in mice. *Pharmacol. Biochem. Behav.* 56(2), 261-264. [https://doi.org/10.1016/s0091-3057\(96\)00218-3](https://doi.org/10.1016/s0091-3057(96)00218-3)
- Flaisher-Grinberg, S., Albelda, N., Gitter, L., Weltman, K., Arad, M., Joel, D., 2009. Ovarian hormones modulate 'compulsive' lever-pressing in female rats. *Horm. Behav.* 55(2), 356-365. <https://doi.org/10.1016/j.yhbeh.2008.10.002>

- Follesa, P., Porcu, P., Sogliano, C., Cinus, M., Biggio, F., Mancuso, L., Mostallino, M.C., Paoletti, A.M., Purdy, R.H., Biggio, G., Concas, A., 2002. Changes in GABA(A) receptor gamma2 subunit gene expression induced by long-term administration of oral contraceptives in rats. *Neuropharmacology*. 42(3), 325-336.
[https://doi.org/10.1016/S0028-3908\(01\)00187-3](https://doi.org/10.1016/S0028-3908(01)00187-3)
- Geyer, M.A., Swerdlow, N.R., 2001. Measurement of startle response, prepulse inhibition, and habituation. *Curr. Protoc. Neurosci.* Chapter 8, Unit 8.7.
<https://doi.org/10.1002/0471142301.ns0807s03>
- Gogos, A., 2013. Natural and synthetic sex hormones: effects on higher-order cognitive function and prepulse inhibition. *Biol. Psychol.* 93(1), 17-23.
<https://doi.org/10.1016/j.biopsycho.2013.02.001>
- Gorzalka, B.B., Dang, S.S., 2012. Minireview. Endocannabinoids and gonadal hormones: bidirectional interactions in physiology and behavior. *Endocrinology*. 153(3), 1016-1024. <https://doi.org/10.1210/en.2011-1643>
- Grobin, A.C., VanDoren, M.J., Porrino, L.J., Morrow, A.L., 2005. Cortical 3 alpha-hydroxy-5 alpha-pregnan-20-one levels after acute administration of Delta 9-tetrahydrocannabinol, cocaine and morphine. *Psychopharmacology (Berl)*. 179(3), 544-550. <https://doi.org/10.1007/s00213-004-2084-3>
- Hengartner, M.P., Angst, J., Ajdacic-Gross, V., Rössler, W., 2020. Cannabis use during adolescence and the occurrence of depression, suicidality and anxiety disorder across adulthood: Findings from a longitudinal cohort study over 30 years. *J. Affect. Disord.* 272, 98-103. <https://doi.org/10.1016/j.jad.2020.03.126>
- Hilderbrand, E.R., Lasek, A.W., 2018. Estradiol enhances ethanol reward in female mice through activation of ER α and ER β . *Horm. Behav.* 98, 159-164.
<https://doi.org/10.1016/j.yhbeh.2018.01.001>
- Holloway, J.L., Beck, K.D., Servatius, R.J., 2011. Facilitated acquisition of the classically conditioned eyeblink response in females is augmented in those taking oral

- contraceptives. *Behav. Brain Res.* 216(1), 301-307.
<https://doi.org/10.1016/j.bbr.2010.08.008>
- Howlett, A.C., Abood, M.E., 2017. CB₁ and CB₂ receptor pharmacology. *Adv. Pharmacol.*, 80, 169-206. <https://doi.org/10.1016/bs.apha.2017.03.007>
- Jimenez-Gomez, C., Osentoski, A., Woods, J.H., 2011. Pharmacological evaluation of the adequacy of marble burying as an animal model of compulsion and/or anxiety. *Behav. Pharmacol.* 22(7), 711-713. <https://doi.org/10.1097/FBP.0b013e32834afebe>
- Klugmann, M., Klippenstein, V., Leweke, F.M., Spanagel, R., Schneider, M., 2011. Cannabinoid exposure in pubertal rats increases spontaneous ethanol consumption and NMDA receptor associated protein levels. *Int. J. Neuropsychopharmacol.* 14(4), 505-517. <https://doi.org/10.1017/S1461145710001562>
- Komaki, A., Hashemi-Firouzi, N., Shojaei, S., Souri, Z., Heidari, S., Shahidi, S., 2015. Study the effect of endocannabinoid system on rat behavior in elevated plus-maze. *Basic Clin. Neurosci.* 6(3), 147-153. <https://pubmed.ncbi.nlm.nih.gov/26904171/>
- Levin, E.R., Vitek, W.S., Hammes, S.R., 2018. Estrogens, progestins, and the female reproductive tract. In Brunton, L.L., Hilal-Dandan, R., & Knollmann, B.C. (Eds.), *Goodman & Gilman's The Pharmacological Basis of Therapeutics, 13th edition*. pp, 803-831. McGraw Hill.
- McEwen, B.S., 1994. Steroid hormone actions on the brain: when is the genome involved? *Horm. Behav.* 28(4), 396-405. <https://doi.org/10.1006/hbeh.1994.1036>
- Molendijk, M.L., de Kloet, E.R., 2015. Immobility in the forced swim test is adaptive and does not reflect depression. *Psychoneuroendocrinology*, 62, 389-391.
<https://doi.org/10.1016/j.psyneuen.2015.08.028>
- Morrish, A.C., Hill, M.N., Riebe, C.J., Gorzalka, B.B., 2009. Protracted cannabinoid administration elicits antidepressant behavioral responses in rats: role of gender and noradrenergic transmission. *Physiol. Behav.* 98(1-2), 118-124.
<https://doi.org/10.1016/j.physbeh.2009.04.023>

- Morrow, A.L., Boero, G., Porcu, P., 2020. A rationale for allopregnanolone treatment of alcohol use disorders: Basic and clinical studies. *Alcohol. Clin. Exp. Res.* 44(2), 320-339. <https://doi.org/10.1111/acer.14253>
- Motzo, C., Porceddu, M.L., Maira, G., Flore, G., Concas, A., Dazzi, L., Biggio, G., 1996. Inhibition of basal and stress-induced dopamine release in the cerebral cortex and nucleus accumbens of freely moving rats by the neurosteroid allopregnanolone. *J. Psychopharmacol.* 10(4), 266-272. <https://doi.org/10.1177/026988119601000402>
- Naysmith, L.F., Williams, S.C.R., Kumari, V., 2022. The influence of stimulus onset asynchrony, task order, sex and hormonal contraception on prepulse inhibition and prepulse facilitation: Methodological considerations for drug and imaging research. *J. Psychopharmacol.* 36(11), 1234-1242. <https://doi.org/10.1177/02698811221133469>
- Orsini, C.A., Blaes, S.L., Hernandez, C.M., Betzhold, S.M., Perera, H., Wheeler, A.R., Ten Eyck, T.W., Garman, T.S., Bizon, J.L., Setlow, B., 2021. Regulation of risky decision making by gonadal hormones in males and females. *Neuropsychopharmacology.* 46(3), 603-613. <https://doi.org/10.1038/s41386-020-00827-0>
- Paul, S.M., Pinna, G., Guidotti, A., 2020. Allopregnanolone: From molecular pathophysiology to therapeutics. A historical perspective. *Neurobiol. Stress.* 12, 100215. <https://doi.org/10.1016/j.ynstr.2020.100215>
- Perciaccante, M., Perciaccante, R.G., 1993. Progestin treatment for obsessive-compulsive disorder. *Psychosomatics.* 34(3), 284-285. [https://doi.org/10.1016/S0033-3182\(93\)71898-8](https://doi.org/10.1016/S0033-3182(93)71898-8)
- Petersen, N., Beltz, A.M., Casto, K.V., Taylor, C.M., Jacobs, E.G., Sundström-Poromaa, I., Pletzer, B., 2023. Towards a more comprehensive neuroscience of hormonal contraceptives. *Nat. Neurosci.* 26(4), 529-531. <https://doi.org/10.1038/s41593-023-01273-z>

- Picazo, O., Fernández-Guasti, A., Lemus, A.E., García, G.A., 1998. A-ring reduced derivatives of two synthetic progestins induce anxiolytic effects in ovariectomized rats. *Brain Res.* 796(1-2), 45-52. [https://doi.org/10.1016/s0006-8993\(98\)00314-x](https://doi.org/10.1016/s0006-8993(98)00314-x)
- Pope, H.G., Gruber, A.J., Hudson, J.I., Cohane, G., Huestis, M.A., Yurgelun-Todd, D., 2003. Early-onset cannabis use and cognitive deficits: what is the nature of the association? *Drug Alcohol Depend.* 69(3), 303-310. [https://doi.org/10.1016/s0376-8716\(02\)00334-4](https://doi.org/10.1016/s0376-8716(02)00334-4)
- Porcu, P., Barron, A.M., Frye, C.A., Walf, A.A., Yang, S.Y., He, X.Y., Morrow, A.L., Panzica, G.C., Melcangi, R.C., 2016. Neurosteroidogenesis today: Novel targets for neuroactive steroid synthesis and action and their relevance for translational research. *J. Neuroendocrinol.* 28(2), 12351. <https://doi.org/10.1111/jne.12351>
- Porcu, P., Mostallino, M.C., Sogliano, C., Santoru, F., Berretti, R., Concas, A., 2012. Long-term administration with levonorgestrel decreases allopregnanolone levels and alters GABA(A) receptor subunit expression and anxiety-like behavior. *Pharmacol. Biochem. Behav.* 102(2), 366-372. <https://doi.org/10.1016/j.pbb.2012.05.011>
- Porcu, P., Serra, M., Concas, A., 2019. The brain as a target of hormonal contraceptives: Evidence from animal studies. *Front. Neuroendocrinol.* 55, 100799. <https://doi.org/10.1016/j.yfrne.2019.100799>
- Porsolt, R.D., Le Pichon, M., Jalfre, M. 1977a. Depression: a new animal model sensitive to antidepressant treatments. *Nature* 266(5604), 730-732. <https://doi.org/10.1038/266730a0>
- Porsolt, R.D., Bertin, A., Jalfre, M., 1977b. Behavioral despair in mice: a primary screening test for antidepressants. *Arch. Int. Pharmacodyn. Ther.* 229(2), 327-336. <https://pubmed.ncbi.nlm.nih.gov/596982/>
- Quigley, J.A., Logsdon, M.K., Turner, C.A., Gonzalez, I.L., Leonardo, N.B., Becker, J.B., 2021. Sex differences in vulnerability to addiction. *Neuropharmacology.* 187, 108491. <https://doi.org/10.1016/j.neuropharm.2021.108491>

- Rougé-Pont, F., Mayo, W., Marinelli, M., Gingras, M., Le Moal, M., Piazza, P.V., 2002. The neurosteroid allopregnanolone increases dopamine release and dopaminergic response to morphine in the rat nucleus accumbens. *Eur. J. Neurosci.* 16(1), 169-173. <https://doi.org/10.1046/j.1460-9568.2002.02084.x>
- Rubino, T., Vigano', D., Realini, N., Guidali, C., Braidà, D., Capurro, V., Castiglioni, C., Cherubino, F., Romualdi, P., Candeletti, S., Sala, M., Parolaro, D., 2008. Chronic delta 9-tetrahydrocannabinol during adolescence provokes sex-dependent changes in the emotional profile in adult rats: behavioral and biochemical correlates. *Neuropsychopharmacology.* 33(11), 2760-2771. <https://doi.org/10.1038/sj.npp.1301664>
- Rupprecht, R., 2003. Neuroactive steroids: mechanisms of action and neuropsychopharmacological properties. *Psychoneuroendocrinology.* 28(2), 139-168. [https://doi.org/10.1016/s0306-4530\(02\)00064-1](https://doi.org/10.1016/s0306-4530(02)00064-1)
- Santorù, F., Berretti, R., Locci, A., Porcu, P., Concas, A., 2014. Decreased allopregnanolone induced by hormonal contraceptives is associated with a reduction in social behavior and sexual motivation in female rats. *Psychopharmacology.* 231(17), 3351-3364. <https://doi.org/10.1007/s00213-014-3539-9>
- Sbisa, A., van den Buuse, M., Gogos, A., 2018. The effect of estrogenic compounds on psychosis-like behaviour in female rats. *PLoS One.* 13(3), e0193853. <https://doi.org/10.1371/journal.pone.0193853>
- Schneider, M., Koch, M., 2003. Chronic pubertal, but not adult chronic cannabinoid treatment impairs sensorimotor gating, recognition memory, and the performance in a progressive ratio task in adult rats. *Neuropsychopharmacology.* 28(10), 1760-1769. <https://doi.org/10.1038/sj.npp.1300225>
- Schneider, M., Schömig, E., Leweke, F.M., 2008. Acute and chronic cannabinoid treatment differentially affects recognition memory and social behavior in pubertal and adult rats. *Addict. Biol.* 13(3-4), 345-357. <https://doi.org/10.1111/j.1369-1600.2008.00117.x>

- Schoch, H., Huerta, M.Y., Ruiz, C.M., Farrell, M.R., Jung, K.M., Huang, J.J., Campbell, R.R., Piomelli, D., Mahler, S.V., 2018. Adolescent cannabinoid exposure effects on natural reward seeking and learning in rats. *Psychopharmacology (Berl)*. 235(1), 121-134.
<https://doi.org/10.1007/s00213-017-4749-8>
- Simone, J., Bogue, E.A., Bhatti, D.L., Day, L.E., Farr, N.A., Grossman, A.M., Holmes, P.V., 2015. Ethinyl estradiol and levonorgestrel alter cognition and anxiety in rats concurrent with a decrease in tyrosine hydroxylase expression in the locus coeruleus and brain-derived neurotrophic factor expression in the hippocampus. *Psychoneuroendocrinology*. 62, 265-278.
<https://doi.org/10.1016/j.psyneuen.2015.08.015>
- Sinnott, R.S., Mark, G.P., Finn, D.A., 2002. Reinforcing effects of the neurosteroid allopregnanolone in rats. *Pharmacol. Biochem. Behav.* 72(4), 923-929.
[https://doi.org/10.1016/s0091-3057\(02\)00776-1](https://doi.org/10.1016/s0091-3057(02)00776-1)
- Spano, M.S., Fadda, P., Frau, R., Fattore, L., Fratta, W., 2010. Cannabinoid self-administration attenuates PCP-induced schizophrenia-like symptoms in adult rats. *Eur. Neuropsychopharmacol.* 20(1), 25-36. <https://doi.org/10.1016/j.euroneuro.2009.09.004>
- Spano, M.S., Fattore, L., Cadeddu, F., Fratta, W., Fadda, P., 2013. Chronic cannabinoid exposure reduces phencyclidine-induced schizophrenia-like positive symptoms in adult rats. *Psychopharmacology (Berl)*. 225(3), 531-542. <https://doi.org/10.1007/s00213-012-2839-1>
- Struik, D., Sanna, F., Fattore, L., 2018. The modulating role of sex and anabolic-androgenic steroid hormones in cannabinoid sensitivity. *Front. Behav. Neurosci.* 12, 249.
<https://doi.org/10.3389/fnbeh.2018.00249>
- Szejko, N., Fremer, C., Müller-Vahl, K.R., 2020. Cannabis improves obsessive-compulsive disorder-case report and review of the literature. *Front. Psychiatry*. 11, 681.
<https://doi.org/10.3389/fpsyt.2020.00681>

United Nations, 2019. Contraceptive Use by Method 2019: Data Booklet

(ST/ESA/SER.A/435). Department of Economic and Social Affairs, Population Division
https://www.un.org/development/desa/pd/sites/www.un.org.development.desa.pd/files/files/documents/2020/Jan/un_2019_contraceptiveusebymethod_databooklet.pdf
(accessed 09 October 2023)

UNODC, United Nations Office on Drugs and Crime, World Drug Report 2022 (United Nations publication, 2022). <https://www.unodc.org/unodc/en/data-and-analysis/world-drug-report-2022.html> (accessed 09 October 2023).

Vaillancourt, C., Cyr, M., Rochford, J., Boksa, P., Di Paolo, T., 2002. Effects of ovariectomy and estradiol on acoustic startle responses in rats. *Pharmacol. Biochem. Behav.* 74(1), 103-109. [https://doi.org/10.1016/s0091-3057\(02\)00967-x](https://doi.org/10.1016/s0091-3057(02)00967-x)

Vallée, M., Vitiello, S., Bellocchio, L., Hébert-Chatelain, E., Monlezun, S., Martin-Garcia, E., Kasanetz, F., Baillie, G.L., Panin, F., Cathala, A., Roullot-Lacarrière, V., Fabre, S., Hurst, D.P., Lynch, D.L., Shore, D.M., Deroche-Gamonet, V., Spampinato, U., Revest, J.-M., Maldonado, R., Reggio, P.H., Ross, R.A., Marsicano, G., Piazza, P.V., 2014. Pregnenolone can protect the brain from cannabis intoxication. *Science*. 343(6166), 94-98. <https://doi.org/10.1126/science.1243985>

Vetter-O'Hagen, C.S., Spear, L.P., 2012. Hormonal and physical markers of puberty and their relationship to adolescent-typical novelty-directed behavior. *Dev. Psychobiol.* 54(5), 523-535. <https://doi.org/10.1002/dev.20610>

Vulink, N.C., Denys, D., Bus, L., Westenberg, H.G., 2006. Female hormones affect symptom severity in obsessive-compulsive disorder. *Int. Clin. Psychopharmacol.* 21(3), 171-175. <https://doi.org/10.1097/01.yic.0000199454.62423.99>

Yoest, K.E., Cummings, J.A., Becker, J.B., 2014. Estradiol, dopamine and motivation. *Cent. Nerv. Syst. Agents Med. Chem.* 14(2), 83-89.
<https://doi.org/10.2174/1871524914666141226103135>

Zanda, M.T., Fadda, P., Antinori, S., Di Chio, M., Fratta, W., Chiamulera, C., Fattore, L.,
2017. Methoxetamine affects brain processing involved in emotional response in rats.
Br. J. Pharmacol. 174(19), 3333-3345. <https://doi.org/10.1111/bph.13952>

Author Contributions Statement

LF, AC and PP: Conceptualization; LF, AP, LC, CC, CS, PP: Investigation; LF, AP, MGP, PP: Data curation and formal analysis; MS: Resources; LF, PP: Funding acquisition, Project administration, Supervision and Writing - original draft; LF, AP, LC, CC, CS, MGP, MS, AC, PP: Writing - review & editing.

Author Confirmation Statement

The manuscript has been read and approved by all authors.

Author Declaration of Interest Statement

Declaration of interest: none.

Funding Information

This work was supported by Fondazione di Sardegna, Italy (Grant IDs 2019.1440 and 2021.0504 to PP and 2020.1072 to LF) and NeuroSardynia ONLUS, Italy.

Figures and Figure Legends

Figure 1. Timeline of the experimental procedures. Upon arrival at postnatal day (PND) 32-35, rats were allowed to acclimate before undergoing surgery for catheter insertion into the jugular vein. Starting on PND 52 they received daily subcutaneous injections of the EE-LNG combination (0.020 and 0.060 mg, respectively). On PND 56 they started daily infusions of WIN 55,212-2 (12.5 μ g/kg). Behavioral tests took place between PND 84 and PND 95 in the following order: locomotor activity (LA, PND 84), novel object recognition (NOR, PND 86), forced swimming (FST, PND 88), marble burying (MB, PND 91), elevated plus maze (EPM, PND 93), and prepulse inhibition (PPI, PND 95). Rats were sacrificed on PND 98 to collect plasma samples for determination of progesterone levels.




Figure 2. Effect of long-term EE-LNG treatment and co-exposure to WIN 55,212-2 on locomotor activity. Spontaneous locomotor activity was assessed on PND 84 and horizontal activity (A), vertical activity (B), and total distance travelled (C), were collected every 10 minutes over the 60 minutes test. Horizontal and vertical activities are expressed as mean counts of photobeam interruptions; distance travelled is in cm. Data represent the mean $\pm$ SEM of values from 9 (Vehicle + Saline), 11 (Vehicle + WIN), 8 (EE-LNG + Saline) or 11 (EE-LNG + WIN) rats/group. Symbols show significant changes ($p < 0.05$) vs. the respective 10 minutes value for every experimental group as follows: *Vehicle + Saline, #Vehicle + WIN 55,212-2, §EE-LNG + Saline, ^EE-LNG + WIN 55,212-2; repeated measures ANOVA followed by Tukey's *post hoc* test. Exact p values are shown in Supplementary Table 2.




Figure 3. Effect of long-term EE-LNG treatment and co-exposure to WIN 55,212-2 in the marble burying test. Neophobia/compulsive behavior was assessed in the marble burying test on PND 91. Data are reported as number of marbles fully (>95%, A) and partially (>67%, B) buried. Panel C illustrates the total number of marbles that triggered burying behavior (fully + partially buried). Each bar represents the mean $\pm$ SEM of values from 9 (Vehicle + Saline), 11 (Vehicle + WIN), 8 (EE-LNG + Saline) or 11 (EE-LNG + WIN) rats/group.




Figure 4. Effect of long-term EE-LNG treatment and co-exposure to WIN 55,212-2 on anxiety- and depressive-like behavior. Anxiety-like behavior was assessed in the elevated plus maze test on PND 93. Data are reported as number of entries (A) and time spent (B) in the open arms of the maze. Depressive-like behavior was assessed in the forced swimming test on PND 88. Data are reported as time (seconds) spent in immobility (C), swimming (D) and climbing (E) behavior. Each bar represents the mean $\pm$ SEM of values from 9 (Vehicle + Saline), 11 (Vehicle + WIN), 8 (EE-LNG + Saline) or 11 (EE-LNG + WIN) rats/group.




Figure 5. Effect of long-term EE-LNG treatment and co-exposure to WIN 55,212-2 on cognitive performance in the novel object recognition and prepulse inhibition tests.

Recognition memory was assessed in the novel object recognition (NOR) test on PND 86.

Data are reported as percent time (A) and percent frequency (B) of exploration of the novel vs. familiar object. Prepulse inhibition (PPI) was assessed on PND 95 and data are reported as percent PPI (C). Each bar represents the mean $\pm$ SEM of values from 9 (Vehicle + Saline), 11 (Vehicle + WIN), 8 (EE-LNG + Saline) or 11 (EE-LNG + WIN) rats/group.





Figure 6. Effect of long-term EE-LNG treatment and co-exposure to WIN 55,212-2 on plasma progesterone levels. Progesterone levels were measured in plasma samples collected on PND 98 at the end of all behavioral tests. Data are expressed as ng/ml and each bar represents the mean $\pm$ SEM of values from 9 (Vehicle + Saline), 11 (Vehicle + WIN), 8 (EE-LNG + Saline) or 11 (EE-LNG + WIN) rats/group. * $p < 0.0001$, two-way ANOVA followed by Tukey's *post hoc* test.



Supplementary Table 1.

Detailed results for the two-way ANOVA repeated measures for each parameter assessed in the locomotor activity test.

Effect	Horizontal activity	Vertical activity	Total distance travelled
EE-LNG treatment	$F_{1,35}=1.19$; $p=0.281$	$F_{1,35}=0.53$; $p=0.470$	$F_{1,35}=1.65$; $p=0.208$
WIN treatment	$F_{1,35}=1.92$; $p=0.174$	$F_{1,35}=6.56$; $p=0.015$	$F_{1,35}=3.62$; $p=0.065$
EE-LNG x WIN	$F_{1,35}=0.06$; $p=0.812$	$F_{1,35}=0.14$; $p=0.708$	$F_{1,35}=0.07$; $p=0.800$
Time	$F_{5,175}=99.96$; $p<0.00001$	$F_{5,175}=41.90$; $p<0.00001$	$F_{5,175}=99.43$; $p<0.00001$
Time x EE-LNG	$F_{5,175}=2.11$; $p=0.067$	$F_{5,175}=1.61$; $p=0.160$	$F_{5,175}=1.75$; $p=0.126$
Time x WIN	$F_{5,175}=1.92$; $p=0.093$	$F_{5,175}=3.53$; $p=0.005$	$F_{5,175}=4.25$; $p=0.001$
Time x EE-LNG x WIN	$F_{5,175}=1.24$; $p=0.292$	$F_{5,175}=1.74$; $p=0.127$	$F_{5,175}=0.84$; $p=0.524$

Supplementary Table 2.

Exact p values for the Tukey *post hoc* analysis for Time effect in the locomotor activity test.

	Vehicle+Saline	EE-LNG+Saline	Vehicle+WIN	EE-LNG+WIN
Horizontal activity				
20 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
30 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
40 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
50 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
60 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
Vertical activity				
20 min	<i>p=0.111</i>	p=0.024	<i>p=0.841</i>	p=0.0004
30 min	p=0.00002	p=0.0008	p=0.009	p=0.0002
40 min	p=0.00002	p=0.0002	<i>p=0.064</i>	p=0.00006
50 min	p=0.00002	p=0.0001	p=0.006	p=0.00002
60 min	p=0.00002	p=0.00002	p=0.025	p=0.00006
Total distance travelled				
20 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
30 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
40 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
50 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002
60 min	p=0.00002	p=0.00002	p=0.00002	p=0.00002

Exact p values refer to significant differences for each time point vs. the respective 10 min value for each experimental group.