

# Targeting tumor associated carbonic anhydrase IX and XII: highly isozyme selective coumarin and psoralen inhibitors

Claudia Melis<sup>a</sup>, Simona Distinto<sup>a\*</sup>, Giulia Bianco<sup>a</sup>, Rita Meleddu<sup>a</sup>, Filippo Cottiglia<sup>a</sup>, Benedetta Fois<sup>a</sup>, Domenico Taverna<sup>b</sup>, Rossella Angius<sup>c</sup>, Stefano Alcaro<sup>d</sup>, Francesco Ortuso<sup>d</sup>, Marco Gaspari<sup>b</sup>, Andrea Angeli<sup>e</sup>, Sonia Del Prete<sup>f</sup>, Clemente Capasso<sup>f</sup>, Claudiu T. Supuran<sup>e\*</sup> and Elias Mac-cioni<sup>a</sup>.

<sup>a</sup> Department of Life and Environmental Sciences, University of Cagliari, Via Ospedale 72, 09124 Cagliari, Italy

<sup>b</sup> Department of Experimental and Clinical Medicine, "Magna Græcia" University of Catanzaro, Catanzaro, Italy

<sup>c</sup> Laboratorio NMR e Tecnologie Bioanalitiche, Sardegna Ricerche, 09010 Pula, Cagliari, Italy

<sup>d</sup> Dipartimento di Scienze della Salute, Università Magna Graecia di Catanzaro, Campus 'S. Venuta', Viale Europa, 88100 Catanzaro, Italy

<sup>e</sup> Dipartimento NEUROFARBA, Sezione di Scienze Farmaceutiche, Università degli Studi di Firenze, Sesto Fiorentino, Florence, Italy

<sup>f</sup> Istituto di Bioscienze e Biorisorse, CNR, Via Pietro Castellino 111, 80131, Napoli, Italy

**KEYWORDS:** *hCA IX, hCA XII, inhibitors, tumor, coumarin, benzenesulfonamide.*

---

**ABSTRACT** A small library of psoralen carboxylic acids and their corresponding benzenesulfonamide derivatives were designed and synthesized to evaluate their activity and selectivity toward tumor associated human carbonic anhydrase (hCA) isoforms IX and XII. Both psoralen acids and sulfonamides exhibited potent inhibition of IX and XII isozymes in the nanomolar concentration range. However, psoralen acids resulted as the most selective in comparison with the corresponding benzenesulfonamide derivatives. Our data indicate that the psoralen scaffold is a promising starting point for the design of highly selective tumor associated hCA inhibitors.

---

Coumarins are a class of heterocyclic compounds widely distributed in nature with a wide variety of biological activity such as antiviral,<sup>1, 2</sup> glucose-lowering agents,<sup>3, 4</sup> anti-mycotic,<sup>5</sup> and antitumor.<sup>6-10</sup>

Calanolides A and B extracted from *Calophyllum lanigerum* showed anti-HIV-1 activity.<sup>11</sup> Furthermore imperatorin has been reported to have anti-inflammatory, antibacterial, antifungal, antiviral, and anticancer activity and might have future clinical application.<sup>12</sup> Noteworthy, other derivatives such as psoralen, bergapten, marmesin, rutaretin are known antitubercular agents.<sup>13</sup> In particular the 7*H*-furo[3,2-*g*]chromen-7-one derivatives, commonly known as psoralens, have been investigated for their ability to suppress proliferation and angiogenesis in human colon cancer cells by targeting HIF-1 $\alpha$  via the mTOR/p70S6K/4E-BP1 and MAPK pathways<sup>7</sup> and to reverse multidrug resistance in both lung cancer A549/D16 and breast cancer MCF-7/ADR cells.<sup>14, 15</sup>

Moreover, the ability of coumarin derivatives to inhibit human carbonic anhydrase isozymes IX and XII (hCA IX and XII) has also been reported.<sup>16-19</sup>

Noteworthy, the chemical accessibility and the ease of functionalization of the coumarin nucleus, allows for synthesizing diverse compound libraries.

The trans-membrane hCA IX and hCA XII isoforms have been associated with tumor progression and invasion.<sup>20-28</sup> In normal conditions, CA IX is expressed in the stomach and few other tissues while it is ectopically induced and highly overexpressed in many hypoxic solid tumors, by a direct transcriptional activation of the *CA9* gene via HIF-1 $\alpha$ . Therefore the design of new and isozyme selective hCA inhibitors is an attractive challenge for medicinal chemists.

In this respect, the design of new psoralen derivatives as hCA IX and XII inhibitors could be advantageous and may lead to the identification of multitarget agents since other cancer related enzymes and pathways could also be hit by such derivatives.<sup>6, 7, 9, 14, 15, 29, 30</sup>


Pursuing on our efforts in the design and synthesis of hCA inhibitors,<sup>31, 32</sup> we have designed and synthesized a small library of coumarin (**EMAC10151 A-D**) and psoralen (**EMAC10152 A-D**) derivatives with the aim to target the hCA IX and XII isoforms. To achieve a deeper insight in

the inhibitory potential of such derivatives we have also synthesized **EMAC10152 A-D** and the corresponding benzenesulfonamide hybrids **EMAC10153 A-D**. The new compounds were synthesized by a versatile multistep synthetic approach (Scheme 1). Methyl 2-(7-hydroxy-4-methyl-2-oxo-2H-chromen-3-yl)acetate **EMAC10150** was obtained by Pechmann condensation of resorcinol with dimethylacetylsuccinate in 98% sulfuric acid. **EMAC10150** was converted into methyl 2-(4-methyl-2-oxo-7-aryloxy-2H-chromen-3-yl)acetate derivatives **EMAC10151 A-D** by Williamson reaction with the appropriate  $\alpha$ -haloketone, in the presence of dry acetone and using potassium carbonate to generate the *in situ* alkoxide ion.

By heating **EMAC10151 A-D** in sodium hydroxide 1 M solution, the simultaneous intramolecular condensation and ester saponification took place to give 2-(5-methyl-7-oxo-3-aryl-7H-furo[3,2-g]chromen-6-yl)acetic acid **EMAC10152 A-D**.

The formation of the 2H-furo[2,3-h]chromen-2-one isomers was not observed, as confirmed by <sup>1</sup>H-NMR spectroscopy (Figure S1 SI). Benzenesulfonamide hybrids **EMAC10153 A-D** were obtained via acyl chloride formation and subsequent reaction with 4-aminobenzenesulfonamide. All compounds were characterized by means of both analytical and spectroscopic methods (Figures S2-S25) and finally evaluated for the inhibition of four hCA isoforms I, II, IX, and XII (Table 1).

### Scheme 1. Synthetic pathway to compounds **EMAC10151 A-D**, **EMAC10152 A-D**, and **EMAC10153 A-D**.



Reagents and conditions (i) dimethylacetylsuccinate, H<sub>2</sub>SO<sub>4</sub> 98%, RT; (ii) dry acetone, K<sub>2</sub>CO<sub>3</sub>,  $\alpha$ -haloketone, reflux, 1-5 h.; (iii) propan-2-ol, NaOH 1 N, reflux, 4 h; (iv) SOCl<sub>2</sub>, RT, dry acetone, 4-aminobenzenesulfonamide, dry pyridine.

**Table 1. Inhibition data towards hCA I, II, IX, and XII of compounds **EMAC10151 A-D**, **EMAC10152 A-D**, and **EMAC10153 A-D**.**

K <sub>i</sub> (nM)					
Compound	R	hCAI	hCAII	hCAIX	hCAXII
10151 A	Cl	>10000	>10000	23.6	446.6
10151 B	CH <sub>3</sub>	>10000	>10000	122.8	56.6
10151 C	H	>10000	>10000	89.7	72.5
10151 D	F	>10000	>10000	84.7	250.0
10152 A	Cl	>10000	>10000	94.7	9.3
10152 B	CH <sub>3</sub>	>10000	>10000	23.0	9.1
10152 C	H	>10000	>10000	17.5	9.4
10152 D	F	>10000	>10000	17.7	7.4
10153 A	Cl	6829.7	55.1	17.8	2.4
10153 B	CH <sub>3</sub>	7069.0	560.0	91.6	3.4
10153 C	H	7016.1	46.6	16.5	3.6
10153 D	F	7148.1	79.5	108.4	49.9
AAZ	//	250	12.1	25.8	5.7

EMAC					
10151 A	Cl	>10000	>10000	23.6	446.6
10151 B	CH <sub>3</sub>	>10000	>10000	122.8	56.6
10151 C	H	>10000	>10000	89.7	72.5
10151 D	F	>10000	>10000	84.7	250.0
10152 A	Cl	>10000	>10000	94.7	9.3
10152 B	CH <sub>3</sub>	>10000	>10000	23.0	9.1
10152 C	H	>10000	>10000	17.5	9.4
10152 D	F	>10000	>10000	17.7	7.4
10153 A	Cl	6829.7	55.1	17.8	2.4
10153 B	CH <sub>3</sub>	7069.0	560.0	91.6	3.4
10153 C	H	7016.1	46.6	16.5	3.6
10153 D	F	7148.1	79.5	108.4	49.9
AAZ	//	250	12.1	25.8	5.7


Most of **EMAC** compounds possess, although with some distinction, selective activity toward the IX and XII isoforms. However, when the benzenesulfonamide moiety is present, a strong reduction of the selectivity becomes evident. In particular, in the case of compound **EMAC10153 D**, the complete loss of selectivity was observed.

On the contrary both **EMAC10151 A-D** and **EMAC10152 A-D** are generally potent and selective inhibitors of the tumor associated hCA IX and XII isoforms. Nevertheless, some considerations should be reported. Regarding the methyl 2-(4-methyl-2-oxo-7-aryloxy-2H-chromen-3-yl)acetate derivatives, **EMAC10151 A-D** a different activity profile could be observed, according to the substitution of the phenyl ring. Thus, the introduction of a halogen atom (Cl or F, **EMAC10151 A** and **EMAC10151 D**) in position 4 of the phenyl substituent, oriented the activity and selectivity toward the IX isozyme. On the contrary, the unsubstituted phenyl or the 4-CH<sub>3</sub>-phenyl moieties led to an increase of the activity toward the hCA XII isoform. Indeed, a reduction of the activity toward hCA IX was observed for compound **EMAC10151 B**. However, although weaker ( $K_i = 122.8$  nM), this derivative exhibited a high selectivity toward the hCA IX and XII isozymes. A similar behavior could be observed for all the psoralen carboxylic acid derivatives **EMAC10152 A-D**. All these compounds were potent and selective tumor associated isoforms IX and XII hCAIs. Their  $K_i$  values toward hCA IX and XII ranged from 94.7 to 7.4 nM, while no inhibition could be observed up to 10000 nM concentration toward the off target hCA I and hCA II isozymes. In more detail, it could be pointed out that, in the case of hCA IX inhibition, the biological activity is more influenced by the size of the substituent on the phenyl ring rather than by its nature. Thus the introduction of a chlorine atom (**EMAC10152 A**) led to the less potent compound within the 10152 series. On the contrary compounds **EMAC10152 C** and **EMAC10152 D**, bearing an unsubstituted phenyl and a fluoro phenyl moiety respectively, resulted as the most potent inhibitors. Interestingly, hCA XII is inhibited at low nanomolar concentrations, regardless of the nature of the substitution.

As mentioned above, in the case of the psoralen-benzenesulfonamide hybrids (**EMAC10153 A-D**), a potent activity toward the hCA XII was observed for most of the tested derivatives. The inhibition of the XII isozyme could still be observed at concentrations ranging from nanomolar to low nanomolar, but a lower selectivity is generally measured toward the off-target isozyme hCA II with respect to **EMAC10151** and **EMAC10152** series. Only in the case of compound **EMAC10153 B**, bearing a methylphenyl moiety on the furan ring, a good selectivity toward hCA IX and XII versus hCA I and II was found.


Computational methods have been applied in order to explain the selectivity of the synthesized coumarin derivatives toward hCA IX and hCA XII isoforms and their inability to inhibit the II isoform, which, in our intent, represents an off-target. To perform our studies we chose the two compounds that showed the highest potency and selectivity toward hCA IX and XII: the acid derivative **EMAC10152 D** and the sulfonamide derivative **EMAC10153 B**. Indeed, these compounds have been docked inside the three crystal structures II (3f8e), IX (5fl4) and XII (4ww8).<sup>33, 34</sup> The previously validated QMPL<sup>35</sup> protocol was applied.<sup>32</sup> Both coumarins derivatives have been prepared considering the possibility that the coumarin moiety can be hydrolyzed by the Zn<sup>2+</sup> activated water molecule of the enzyme cavity, which acts as a very potent nucleophile.<sup>33</sup> Therefore, the compounds were ionized at pH 7.4. However, sulfonamide group was considered both unionized, at pH 7.4, and ionized taking into account the micro basic pH inside the binding pocket (Figure S26 SI). All molecules were subjected to conformational analysis and the global minimum of each was selected for carrying out docking experiments. The putative binding mode suggested by docking simulations well explain the selectivity of our compounds. In fact, both compounds are not able to enter deeply in the active site of hCA II because of the steric hindrance of Phe 131 (Figure 1 and Figure S27). On the contrary, they perfectly fit into the active site of hCA IX and hCA XII, where Phe 131 is substituted by Val 130 and Ala 129, respectively (Figure 1). In particular the suggested binding mode shows that the coumarin moiety of compound **EMAC10152 D** can reach the catalytic site and be hydrolyzed. Conversely, like most of sulfonamide derivatives, **EMAC10153 B** acts by interacting with the sulfonamide moiety, while the coumarin portion, oriented

outside the pocket, cannot be hydrolyzed (Figure 1). According to the above suggested mechanism, only the complexes with hydrolyzed **EMAC10152 D** and closed **EMAC10153 B** were subjected to a post-docking procedure based on energy minimization in order to take into account the induced fit phenomena that occur upon ligand binding.<sup>36</sup> The docking results are in agreement with experimental data regarding the isoform selectivity. In fact, EMAC compounds bound to hCA IX and XII showed lower G-score compared to the complex with -isoform II (Table S1), indicating a major stability of the complexes with hCA IX and XII in agreement with the experimental results (Table 1).



**Figure 1.** Three dimensional representation of docking results of a) **EMAC10152 D** in cyan and b) **EMAC10153 B** in pink with the three hCA isoforms represented as surface: in grey hCA II, in beige hCA IX and in light-blue hCA XII. The active site is highlighted in pale yellow.

The analysis of the interaction mode with isoforms IX and XII (Figure 2) shows that **EMAC10152 D** and **EMAC10153 B** are able to establish a wide array of hydrogen bonds with crucial residues of the binding site and several hydrophobic interactions. In more detail, the carboxylic function of **EMAC10152 D** interacts as a bidentate chelator with Zn<sup>2+</sup>. Conversely, in the case of **EMAC10153 A**, the sulfonamide group plays a key role by interacting with Zn<sup>2+</sup> and Thr residues, as it has been generally observed for this class of compounds.<sup>37, 38</sup>



**Figure 2.** Three dimensional representation of the putative binding mode obtained by docking experiment of EMAC10152 D and EMAC10153 B into a) hCA IX (beige) and b) hCA XII (light blue), and the relative 2D representation of the complexes stabilizing interactions with the residues of the binding site.

Therefore, a putative binding mode seems to confirm the importance of the coumarin moiety in both closed and hydrolyzed conformation, although the different functionalization of EMAC compounds with respect to previously reported coumarins,<sup>33</sup> led to a different target recognition that involves the interaction with the zinc ion. Hence, this could represent an innovative mode of action. In addition, it was confirmed that a bulky moiety, linked to the catalytic site binding scaffold, allows selectivity over the off-target II isoform. In fact, in the case of the II isoform, the steric hindrance of Phe131 prevents the ligand-target interaction.

According to these results both psoralen and the hybrid compounds could be considered as promising scaffolds for the design of high selective inhibitors of the hCAs IX and XII isoforms. However, while very high selectivity was always achieved by coumarin (EMAC10151) and psoralen (EMAC10152) derivatives, the same behavior was not observed in the case of the hybrid compounds EMAC10153. Probably, this is mainly due to the presence of the strong  $\text{Zn}^{2+}$  binder benzenesulfonamide which, on the one hand, may lead to very potent compounds but, on the other, could affect selectivity.

Conversely, hydrolyzed coumarin EMAC10151 and psoralen EMAC10152 most likely act as bidentate  $\text{Zn}^{2+}$  chelators, while hybrid compounds EMAC10153 may preferentially act as traditional  $\text{Zn}^{2+}$ -binders. These results were very encouraging and pushed us to further investigate these derivatives in order to identify potential candidates for the treatment of hypoxic tumors.

## ASSOCIATED CONTENT

### Supporting Information

Experimental procedures and compounds characterization are reported (PDF). The Supporting Information is available free of charge on the ACS Publications website at DOI:

## AUTHOR INFORMATION

### Corresponding Author

claudiu.supuran@unifi.it, s.distinto@unica.it.

### Author Contributions

The authors declare no competing financial interest.

## ABBREVIATIONS

HIF-1 $\alpha$ , Hypoxia-inducible factor 1-alpha, mTOR, mammalian target of rapamycin; p70S6K, Ribosomal protein S6 kinase beta-1; 4E-BP1, Eukaryotic translation initiation factor 4E (eIF4E)-binding protein 1; MAPK, mitogen-activated protein kinase; hCA, human Carbonic anhydrase; AAZ, acetazolamide.

## REFERENCES

- Ong, E. B. B.; Watanabe, N.; Saito, A.; Futamura, Y.; Abd El Galil, K. H.; Koito, A.; Najimudin, N.; Osada, H.; Vipirinin, a coumarin-based HIV-1 VPR inhibitor, interacts with a hydrophobic region of VPR. *J. Biol. Chem.* **2011**, *286* (16), 14049-14056.
- Lin, P.-H.; Ke, Y.-Y.; Su, C.-T.; Shiao, H.-Y.; Hsieh, H.-P.; Chao, Y.-K.; Lee, C.-N.; Kao, C.-L.; Chao, Y.-S.; Chang, S.-Y., Inhibition of HIV-1 tat-mediated transcription by a coumarin derivative, BPRHIV001, through the Akt pathway. *J. Virol.* **2011**, *85* (17), 9114-9126.

3. Han, J.; Sun, L.; Huang, X.; Li, Z.; Zhang, C.; Qian, H.; Huang, W., Novel coumarin modified GLP-1 derivatives with enhanced plasma stability and prolonged in vivo glucose-lowering ability. *Br. J. Pharmacol.* **2014**, *171* (23), 5252-5264.
4. Wang, G.; Peng, Z.; He, D.; Yan, C.; Liu, W. Coumarin-isatin type compound useful in treatment of diabetes mellitus and its preparation. CN105237521A, 2016.
5. Mercer, D. K.; Robertson, J.; Wright, K.; Miller, L.; Smith, S.; Stewart, C. S.; O'Neil, D. A., A prodrug approach to the use of coumarins as potential therapeutics for superficial mycoses. *PLoS One* **2013**, *8* (11), e80760.
6. Musa, M. A.; Cooperwood, J. S.; Khan, M. O. F., A review of coumarin derivatives in pharmacotherapy of breast cancer. *Curr. Med. Chem.* **2008**, *15* (26), 2664-2679.
7. Mi, C.; Ma, J.; Wang, K. S.; Zuo, H. X.; Wang, Z.; Li, M. Y.; Piao, L. X.; Xu, G. H.; Li, X.; Quan, Z. S.; Jin, X., Imperatorin suppresses proliferation and angiogenesis of human colon cancer cell by targeting HIF- $\alpha$  via the mTOR/p70S6K/4E-BP1 and MAPK pathways. *J. Ethnopharmacol.* **2017**, *203*, 27-38.
8. Xia, W.; Gooden, D.; Liu, L.; Zhao, S.; Soderblom, E. J.; Toone, E. J.; Beyer, W. F.; Walder, H.; Spector, N. L., Photo-activated psoralen binds the ErbB2 catalytic kinase domain, blocking ErbB2 signaling and triggering tumor cell apoptosis. *PLoS One* **2014**, *9* (2), e88983/1-e88983/10, 10 pp.
9. Leanza, L.; Romio, M.; Becker, K. A.; Azzolini, M.; Trentin, L.; Manago, A.; Venturini, E.; Zaccagnino, A.; Mattarei, A.; Carraretto, L.; Urbani, A.; Kadow, S.; Biasutto, L.; Martini, V.; Severin, F.; Peruzzo, R.; Trimarco, V.; Egberts, J.-H.; Hauser, C.; Visentin, A.; Semenzato, G.; Kalthoff, H.; Zoratti, M.; Gulbins, E.; Paradisi, C.; Szabo, I., Direct pharmacological targeting of a mitochondrial ion channel selectively kills tumor cells in vivo. *Cancer Cell* **2017**, *31* (4), 516-531.e10.
10. Min, K.-J.; Um, H. J.; Seo, S. U.; Woo, S. M.; Kim, S.; Park, J.-W.; Lee, H.-S.; Kim, S. H.; Choi, Y. H.; Lee, T.-J.; Kwon, T. K., Angelicin potentiates TRAIL-induced apoptosis in renal carcinoma Caki cells through activation of caspase 3 and down-regulation of c-FLIP expression. *Drug Dev. Res.* **2018**, *79* (1), 3-10.
11. Gustafson, K. R.; Fuller, R. W.; Cardellina, J. H., II; McMahan, J. B.; Currens, M. J.; Boyd, M. R.; Kashman, Y.; Buckheit, R. W., Jr.; Hughes, S. H.; Cragg, G. M., The Calanolides, a novel HIV-inhibitory class of coumarin derivatives from the tropical rainforest tree, *Calophyllum lanigerum*. *J. Med. Chem.* **1992**, *35* (15), 2735-2743.
12. Koziol, E.; Skalicka-Woźniak, K., Imperatorin-pharmacological meaning and analytical clues: profound investigation. *Phytochem. Rev.* **2016**, *15* (4), 627-649.
13. Chiang, C. C.; Cheng, M. J.; Peng, C. F.; Huang, H. Y.; Chen, I. S., A novel dimeric coumarin analog and antimycobacterial constituents from *Fatoua pilosa*. *Chemistry and Biodiversity* **2010**, *7* (7), 1728-1736.
14. Hsieh, M.-J.; Chen, M.-K.; Yu, Y.-Y.; Sheu, G.-T.; Chiou, H.-L., Psoralen reverses docetaxel-induced multidrug resistance in A549/D16 human lung cancer cells lines. *Phytomedicine* **2014**, *21* (7), 970-977.
15. Jiang, J.; Wang, X.; Cheng, K.; Zhao, W.; Hua, Y.; Xu, C.; Yang, Z., Psoralen reverses the P-glycoprotein-mediated multidrug resistance in human breast cancer MCF-7/ADR cells. *Mol. Med. Rep.* **2016**, *13* (6), 4745-4750.
16. Touisni, N.; Maresca, A.; McDonald, P. C.; Lou, Y.; Scozzafava, A.; Dedhar, S.; Winum, J.-Y.; Supuran, C. T., Glycosyl coumarin carbonic anhydrase IX and XII inhibitors strongly attenuate the growth of primary breast tumors. *J. Med. Chem.* **2011**, *54* (24), 8271-8277.
17. Maresca, A.; Supuran, C. T., Coumarins incorporating hydroxy- and chloro-moieties selectively inhibit the transmembrane, tumor-associated carbonic anhydrase isoforms IX and XII over the cytosolic ones I and II. *Bioorg. Med. Chem. Lett.* **2010**, *20* (15), 4511-4.
18. Nocentini, A.; Carta, F.; Ceruso, M.; Bartolucci, G.; Supuran, C. T., Click-tailed coumarins with potent and selective inhibitory action against the tumor-associated carbonic anhydrases IX and XII. *Bioorg. Med. Chem.* **2015**, *23* (21), 6955-6966.
19. De Luca, L.; Mancuso, F.; Ferro, S.; Buemi, M. R.; Angeli, A.; Del Prete, S.; Capasso, C.; Supuran, C. T.; Gitto, R., Inhibitory effects and structural insights for a novel series of coumarin-based compounds that selectively target human CA IX and CA XII carbonic anhydrases. *Eur. J. Med. Chem.* **2018**, *143*, 276-282.
20. Supuran, C. T., Inhibition of carbonic anhydrase IX as a novel anticancer mechanism. *World J. Clin. Oncol.* **2012**, *3* (7), 98-103.
21. Winum, J.-Y.; Rami, M.; Scozzafava, A.; Montero, J.-L.; Supuran, C., Carbonic anhydrase IX: a new druggable target for the design of antitumor agents. *Med. Res. Rev.* **2008**, *28* (3), 445-463.
22. Cecchi, A.; Hulikova, A.; Pastorek, J.; Pastorekova, S.; Scozzafava, A.; Winum, J. Y.; Montero, J. L.; Supuran, C. T., Carbonic anhydrase inhibitors. Design of fluorescent sulfonamides as probes of tumor-associated carbonic anhydrase IX that inhibit isozyme IX-mediated acidification of hypoxic tumors. *J. Med. Chem.* **2005**, *48* (15), 4834-41.
23. Supuran, C. T.; Winum, J. Y., Carbonic anhydrase IX inhibitors in cancer therapy: an update. *Future Med Chem* **2015**, *7* (11), 1407-14.
24. Araste, F.; Ebrahimzadeh, W.; Rasooli, I.; Rajabibazl, M.; Mousavi Gargari, S. L., A novel VHH nanobody against the active site (the CA domain) of tumor-associated, carbonic anhydrase isoform IX and its usefulness for cancer diagnosis. *Biotechnol. Lett.* **2014**, *36* (1), 21-28.
25. De Simone, G.; Supuran, C. T., Carbonic anhydrase IX: Biochemical and crystallographic characterization of a novel antitumor target. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* **2010**, *1804* (2), 404-409.
26. Horie, K.; Kawakami, K.; Fujita, Y.; Sugaya, M.; Kameyama, K.; Mizutani, K.; Deguchi, T.; Ito, M., Exosomes expressing carbonic anhydrase 9 promote angiogenesis. *Biochem. Biophys. Res. Commun.* **2017**, *492* (3), 356-361.
27. Supuran, C. T., Structure and function of carbonic anhydrases. *Biochem. J.* **2016**, *473* (14), 2023-32.
28. Jiang, J.; Zhao, J. H.; Wang, X. L.; Guo, X. J.; Yang, J.; Bai, X.; Jin, S. Y.; Ge, R. L., Correlation between carbonic anhydrase IX (CA-9), XII (CA-12) and hypoxia inducible factor-2 $\alpha$  (HIF-2 $\alpha$ ) in breast cancer. *Neoplasma* **2015**, *62* (3), 456-463.
29. Rajabi, M.; Hossaini, Z.; Khalilzadeh, M. A.; Datta, S.; Halder, M.; Mousa, S. A., Synthesis of a new class of furo[3,2-c]coumarins and its anticancer activity. *J. Photochem. Photobiol. B* **2015**, *148*, 66-72.
30. Abdel Hafez, O. M.; Amin, K. M.; Abdel-Latif, N. A.; Mohamed, T. K.; Ahmed, E. Y.; Maher, T., Synthesis and



- antitumor activity of some new xanthotoxin derivatives. *Eur. J. Med. Chem.* **2009**, *44* (7), 2967-74.
31. Melis, C.; Meleddu, R.; Angeli, A.; Distinto, S.; Bianco, G.; Capasso, C.; Cottiglia, F.; Angius, R.; Supuran, C. T.; Maccioni, E., Isatin: a privileged scaffold for the design of carbonic anhydrase inhibitors. *J. Enzyme Inhib. Med. Chem.* **2017**, *32* (1), 68-73.
32. Bianco, G.; Meleddu, R.; Distinto, S.; Cottiglia, F.; Gaspari, M.; Melis, C.; Corona, A.; Angius, R.; Angeli, A.; Taverna, D.; Alcaro, S.; Leitans, J.; Kazaks, A.; Tars, K.; Supuran, C. T.; Maccioni, E., N-Acylbenzenesulfonamide dihydro-1,3,4-oxadiazole hybrids: seeking selectivity toward carbonic anhydrase isoforms. *ACS Med. Chem. Lett.* **2017**, *8* (8), 792-796.
33. Maresca, A.; Temperini, C.; Vu, H.; Pham, N. B.; Poulsen, S. A.; Scozzafava, A.; Quinn, R. J.; Supuran, C. T., Non-zinc mediated inhibition of carbonic anhydrases: coumarins are a new class of suicide inhibitors. *J. Am. Chem. Soc.* **2009**, *131* (8), 3057-62.
34. Leitans, J.; Kazaks, A.; Balode, A.; Ivanova, J.; Zalubovskis, R.; Supuran, C. T.; Tars, K., Efficient expression and crystallization system of cancer-associated carbonic anhydrase isoform IX. *J. Med. Chem.* **2015**, *58* (22), 9004-9.
35. Chung, J. Y.; Hah, J. M.; Cho, A. E., Correlation between performance of QM/MM docking and simple classification of binding sites. *J. Chem. Inf. Model.* **2009**, *49* (10), 2382-7.
36. Kollman, P. A.; Massova, I.; Reyes, C.; Kuhn, B.; Huo, S.; Chong, L.; Lee, M.; Lee, T.; Duan, Y.; Wang, W.; Donini, O.; Cieplak, P.; Srinivasan, J.; Case, D. A.; Cheatham, T. E., Calculating structures and free energies of complex molecules: combining molecular mechanics and continuum models. *Acc. Chem. Res.* **2000**, *33* (12), 889-897.
37. Meleddu, R.; Maccioni, E.; Distinto, S.; Bianco, G.; Melis, C.; Alcaro, S.; Cottiglia, F.; Ceruso, M.; Supuran, C. T., New 4-[(3-cyclohexyl-4-aryl-2,3-dihydro-1,3-thiazol-2-ylidene)amino]benzene-1-sulfonamides, synthesis and inhibitory activity toward carbonic anhydrase I, II, IX, XII. *Bioorg. Med. Chem. Lett.* **2015**, *25* (16), 3281-4.
38. Supuran, C. T., Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nat. Rev. Drug Discov.* **2008**, *7* (2), 168.

## Targeting tumor associated carbonic anhydrase IX and XII: highly isozyme selective coumarin and psoralen inhibitors

Claudia Melis, Simona Distinto, Giulia Bianco, Rita Meleddu, Filippo Cottiglia, Benedetta Fois, Domenico Taverna, Rossella Angius, Stefano Alcaro, Francesco Ortuso, Marco Gaspari, Andrea Angeli, Sonia Del Prete, Clemente Capasso, Claudiu T. Supuran and Elias Maccioni.

