

Search for the rare decay of charmed baryon Λ_c^+ into the $p\mu^+\mu^-$ final state

R. Aaij *et al.**
(LHCb Collaboration)

 (Received 22 July 2024; accepted 27 August 2024; published 24 September 2024)

A search for the nonresonant $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ decay is performed using proton-proton collision data recorded at a center-of-mass energy of 13 TeV by the LHCb experiment, corresponding to an integrated luminosity of 5.4 fb^{-1} . No evidence for the decay is found in the dimuon invariant-mass regions where the expected contributions of resonances is subdominant. The upper limit on the branching fraction of the $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ decay is determined to be $2.9(3.2) \times 10^{-8}$ at 90%(95%) confidence level. The branching fractions in the dimuon invariant-mass regions dominated by the η , ρ and ω resonances are also determined.

DOI: [10.1103/PhysRevD.110.052007](https://doi.org/10.1103/PhysRevD.110.052007)

I. INTRODUCTION

The flavor-changing neutral-current decay¹ $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ is heavily suppressed in the Standard Model (SM) as it is forbidden at tree level and is affected by the Glashow-Iliopoulos-Maiani mechanism [1]. The phenomenology of the decay is discussed in Ref. [2] and the SM predictions for q^2 differential branching fractions, where q^2 is the dimuon invariant-mass squared, are calculated in Ref. [3]. The branching fraction (\mathcal{B}) associated with the short-distance contributions to the $c \rightarrow u\mu^+\mu^-$ transition is expected to be below 10^{-8} in the SM, with an enhancement up to 10^{-6} due to long-distance processes involving intermediate vector-meson resonances that decay to a lepton pair [4,5]. The estimation of resonant SM contributions relies on the measured values for the corresponding branching fractions and the isospin relation between ρ and ω states, to overcome the lack of experimental information on $\mathcal{B}(\Lambda_c^+ \rightarrow p\rho)$. Strong phases between the resonances, and thus the effect of interference, are *a priori* unknown.

Interesting deviations from the SM predictions have been observed in the equivalent type of transition for the b quark, $b \rightarrow s\ell\ell$, in particular in angular observables [6]. These anomalies motivate the extension of this physics program into the charm sector, which is far less explored both experimentally and theoretically. The first results, such as an angular analysis of charm mesons [7], are already available. A discussion of the discovery potential of effects

beyond the SM in rare charm-baryon decays can be found in Ref. [8].

The LHCb collaboration previously searched for the $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ decay using the proton-proton (pp) data collected in 2011–2012 [9] and placed an upper limit of 7.7×10^{-8} at 90% confidence level (CL), thereby improving the result from the *BABAR* experiment [10] by two orders of magnitude. The analysis of data collected in 2016–2018 presented in this paper follows the methodology of the previous analysis, and relies on improved techniques with the use of the CL_s method [11] implementation based on unbinned likelihoods, the incorporation of background due to particle misidentification into the data model, a different definition of the short-distance signal and a more detailed treatment of the entire range of dimuon invariant mass where resonances are expected to dominate. Accordingly, the new results for the short distance contribution should not be combined with the results based on the 2011–2012 data.

The measurement is carried out in different regions of dimuon invariant mass $m_{\mu\mu}$ and the $\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-)$ decay is used for normalization since it has the same final state as the signal and is relatively abundant.

II. DETECTOR AND SIMULATION

The LHCb detector [12,13] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [14], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [15] placed downstream of the magnet. The tracking system provides a measurement of the momentum, p , of charged particles with a relative

*Full author list given at the end of the article.

¹The charge-conjugate mode is implied throughout the paper.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/ c . The minimum distance of a track to a primary pp collision vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV/ c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [16]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multi-wire proportional chambers [17]. The online event selection is performed by a trigger system [18], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction.

Simulated samples of Λ_c^+ baryons are produced to model the effects of the detector acceptance and the imposed selection requirements. The pp collisions are generated using Pythia [19] with a specific LHCb configuration [20]. Decays of unstable particles are described by EvtGen [21], in which final state radiation is generated using Photos [22]. In the case of $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ decays, a model with final state particles equally distributed in the phase space is used. The interaction of the generated particles with the detector, and its response, is implemented using the Geant4 toolkit [23] as described in Ref. [24]. The underlying pp interaction and Λ_c^+ baryon are reused multiple times to speed up the simulation process, with independently generated Λ_c^+ decay products [25].

III. CANDIDATE SELECTION

The selection criteria largely follow those described in Ref. [9]. The Λ_c^+ candidates are reconstructed by combining a pair of oppositely charged tracks identified as muons with an additional track identified as a proton. The three tracks are required to be compatible with originating from a common vertex and have significant χ_{IP}^2 with respect to all PVs. Here, χ_{IP}^2 is defined as the difference in the vertex-fit χ^2 of a given PV, reconstructed with and without the track under consideration. The Λ_c^+ candidates must have a decay vertex displaced from any PV and be compatible with originating from one of the PVs.

The background formed by random combinations of tracks, referred to as combinatorial background, is suppressed using a multivariate classifier. A boosted decision tree (BDT) algorithm [26], as implemented in the TMVA toolkit [27,28], is trained using supervised learning with tenfold cross-validation [29] to achieve an unbiased classifier response. The BDT discrimination is done in two steps. A first BDT classifier employs topological and kinematic variables of the Λ_c^+ candidates. A relatively loose requirement on the response of that BDT is applied as the last preselection step to enhance the purity of the

normalization data sample. After preselection, simulated samples of $\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-)$ decays are compared with data, and weights are determined to improve the agreement between data and simulation for the Λ_c^+ transverse momentum and the number of tracks in the event. A second BDT classifier is trained using a set of variables that includes those of the first classifier and, in addition, kinematic, vertex and isolation variables related to the decay products. For the two classifiers, simulated events containing $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ decays are used as a proxy for the signal, while the background is modeled using sideband events with a $p\mu^+\mu^-$ invariant mass within [2140, 2251] and [2321, 2430] MeV/ c^2 . The optimization of final requirements is performed in three dimensions: the response value of the second BDT classifier and two variables characterizing the proton and muon particle identification (PID). In this optimization, the simulation is corrected for imperfections in modeling of the PID performance [30]. The final requirements are determined using pseudoexperiments by finding the best expected upper limit (using the CL_s method) on the ratio of branching fractions of the signal relative to the normalization channel. At this stage, only two background sources are expected to significantly contribute: combinatorial background and $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ decays where both pions are misidentified as muons. In each pseudoexperiment, pseudodata for the signal and these two background components are drawn from the three-dimensional probability density functions determined after the preselection stage.

Six dimuon invariant-mass ranges are considered as presented in Table I. The nonresonant signal decay is searched for in two $m_{\mu\mu}$ regions, *low- m* and *high- m* , where the expected contributions of resonances are subdominant. In the following, the combination of these two ranges is referred to as the *signal* region. The non *signal* range is divided into regions where either the η , ρ , ω or ϕ resonance dominates. The branching fraction results quoted for a given resonant region are labeled by the resonant decay expected to dominate in that region. However, due to the relatively low sample size, no attempt to separate the resonant contributions within a given dimuon invariant-

TABLE I. Definitions of the dimuon invariant-mass ranges. The lower (upper) bound of the *low- m* (*high- m*) region corresponds to the phase-space limits.

Region	Range [MeV/ c^2]
<i>low-m</i>	$211.32 < m_{\mu^+\mu^-} < 507.86$
<i>high-m</i>	$1059.46 < m_{\mu^+\mu^-} < 1348.13$
η	$507.86 < m_{\mu^+\mu^-} < 587.86$
ω	$742.65 < m_{\mu^+\mu^-} < 822.65$
ρ	$587.86 < m_{\mu^+\mu^-} < 742.65$
	or $822.65 < m_{\mu^+\mu^-} < 965.20$
ϕ	$979.46 < m_{\mu^+\mu^-} < 1059.46$

mass range is made. Therefore it has to be emphasized that the results must be interpreted as corresponding to that particular $m_{\mu\mu}$ region rather than to a specific resonant decay.

IV. NORMALIZATION AND INVARIANT-MASS FIT

The $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ branching fraction is determined from the measured signal yield N_{sig} according to

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) = \alpha \times N_{\text{sig}}, \quad (1)$$

where the normalization factor α is given by

$$\alpha = \frac{\epsilon_{\text{norm}}}{\epsilon_{\text{sig}} \times N_{\text{norm}}} \times \mathcal{B}(\Lambda_c^+ \rightarrow p\phi) \times \mathcal{B}(\phi \rightarrow \mu^+\mu^-). \quad (2)$$

Here, N_{norm} is the yield of $\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-)$ decays, while ϵ_{sig} and ϵ_{norm} represent the corresponding total efficiencies for signal and normalization channels, respectively. In the case of the ratio of branching fractions $\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)/\mathcal{B}(\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-))$, the corresponding normalization factor α takes the form $\epsilon_{\text{norm}}/(\epsilon_{\text{sig}} \times N_{\text{norm}})$. The ratio of efficiencies $\epsilon_{\text{norm}}/\epsilon_{\text{sig}}$ for the signal regions is compatible with unity within statistical uncertainty.

An extended unbinned maximum-likelihood fit to the $p\mu^+\mu^-$ invariant mass is performed in each $m_{\mu\mu}$ region, where the signal shape of Λ_c^+ candidates is described by a Crystal Ball function [31]. The background is described using two distinct components: the combinatorial background, modeled by an exponential shape, and the $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ peaking background, represented by the sum of a Crystal Ball function and a Gaussian function. The shapes of both the signal and the $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ background are determined from simulation for each $m_{\mu\mu}$ region and fixed in the fit to data. Weights are applied to simulated $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ events to match the phase-space distribution observed in Ref. [32].

In each $m_{\mu\mu}$ region, there are four parameters in the fit. Three of them are free parameters: the signal yield, the combinatorial background yield and the exponent describing the shape of the combinatorial background. The fourth one corresponds to the $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ background. Its total yield in all regions is a free parameter, and the fraction in each region is constrained based on simulation.

The results of the fit to the data for the *signal* region and for the *low- m* and *high- m* regions are presented in Fig. 1. The results of the fit for the normalization mode and other regions dominated by resonances are presented in Fig. 2. The yields and the Λ_c^+ signal significance for the six $m_{\mu\mu}$ regions are reported in Table II, where the significance is




FIG. 1. Distributions of $p\mu^+\mu^-$ invariant mass for (top) *signal*, (bottom left) *low- m* and (bottom right) *high- m* regions. The results of the fit are overlaid.




FIG. 2. Distributions of $p\mu^+\mu^-$ invariant mass for (top left) ϕ , (top right) ω , (bottom left) ρ and (bottom right) η regions. The results of the fit are overlaid.

determined using Wilks' theorem [33]. For the signal regions the upper limits for the corresponding branching fractions are determined. The residual contribution from resonant decays into signal regions is estimated from simulation neglecting interference effects. For the *high-m* region, where the significance is the highest, only 2.8 events are attributed to $\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-)$ decays while for the *low-m* region the expected contribution is negligible.

V. SYSTEMATIC UNCERTAINTIES

The systematic uncertainties taken into account in the determination of the upper limit on the branching fraction are summarized in Table III. The two main contributions are the statistical uncertainty of the simulation samples

used to determine the efficiency ratio and the uncertainty on the yield of the normalization mode. The latter is dominated by the statistical uncertainty on the yield returned from the fit. Other sources of systematic uncertainty on the efficiency ratio are associated with the following:

- (1) The level of agreement between data and simulation. The maximum difference between ratios of efficiencies determined for a few variants of the baseline procedure of weighting simulation to match the data is taken as a systematic uncertainty.
- (2) The efficiency determination from simulation. The difference in the efficiency obtained including or excluding Λ_c^+ candidates, for which one or more decay products are not correctly matched to the generated particles, is taken as a systematic

TABLE II. Results of the fit to the $p\mu^+\mu^-$ invariant-mass distribution for various $m_{\mu\mu}$ regions. The significance of the $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ component is also given.

Region	$\Lambda_c^+ \rightarrow p\mu^+\mu^-$ yield	$\Lambda_c^+ \rightarrow p\pi^+\pi^-$ yield	Combinatorial yield	Significance $\Lambda_c^+ \rightarrow p\mu^+\mu^-$
<i>signal</i>	18 ± 10	3 ± 7	681 ± 28	2.0σ
<i>low-m</i>	1 ± 5	4 ± 4	241 ± 17	0.3σ
<i>high-m</i>	21 ± 8	4 ± 4	432 ± 22	2.8σ
η	12 ± 5	2.2 ± 1.6	84 ± 10	3.0σ
ρ	43 ± 10	20 ± 6	382 ± 22	5.6σ
ω	81 ± 10	4.8 ± 2.1	101 ± 11	$>7\sigma$
ϕ	423 ± 22	3.8 ± 2.4	173 ± 15	$>7\sigma$

TABLE III. Summary of systematic uncertainties for the signal regions and for the η , ρ , and ω resonant regions.

Uncertainty source [%]	<i>signal</i>	<i>low-m</i>	<i>high-m</i>	η	ρ	ω
Normalization channel	5.3	5.3	5.3	5.3	5.3	5.3
Efficiency ratio (stat.)	5.7	8.2	6.8	12.9	5.3	4.6
Efficiency ratio (syst.)	1.7	1.7	1.7	1.7	1.7	1.7
Shape of signal	1.6	3.4	1.1	0.6	1.7	1.2
Shape of $\Lambda_c^+ \rightarrow p\pi^+\pi^-$	0.1	3.2	0.2	0.4	5.3	1.2
Shape of combinatorial	0.1	0.3	0.2	0.1	0.0	0.0
$\Lambda_c^+ \rightarrow p\pi^+\pi^-$ decay model	0.1	0.1	0.1	0.1	0.1	0.1
Fit bias	0.1	0.1	0.2	0.3	0.1	0.1
Total	8.1	11.0	8.9	14.1	9.5	7.4

uncertainty. The procedure is applied consistently for the signal and normalization modes.

- (3) The correction of PID variables in simulation. The difference of the efficiency ratio for the baseline and an alternative parametrization of the PID calibration procedure [30] is taken as a systematic uncertainty.

These three sources added in quadrature amount to a relative uncertainty of 1.7%.

The systematic uncertainties related to the fit model are estimated using pseudoexperiments. An alternative model for each component is used to generate pseudodata, which are then fitted using the baseline model described in Sec. IV. A Johnson function [34] is used as an alternative shape for the signal, while the $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ background is parametrized as the sum of a Johnson and a Gaussian function, and a linear function is used as an alternative model for the combinatorial background. The difference between the generated signal yield and the average fitted yield is taken as systematic uncertainty. In addition, the uncertainty related to the $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ decay model is determined by applying different corrections to the $\Lambda_c^+ \rightarrow p\pi^+\pi^-$ simulated sample and determining an alternative invariant-mass shape. The bias due to maximum-likelihood fit is determined from the same data model used for pseudodata generation and fitting. The signal yield difference between the generated value and the average of fitted values from the pseudoexperiments is taken as a systematic uncertainty.

The systematic uncertainties associated with the ratio of branching fractions in the resonant regions are determined

in similar way as for the signal regions and are presented in Table III.

VI. RESULTS

A. Signal regions

No significant contribution of $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ decays is observed in either of the two signal regions. Upper limits on the branching fractions and their ratios are determined using the CL_s method implemented in the GammaCombo package [35]. The systematic uncertainties are included in the determination of CL_s. The following upper limit in the *signal* region is determined to be

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi)\mathcal{B}(\phi \rightarrow \mu^+\mu^-)} < 0.09(0.10) \text{ at } 90\%(95\%) \text{ CL.}$$

Using the values of the branching fractions for $\Lambda_c^+ \rightarrow p\phi$ and $\phi \rightarrow \mu^+\mu^-$ decays from Ref. [36] and including their uncertainties in the CL_s determination, an upper limit on the branching fraction is determined to be

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 2.9(3.2) \times 10^{-8} \text{ at } 90\%(95\%) \text{ CL.}$$

The CL_s as a function of branching fraction is shown in Fig. 3. The limits are determined for *low-m* and *high-m* regions separately with the same assumptions as for the *signal* region. The following upper limits are determined:

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi)\mathcal{B}(\phi \rightarrow \mu^+\mu^-)} < 0.029(0.034) \text{ at } 90\%(95\%) \text{ CL } (low-m),$$

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi)\mathcal{B}(\phi \rightarrow \mu^+\mu^-)} < 0.094(0.10) \text{ at } 90\%(95\%) \text{ CL } (high-m),$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 0.93(1.1) \times 10^{-8} \text{ at } 90\%(95\%) \text{ CL } (low-m),$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 3.0(3.3) \times 10^{-8} \text{ at } 90\%(95\%) \text{ CL } (high-m).$$

The upper limit placed using data from the *high-m* region is driven by the signal yield whose significance is close to but lower than 3σ , while in the *low-m* region the invariant-mass distribution is fully compatible with no signal.

Assuming a decay model with final state particles equally distributed in the phase space, the values of upper limits extrapolated from the *signal* region to the full phase space are determined to be

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi)\mathcal{B}(\phi \rightarrow \mu^+\mu^-)} < 0.23(0.25) \text{ at } 90\%(95\%) \text{ CL},$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 7.3(8.2) \times 10^{-8} \text{ at } 90\%(95\%) \text{ CL}.$$

In spite of the enlarged data sample, the extrapolated upper limit on the branching fraction is only slightly better than that determined for the 2011–2012 data [9], $7.7(9.6) \times 10^{-8}$ at 90%(95%) CL. This relatively small improvement is driven mainly by the 2.8σ effect observed in the *high-m* region in this analysis.

B. Resonant regions

Excluding the normalization channel, the significance of the Λ_c^+ signal exceeds the 5σ threshold in the ρ and ω regions while the significance for the $\Lambda_c^+ \rightarrow p\eta(\rightarrow \mu^+\mu^-)$ decay is at 3σ . The corresponding branching fractions are determined using Eqs. (1) and (2) where the decay parameters for the nonresonant decays are replaced by those for the resonant decays. It has to be emphasized that disentangling the resonant components to account for interference effects has not been attempted. The efficiency

for a given $m_{\mu\mu}$ region is determined using simulation samples of the corresponding resonant decay, thus neglecting possible contributions from other resonances. Under this assumption, the ratio of branching fractions with respect to the normalization channel are determined for the ω , ρ and η regions to be

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\omega(\rightarrow \mu^+\mu^-))}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-))} = 0.240 \pm 0.030(\text{stat}) \pm 0.018(\text{syst}),$$

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\rho(\rightarrow \mu^+\mu^-))}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-))} = 0.229 \pm 0.051(\text{stat}) \pm 0.022(\text{syst}),$$

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\eta(\rightarrow \mu^+\mu^-))}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-))} = 0.032 \pm 0.013(\text{stat}) \pm 0.004(\text{syst}).$$

The corresponding branching fractions are determined to be

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\omega) = (9.82 \pm 1.23(\text{stat}) \pm 0.73(\text{syst}) \pm 2.79(\text{ext})) \times 10^{-4},$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\rho) = (1.52 \pm 0.34(\text{stat}) \pm 0.14(\text{syst}) \pm 0.24(\text{ext})) \times 10^{-3},$$

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\eta) = (1.67 \pm 0.69(\text{stat}) \pm 0.23(\text{syst}) \pm 0.34(\text{ext})) \times 10^{-3},$$

where the second uncertainties are induced by the ratio of branching fractions presented in Table III and the third




FIG. 3. Observed and expected (background only hypothesis) limits for $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ branching fraction.

uncertainties are due to the following branching fractions: $\mathcal{B}(\Lambda_c^+ \rightarrow p\phi)$, $\mathcal{B}(\phi \rightarrow \mu^+\mu^-)$, $\mathcal{B}(\omega \rightarrow \mu^+\mu^-)$, $\mathcal{B}(\rho \rightarrow \mu^+\mu^-)$ and $\mathcal{B}(\eta \rightarrow \mu^+\mu^-)$ [36], which were used as external inputs.

VII. SUMMARY

A search for the nonresonant $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ decay is presented using data samples of proton-proton collisions collected by the LHCb experiment in 2016–2018, corresponding to an integrated luminosity of 5.4 fb^{-1} . In the dimuon invariant-mass regions where the contributions of resonant decays is subdominant, *i.e.* below the η and above the ϕ resonances, the following upper limits for the branching fractions are derived: $\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 0.93(1.1) \times 10^{-8}$ at 90% (95%) CL and $\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 3.0(3.3) \times 10^{-8}$ at 90% (95%) CL, respectively where $\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-)$ as a normalization channel. Combining these two regions, one finds $\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 2.9(3.2) \times 10^{-8}$ at 90% (95%) CL. For the reasons described

in the introduction, the results for the short-distance contribution presented in this paper should not be combined with the results published for 2011–2012 data [9].

In a narrow region around the ω resonance, a Λ_c^+ signal is observed with a significance of more than 7σ , corresponding to a branching fraction $(9.82 \pm 1.23(\text{stat}) \pm 0.73(\text{syst}) \pm 2.79(\text{ext})) \times 10^{-4}$, compatible with and more precise than the result of the previous analysis for the data collected in 2011–2012 [9].

Excluding the region around the ω meson, a signal in the ρ region is found with a significance of 5.6σ . The corresponding branching fraction determined with respect to the normalization channel is measured to be $(1.52 \pm 0.34(\text{stat}) \pm 0.14(\text{syst}) \pm 0.24(\text{ext})) \times 10^{-3}$, giving strong evidence for the $\Lambda_c^+ \rightarrow p\rho$ decay, qualified by the fact that interference effects are not accounted for. For the η region the branching fraction was measured to be $(1.67 \pm 0.69(\text{stat}) \pm 0.23(\text{syst}) \pm 0.34(\text{ext})) \times 10^{-3}$ with a significance of 3.0σ . The value is compatible with the world average [36] but less precise.

ACKNOWLEDGMENTS

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ

and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MCID/IFA (Romania); MICIU and AEI (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), and Polish WLCG (Poland). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); Minciencias (Colombia); EPLANET, Marie Skłodowska-Curie Actions, ERC and NextGenerationEU (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); AvH Foundation (Germany); ICSC (Italy); Severo Ochoa and María de Maeztu Units of Excellence, GVA, XuntaGal, GENCAT, InTalent-Inditex and Prog. Atracción Talento CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

-
- [1] S. Glashow, J. Iliopoulos, and L. Maiani, *Phys. Rev. D* **2**, 1285 (1970).
- [2] S. Meinel, *Phys. Rev. D* **97**, 034511 (2018).
- [3] M. Golz, G. Hiller, and T. Magorsch, *J. High Energy Phys.* **09** (2021) 208.
- [4] S. Fajfer, S. Prelovsek and P. Singer, *Phys. Rev. D* **64**, 114009 (2001).
- [5] G. Burdman, E. Golowich, J. Hewett, and S. Pakvasa, *Phys. Rev. D* **66**, 014009 (2002).
- [6] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **126**, 161802 (2021).
- [7] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **128**, 221801 (2022).
- [8] H. Gisbert, M. Golz, and D. Mitzel, *Mod. Phys. Lett. A* **36**, 2130002 (2021).
- [9] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. D* **97**, 091101 (2018).
- [10] J. P. Lees *et al.* (BABAR Collaboration), *Phys. Rev. D* **84**, 072006 (2011).
- [11] A. L. Read, *J. Phys. G* **28**, 2693 (2002).
- [12] A. A. Alves Jr. *et al.* (LHCb Collaboration), *J. Instrum.* **3**, S08005 (2008).
- [13] LHCb Collaboration, *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
- [14] R. Aaij *et al.*, *J. Instrum.* **9**, P09007 (2014).
- [15] P. d’Argent *et al.*, *J. Instrum.* **12**, P11016 (2017).
- [16] M. Adinolfi *et al.*, *Eur. Phys. J. C* **73**, 2431 (2013).
- [17] A. A. Alves Jr. *et al.*, *J. Instrum.* **8**, P02022 (2013).
- [18] R. Aaij *et al.*, *J. Instrum.* **8**, P04022 (2013).
- [19] T. Sjöstrand, S. Mrenna, and P. Skands, *Comput. Phys. Commun.* **178**, 852 (2008); *J. High Energy Phys.* **05** (2006) 026.
- [20] I. Belyaev *et al.*, *J. Phys. Conf. Ser.* **331**, 032047 (2011).
- [21] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [22] N. Davidson, T. Przedzinski, and Z. Was, *Comput. Phys. Commun.* **199**, 86 (2016).
- [23] J. Allison *et al.* (Geant4 Collaboration), *IEEE Trans. Nucl. Sci.* **53**, 270 (2006); S. Agostinelli *et al.* (Geant4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [24] M. Clemencic, G. Corti, S. Easo, C. R. Jones, S. Miglioranza, M. Pappagallo, and P. Robbe, *J. Phys. Conf. Ser.* **331**, 032023 (2011).

- [25] D. Müller, M. Clemencic, G. Corti, and M. Gersabeck, *Eur. Phys. J. C* **78**, 1009 (2018).
- [26] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and Regression Trees* (Wadsworth international group, Belmont, California, USA, 1984).
- [27] H. Voss, A. Hoecker, J. Stelzer, and F. Tegenfeldt, *Proc. Sci. ACAT2007* (2007) 040.
- [28] A. Hoecker *et al.*, [arXiv:physics/0703039](https://arxiv.org/abs/physics/0703039).
- [29] A. Blum, A. Kalai, and J. Langford, Beating the hold-out: Bounds for k-fold and progressive cross-validation, in *COLT '99 Proceedings of the Twelfth Annual Conference on Computational Learning Theory* (Association for Computing Machinery, New York, NY, 1999), pp. 203–208, <https://dl.acm.org/doi/proceedings/10.1145/307400>.
- [30] R. Aaij *et al.*, *Eur. Phys. J. Tech. Instrum.* **6**, 1 (2019).
- [31] T. Skwarnicki, A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances, Ph.D. thesis, Institute of Nuclear Physics, Krakow, 1986 [DESY-F31-86-02], <http://inspirehep.net/record/230779/>.
- [32] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **03** (2018) 182.
- [33] S. S. Wilks, *Ann. Math. Stat.* **9**, 60 (1938).
- [34] N. Johnson, *Biom. J.* **36**, 149 (1949).
- [35] M. Kenzie *et al.*, GammaCombo: A statistical analysis framework for combining measurements, fitting datasets and producing confidence intervals, [10.5281/zenodo.3371421](https://doi.org/10.5281/zenodo.3371421).
- [36] R. L. Workman *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).

R. Aaij³⁶, A. S. W. Abdelmotteleb⁵⁵, C. Abellan Beteta⁴⁹, F. Abudinén⁵⁵, T. Ackernley⁵⁹, A. A. Adefisoye⁶⁷, B. Adeva⁴⁵, M. Adinolfi⁵³, P. Adlarson⁷⁹, C. Agapopoulou¹³, C. A. Aidala⁸⁰, Z. Ajaltouni¹¹, S. Akar⁶⁴, K. Akiba³⁶, P. Albicocco²⁶, J. Albrecht¹⁸, F. Alessio⁴⁷, M. Alexander⁵⁸, Z. Aliouche⁶¹, P. Alvarez Cartelle⁵⁴, R. Amalric¹⁵, S. Amato³, J. L. Amey⁵³, Y. Amhis^{13,47}, L. An⁶, L. Anderlini²⁵, M. Andersson⁴⁹, A. Andreianov⁴², P. Andreola⁴⁹, M. Andreotti²⁴, D. Andreou⁶⁷, A. Anelli^{29,b}, D. Ao⁷, F. Archilli^{35,c}, M. Argenton²⁴, S. Arguedas Cuendis⁹, A. Artamonov⁴², M. Artuso⁶⁷, E. Aslanides¹², M. Atzeni⁶³, B. Audurier¹⁴, D. Bacher⁶², I. Bachiller Perea¹⁰, S. Bachmann²⁰, M. Bachmayer⁴⁸, J. J. Back⁵⁵, P. Baladron Rodriguez⁴⁵, V. Balagura¹⁴, W. Baldini²⁴, H. Bao⁷, J. Baptista de Souza Leite⁵⁹, M. Barbetti^{25,d}, I. R. Barbosa⁶⁸, R. J. Barlow⁶¹, M. Barnyakov²³, S. Barsuk¹³, W. Barter⁵⁷, M. Bartolini⁵⁴, J. Bartz⁶⁷, F. Baryshnikov⁴², J. M. Basels¹⁶, G. Bassi^{33,e}, B. Batsukh⁵, A. Bay⁴⁸, A. Beck⁵⁵, M. Becker¹⁸, F. Bedeschi³³, I. B. Bediaga², S. Belin⁴⁵, V. Bellee⁴⁹, K. Belous⁴², I. Belov²⁷, I. Belyaev³⁴, G. Benane¹², G. Bencivenni²⁶, E. Ben-Haim¹⁵, A. Berezhnoy⁴², R. Bernet⁴⁹, S. Bernet Andres⁴³, A. Bertolin³¹, C. Betancourt⁴⁹, F. Betti⁵⁷, J. Bex⁵⁴, I. A. Bezshyiko⁴⁹, J. Bhom³⁹, M. S. Bieker¹⁸, N. V. Biesuz²⁴, P. Billoir¹⁵, A. Biolchini³⁶, M. Birch⁶⁰, F. C. R. Bishop¹⁰, A. Bitadze⁶¹, A. Bizzeti¹⁰, T. Blake⁵⁵, F. Blanc⁴⁸, J. E. Blank¹⁸, S. Blusk⁶⁷, V. Bocharnikov⁴², J. A. Boelhaeve¹⁸, O. Boente Garcia¹⁴, T. Boettcher⁶⁴, A. Bohare⁵⁷, A. Boldyrev⁴², C. S. Bolognani⁷⁶, R. Bolzonella^{24,f}, N. Bondar⁴², F. Borgato^{31,47,g}, S. Borghi⁶¹, M. Borsato^{29,b}, J. T. Borsuk³⁹, S. A. Bouchiba⁴⁸, T. J. V. Bowcock⁵⁹, A. Boyer⁴⁷, C. Bozzi²⁴, M. J. Bradley⁶⁰, A. Brea Rodriguez⁴⁸, N. Breer¹⁸, J. Brodzicka³⁹, A. Brossa Gonzalo^{45,55,44,a}, J. Brown⁵⁹, D. Brundu³⁰, E. Buchanan⁵⁷, A. Buonaura⁴⁹, L. Buonincontri^{31,g}, A. T. Burke⁶¹, C. Burr⁴⁷, A. Butkevich⁴², J. S. Butter⁵⁴, J. Buytaert⁴⁷, W. Byczynski⁴⁷, S. Cadeddu³⁰, H. Cai⁷², R. Calabrese^{24,f}, S. Calderon Ramirez⁹, L. Calefice⁴⁴, S. Cali²⁶, M. Calvi^{29,b}, M. Calvo Gomez⁴³, P. Camargo Magalhaes^{2,h}, J. I. Cambon Bouzas⁴⁵, P. Campana²⁶, D. H. Campora Perez⁷⁶, A. F. Campoverde Quezada⁷, S. Capelli²⁹, L. Capriotti²⁴, R. Caravaca-Mora⁹, A. Carbone^{23,i}, L. Carcedo Salgado⁴⁵, R. Cardinale^{27,j}, A. Cardini³⁰, P. Carniti^{29,b}, L. Carus²⁰, A. Casais Vidal⁶³, R. Caspary²⁰, G. Casse⁵⁹, J. Castro Godinez⁹, M. Cattaneo⁴⁷, G. Cavallero^{24,47}, V. Cavallini^{24,f}, S. Celani²⁰, D. Cervenkov⁶², S. Cesare^{28,k}, A. J. Chadwick⁵⁹, I. Chahrour⁸⁰, M. Charles¹⁵, Ph. Charpentier⁴⁷, C. A. Chavez Barajas⁵⁹, M. Chefdeville¹⁰, C. Chen¹², S. Chen⁵, Z. Chen⁷, A. Chernov³⁹, S. Chernyshenko⁵¹, V. Chobanova⁷⁸, S. Cholak⁴⁸, M. Chrzaszcz³⁹, A. Chubykin⁴², V. Chulikov⁴², P. Ciambrone²⁶, X. Cid Vidal⁴⁵, G. Ciezarek⁴⁷, P. Cifra⁴⁷, P. E. L. Clarke⁵⁷, M. Clemencic⁴⁷, H. V. Cliff⁵⁴, J. Closier⁴⁷, C. Cocha Toapaxi²⁰, V. Coco⁴⁷, J. Cogan¹², E. Cogneras¹¹, L. Cojocariu⁴¹, P. Collins⁴⁷, T. Colombo⁴⁷, A. Comerma-Montells⁴⁴, L. Congedo²², A. Contu³⁰, N. Cooke⁵⁸, I. Corredoira⁴⁵, A. Correia¹⁵, G. Corti⁴⁷, J. J. Cottee Meldrum⁵³, B. Couturier⁴⁷, D. C. Craik⁴⁹, M. Cruz Torres^{2,l}, E. Curras Rivera⁴⁸, R. Currie⁵⁷, C. L. Da Silva⁶⁶, S. Dadabaev⁴², L. Dai⁶⁹, X. Dai⁶, E. Dall'Occo¹⁸, J. Dalseno⁴⁵, C. D'Ambrosio⁴⁷, J. Daniel¹¹, A. Danilina⁴², P. d'Argent²², A. Davidson⁵⁵, J. E. Davies⁶¹, A. Davis⁶¹, O. De Aguiar Francisco⁶¹, C. De Angelis^{30,m}, F. De Benedetti⁴⁷, J. de Boer³⁶, K. De Bruyn⁷⁵, S. De Capua⁶¹, M. De Cian^{20,47}, U. De Freitas Carneiro Da Graca^{2,n}, E. De Lucia²⁶

J. M. De Miranda², L. De Paula³, M. De Serio^{22,o}, P. De Simone²⁶, F. De Vellis¹⁸, J. A. de Vries⁷⁶, F. Debernardis²², D. Decamp¹⁰, V. Dedu¹², L. Del Buono¹⁵, B. Delaney⁶³, H.-P. Dembinski¹⁸, J. Deng⁸, V. Denysenko⁴⁹, O. Deschamps¹¹, F. Dettori^{30,m}, B. Dey⁷⁴, P. Di Nezza²⁶, I. Diachkov⁴², S. Didenko⁴², S. Ding⁶⁷, L. Dittmann²⁰, V. Dobishuk⁵¹, A. D. Docheva⁵⁸, C. Dong⁴, A. M. Donohoe²¹, F. Dordei³⁰, A. C. dos Reis², A. D. Dowling⁶⁷, W. Duan⁷⁰, P. Duda⁷⁷, M. W. Dudek³⁹, L. Dufour⁴⁷, V. Duk³², P. Durante⁴⁷, M. M. Duras⁷⁷, J. M. Durham⁶⁶, O. D. Durmus⁷⁴, A. Dziurda³⁹, A. Dzyuba⁴², S. Easo⁵⁶, E. Eckstein¹⁷, U. Egede¹, A. Egorychev⁴², V. Egorychev⁴², S. Eisenhardt⁵⁷, E. Ejopu⁶¹, L. Eklund⁷⁹, M. Elashri⁶⁴, J. Ellbracht¹⁸, S. Ely⁶⁰, A. Ene⁴¹, E. Eppele⁶⁴, J. Eschle⁶⁷, S. Esen²⁰, T. Evans⁶¹, F. Fabiano^{30,47,m}, L. N. Falcao², Y. Fan⁷, B. Fang⁷², L. Fantini^{32,p}, M. Faria⁴⁸, K. Farmer⁵⁷, D. Fazzini^{29,b}, L. Felkowski⁷⁷, M. Feng^{5,7}, M. Feo^{18,47}, M. Fernandez Gomez⁴⁵, A. D. Fernandez⁶⁵, F. Ferrari²³, F. Ferreira Rodrigues³, M. Ferrillo⁴⁹, M. Ferro-Luzzi⁴⁷, S. Filippov⁴², R. A. Fini²², M. Fiorini^{24,f}, K. L. Fischer⁶², D. S. Fitzgerald⁸⁰, C. Fitzpatrick⁶¹, F. Fleuret¹⁴, M. Fontana²³, L. F. Foreman⁶¹, R. Forty⁴⁷, D. Foulds-Holt⁵⁴, M. Franco Sevilla⁶⁵, M. Frank⁴⁷, E. Franzoso^{24,f}, G. Frau²⁰, C. Frei⁴⁷, D. A. Friday⁶¹, J. Fu⁷, Q. Fuehring¹⁸, Y. Fujii¹, T. Fulghesu¹⁵, E. Gabriel³⁶, G. Galati²², M. D. Galati³⁶, A. Gallas Torreira⁴⁵, D. Galli^{23,i}, S. Gambetta⁵⁷, M. Gandelman³, P. Gandini²⁸, B. Ganie⁶¹, H. Gao⁷, R. Gao⁶², Y. Gao⁸, Y. Gao⁶, Y. Gao⁸, M. Garau^{30,m}, L. M. Garcia Martin⁴⁸, P. Garcia Moreno⁴⁴, J. Garcia Pardiñas⁴⁷, K. G. Garg⁸, L. Garrido⁴⁴, C. Gaspar⁴⁷, R. E. Geertsema³⁶, L. L. Gerken¹⁸, E. Gersabeck⁶¹, M. Gersabeck⁶¹, T. Gershon⁵⁵, Z. Ghorbanimoghaddam⁵³, L. Giambastiani^{31,g}, F. I. Giasemis^{15,q}, V. Gibson⁵⁴, H. K. Giemza⁴⁰, A. L. Gilman⁶², M. Giovannetti²⁶, A. Gioventù⁴⁴, P. Gironella Gironell⁴⁴, C. Giugliano^{24,f}, M. A. Giza³⁹, E. L. Gkougkousis⁶⁰, F. C. Glaser^{13,20}, V. V. Gligorov^{15,47}, C. Göbel⁶⁸, E. Golobardes⁴³, D. Golubkov⁴², A. Golutvin^{60,42,47}, A. Gomes^{2,a,r}, S. Gomez Fernandez⁴⁴, F. Goncalves Abrantes⁶², M. Goncerz³⁹, G. Gong⁴, J. A. Gooding¹⁸, I. V. Gorelov⁴², C. Gotti²⁹, J. P. Grabowski¹⁷, L. A. Granado Cardoso⁴⁷, E. Graugés⁴⁴, E. Graverini^{48,s}, L. Grazette⁵⁵, G. Graziani¹, A. T. Grecu⁴¹, L. M. Greeven³⁶, N. A. Grieser⁶⁴, L. Grillo⁵⁸, S. Gromov⁴², C. Gu¹⁴, M. Guarise²⁴, M. Guittiere¹³, V. Guliaeva⁴², P. A. Günther²⁰, A.-K. Guseinov⁴⁸, E. Gushchin⁴², Y. Guz^{6,42,47}, T. Gys⁴⁷, K. Habermann¹⁷, T. Hadavizadeh¹, C. Hadjivasilou⁶⁵, G. Haefeli⁴⁸, C. Haen⁴⁷, J. Haimberger⁴⁷, M. Hajheidari⁴⁷, M. M. Halvorsen⁴⁷, P. M. Hamilton⁶⁵, J. Hammerich⁵⁹, Q. Han⁸, X. Han²⁰, S. Hansmann-Menzemer²⁰, L. Hao⁷, N. Harnew⁶², M. Hartmann¹³, J. He^{7,t}, F. Hemmer⁴⁷, C. Henderson⁶⁴, R. D. L. Henderson^{1,55}, A. M. Hennequin⁴⁷, K. Hennessy⁵⁹, L. Henry⁴⁸, J. Herd⁶⁰, P. Herrero Gascon²⁰, J. Heuel¹⁶, A. Hicheur³, G. Hijano Mendizabal⁴⁹, D. Hill⁴⁸, S. E. Hollitt¹⁸, J. Horswill⁶¹, R. Hou⁸, Y. Hou¹¹, N. Howarth⁵⁹, J. Hu²⁰, J. Hu⁷⁰, W. Hu⁶, X. Hu⁴, W. Huang⁷, W. Hulsbergen³⁶, R. J. Hunter⁵⁵, M. Hushchyn⁴², D. Hutchcroft⁵⁹, D. Ilin⁴², P. Ilten⁶⁴, A. Inglessi⁴², A. Iniukhin⁴², A. Ishteev⁴², K. Ivshin⁴², R. Jacobsson⁴⁷, H. Jage¹⁶, S. J. Jaimes Elles^{46,73}, S. Jakobsen⁴⁷, E. Jans³⁶, B. K. Jashal⁴⁶, A. Jawahery^{65,47}, V. Jevtic¹⁸, E. Jiang⁶⁵, X. Jiang^{5,7}, Y. Jiang⁷, Y. J. Jiang⁶, M. John⁶², D. Johnson⁵², C. R. Jones⁵⁴, T. P. Jones⁵⁵, S. Joshi⁴⁰, B. Jost⁴⁷, N. Jurik⁴⁷, I. Juszczak³⁹, D. Kaminaris⁴⁸, S. Kandybei⁵⁰, Y. Kang⁴, C. Kar¹¹, M. Karacson⁴⁷, D. Karpenkov⁴², A. Kauniskangas⁴⁸, J. W. Kautz⁶⁴, F. Keizer⁴⁷, M. Kenzie⁵⁴, T. Ketel³⁶, B. Khanji⁶⁷, A. Kharisova⁴², S. Kholodenko^{33,47}, G. Khreich¹³, T. Kirm¹⁶, V. S. Kirsebom^{29,b}, O. Kitouni⁶³, S. Klaver³⁷, N. Kleijne^{33,e}, K. Klimaszewski⁴⁰, M. R. Kmiec⁴⁰, S. Kolliiev⁵¹, L. Kolk¹⁸, A. Konoplyannikov⁴², P. Kopciewicz^{38,47}, P. Koppenburg³⁶, M. Korolev⁴², I. Kostiuik³⁶, O. Kot⁵¹, S. Kotriakhova⁴², A. Kozachuk⁴², P. Kravchenko⁴², L. Kravchuk⁴², M. Kreps⁵⁵, P. Krokovny⁴², W. Krupa⁶⁷, W. Krzemien⁴⁰, O. K. Kshyvanskyi⁵¹, J. Kubat²⁰, S. Kubis⁷⁷, M. Kucharczyk³⁹, V. Kudryavtsev⁴², E. Kulikova⁴², A. Kupsc⁷⁹, B. K. Kutsenko¹², D. Lacarrere⁴⁷, A. Lai³⁰, A. Lampis³⁰, D. Lancierini⁵⁴, C. Landesa Gomez⁴⁵, J. J. Lane¹, R. Lane⁵³, C. Langenbruch²⁰, J. Langer¹⁸, O. Lantwin⁴², T. Latham⁵⁵, F. Lazzari^{33,s}, C. Lazzeroni⁵², R. Le Gac¹², R. Lefèvre¹¹, A. Leflat⁴², S. Legotin⁴², M. Lehuraux⁵⁵, E. Lemos Cid⁴⁷, O. Leroy¹², T. Lesiak³⁹, B. Leverington²⁰, A. Li⁴, H. Li⁷⁰, K. Li⁸, L. Li⁶¹, P. Li⁴⁷, P.-R. Li⁷¹, Q. Li^{5,7}, S. Li⁸, T. Li^{5,u}, T. Li⁷⁰, Y. Li⁸, Y. Li⁵, Z. Lian⁴, X. Liang⁶⁷, S. Liralon⁴⁶, C. Lin⁷, T. Lin⁵⁶, R. Lindner⁴⁷, V. Lisovskyi⁴⁸, R. Litvinov³⁰, F. L. Liu¹, G. Liu⁷⁰, K. Liu⁷¹, S. Liu^{5,7}, Y. Liu⁵⁷, Y. Liu⁷¹, Y. L. Liu⁶⁰, A. Lobo Salvia⁴⁴, A. Loi³⁰, J. Lomba Castro⁴⁵, T. Long⁵⁴, J. H. Lopes³, A. Lopez Huertas⁴⁴, S. López Soliño⁴⁵, C. Lucarelli^{25,d}, D. Lucchesi^{31,g}, M. Lucio Martinez⁷⁶, V. Lukashenko^{36,51}, Y. Luo⁶, A. Lupato³¹, E. Luppi^{24,f}, K. Lynch²¹, X.-R. Lyu⁷, G. M. Ma⁴, R. Ma⁷, S. Maccolini¹⁸, F. Machefert¹³, F. Maciuc⁴¹, B. Mack⁶⁷, I. Mackay⁶², L. M. Mackey⁶⁷, L. R. Madhan Mohan⁵⁴

M. J. Madurai⁵², A. Maevskiy⁴², D. Magdalinski³⁶, D. Maisuzenko⁴², M. W. Majewski³⁸, J. J. Malczewski³⁹, S. Malde⁶², L. Malentacca⁴⁷, A. Malinin⁴², T. Maltsev⁴², G. Manca^{30,m}, G. Mancinelli¹², C. Mancuso^{28,13,k}, R. Manera Escalero⁴⁴, D. Manuzzi²³, D. Marangotto^{28,k}, J. F. Marchand¹⁰, R. Marchevski⁴⁸, U. Marconi²³, S. Mariani⁴⁷, C. Marin Benito⁴⁴, J. Marks²⁰, A. M. Marshall⁵³, G. Martelli^{32,p}, G. Martellotti³⁴, L. Martinazzoli⁴⁷, M. Martinelli^{29,b}, D. Martinez Santos⁴⁵, F. Martinez Vidal⁴⁶, A. Massafferri², R. Matev⁴⁷, A. Mathad⁴⁷, V. Matiunin⁴², C. Matteuzzi⁶⁷, K. R. Mattioli¹⁴, A. Mauri⁶⁰, E. Maurice¹⁴, J. Mauricio⁴⁴, P. Mayencourt⁴⁸, M. Mazurek⁴⁰, M. McCann⁶⁰, L. McConnell²¹, T. H. McGrath⁶¹, N. T. McHugh⁵⁸, A. McNab⁶¹, R. McNulty²¹, B. Meadows⁶⁴, G. Meier¹⁸, D. Melnychuk⁴⁰, F. M. Meng⁴, M. Merk^{36,76}, A. Merli⁴⁸, L. Meyer Garcia⁶⁵, D. Miao^{5,7}, H. Miao⁷, M. Mikhasenko^{17,v}, D. A. Milanese⁷³, A. Minotti^{29,b}, E. Minucci⁶⁷, T. Miralles¹¹, B. Mitreska¹⁸, D. S. Mitzel¹⁸, A. Modak⁵⁶, A. Mödden¹⁸, R. A. Mohammed⁶², R. D. Moise¹⁶, S. Mokhnenko⁴², T. Mombächer⁴⁷, M. Monk^{55,1}, S. Monteil¹¹, A. Morcillo Gomez⁴⁵, G. Morello²⁶, M. J. Morello^{33,e}, M. P. Morgenthaler²⁰, A. B. Morris⁴⁷, A. G. Morris¹², R. Mountain⁶⁷, H. Mu⁴, Z. M. Mu⁶, E. Muhammad⁵⁵, F. Muheim⁵⁷, M. Mulder⁷⁵, K. Müller⁴⁹, F. Muñoz-Rojas⁹, R. Murta⁶⁰, P. Naik⁵⁹, T. Nakada⁴⁸, R. Nandakumar⁵⁶, T. Nanut⁴⁷, I. Nasteva³, M. Needham⁵⁷, N. Neri^{28,k}, S. Neubert¹⁷, N. Neufeld⁴⁷, P. Neustroev⁴², J. Nicolini^{18,13}, D. Nicotra⁷⁶, E. M. Niel⁴⁸, N. Nikitin⁴², P. Nogarolli³, P. Nogga¹⁷, N. S. Nolte⁶³, C. Normand⁵³, J. Novoa Fernandez⁴⁵, G. Nowak⁶⁴, C. Nunez⁸⁰, H. N. Nur⁵⁸, A. Oblakowska-Mucha³⁸, V. Obraztsov⁴², T. Oeser¹⁶, S. Okamura^{24,47,f}, A. Okhotnikov⁴², O. Okhrimenko⁵¹, R. Oldeman^{30,m}, F. Oliva⁵⁷, M. Olocco¹⁸, C. J. G. Onderwater⁷⁶, R. H. O'Neil⁵⁷, J. M. Otalora Goicochea³, P. Owen⁴⁹, A. Oyanguren⁴⁶, O. Ozcelik⁵⁷, K. O. Padeken¹⁷, B. Pagare⁵⁵, P. R. Pais²⁰, T. Pajero⁴⁷, A. Palano²², M. Palutan²⁶, G. Panshin⁴², L. Paolucci⁵⁵, A. Papanestis⁵⁶, M. Pappagallo^{22,o}, L. L. Pappalardo^{24,f}, C. Pappenheimer⁶⁴, C. Parkes⁶¹, B. Passalacqua²⁴, G. Passaleva²⁵, D. Passaro^{33,e}, A. Pastore²², M. Patel⁶⁰, J. Patoc⁶², C. Patrignani^{23,i}, A. Paul⁶⁷, C. J. Pawley⁷⁶, A. Pellegrino³⁶, J. Peng^{5,7}, M. Pepe Altarelli²⁶, S. Perazzini²³, D. Pereima⁴², H. Pereira Da Costa⁶⁶, A. Pereiro Castro⁴⁵, P. Perret¹¹, A. Perro⁴⁷, K. Petridis⁵³, A. Petrolini^{27,j}, J. P. Pfaller⁶⁴, H. Pham⁶⁷, L. Pica^{33,e}, M. Piccini³², B. Pietrzyk¹⁰, G. Pietrzyk¹³, D. Pinci³⁴, F. Pisani⁴⁷, M. Pizzichemi^{29,b}, V. Placinta⁴¹, M. Plo Casasus⁴⁵, F. Polci^{15,47}, M. Poli Lener²⁶, A. Poluektov¹², N. Polukhina⁴², I. Polyakov⁴⁷, E. Polycarpo³, S. Ponce⁴⁷, D. Popov⁷, S. Poslavskii⁴², K. Prasanth⁵⁷, C. Prouve⁴⁵, V. Pugatch⁵¹, G. Punzi^{33,s}, S. Qasim⁴⁹, W. Qian⁷, N. Qin⁴, S. Qu⁴, R. Quagliani⁴⁷, R. I. Rabadan Trejo⁵⁵, J. H. Rademacker⁵³, M. Rama³³, M. Ramírez García⁸⁰, V. Ramos De Oliveira⁶⁸, M. Ramos Pernas⁵⁵, M. S. Rangel³, F. Ratnikov⁴², G. Raven³⁷, M. Rebollo De Miguel⁴⁶, F. Redi^{28,w}, J. Reich⁵³, F. Reiss⁶¹, Z. Ren⁷, P. K. Resmi⁶², R. Ribatti^{33,e}, G. R. Ricart^{14,81}, D. Riccardi^{33,e}, S. Ricciardi⁵⁶, K. Richardson⁶³, M. Richardson-Slipper⁵⁷, K. Rinnert⁵⁹, P. Robbe¹³, G. Robertson⁵⁸, E. Rodrigues⁵⁹, E. Rodriguez Fernandez⁴⁵, J. A. Rodriguez Lopez⁷³, E. Rodriguez Rodriguez⁴⁵, A. Rogovskiy⁵⁶, D. L. Rolf⁴⁷, P. Roloff⁴⁷, V. Romanovskiy⁴², M. Romero Lamas⁴⁵, A. Romero Vidal⁴⁵, G. Romolini²⁴, F. Ronchetti⁴⁸, M. Rotondo²⁶, S. R. Roy²⁰, M. S. Rudolph⁶⁷, T. Ruf⁴⁷, M. Ruiz Diaz²⁰, R. A. Ruiz Fernandez⁴⁵, J. Ruiz Vidal^{79,x}, A. Ryzhikov⁴², J. Ryzka³⁸, J. J. Saavedra-Arias⁹, J. J. Saborido Silva⁴⁵, R. Sadek¹⁴, N. Sagidova⁴², D. Sahoo⁷⁴, N. Sahoo⁵², B. Saitta^{30,m}, M. Salomoni^{29,47,b}, C. Sanchez Gras³⁶, I. Sanderswood⁴⁶, R. Santacesaria³⁴, C. Santamarina Rios⁴⁵, M. Santimaria^{26,47}, L. Santoro², E. Santovetti³⁵, A. Saputi^{24,47}, D. Saranin⁴², G. Sarpis⁵⁷, M. Sarpis⁶¹, C. Satriano^{34,y}, A. Satta³⁵, M. Saur⁶, D. Savrina⁴², H. Sazak¹⁶, L. G. Scantlebury Smead⁶², A. Scarabotto¹⁸, S. Schael¹⁶, S. Scherl⁵⁹, M. Schiller⁵⁸, H. Schindler⁴⁷, M. Schmelling¹⁹, B. Schmidt⁴⁷, S. Schmitt¹⁶, H. Schmitz¹⁷, O. Schneider⁴⁸, A. Schopper⁴⁷, N. Schulte¹⁸, S. Schulte⁴⁸, M. H. Schune¹³, R. Schwemmer⁴⁷, G. Schwering¹⁶, B. Sciascia²⁶, A. Sciuccati⁴⁷, S. Sellam⁴⁵, A. Semennikov⁴², T. Senger⁴⁹, M. Senghi Soares³⁷, A. Sergi²⁷, N. Serra⁴⁹, L. Sestini³¹, A. Seuthe¹⁸, Y. Shang⁶, D. M. Shangase⁸⁰, M. Shapkin⁴², R. S. Sharma⁶⁷, I. Shchemerov⁴², L. Shchutska⁴⁸, T. Shears⁵⁹, L. Shekhtman⁴², Z. Shen⁶, S. Sheng^{5,7}, V. Shevchenko⁴², B. Shi⁷, Q. Shi⁷, E. B. Shields^{29,b}, Y. Shimizu¹³, E. Shmanin⁴², R. Shorkin⁴², J. D. Shupperd⁶⁷, R. Silva Coutinho⁶⁷, G. Simi^{31,g}, S. Simone^{22,o}, N. Skidmore⁵⁵, T. Skwarnicki⁶⁷, M. W. Slater⁵², J. C. Smallwood⁶², E. Smith⁶³, K. Smith⁶⁶, M. Smith⁶⁰, A. Snoch³⁶, L. Soares Lavoura⁵⁷, M. D. Sokoloff⁶⁴, F. J. P. Soler⁵⁸, A. Solomin^{42,53}, A. Solovov⁴², I. Solovyev⁴², R. Song¹, Y. Song⁴⁸, Y. Song⁴, Y. S. Song⁶, F. L. Souza De Almeida⁶⁷, B. Souza De Paula³, E. Spadaro Norella^{28,k}, E. Spedicato²³, J. G. Speer¹⁸, E. Spiridenkov⁴², P. Spradlin⁵⁸, V. Sriskaran⁴⁷, F. Stagni⁴⁷, M. Stahl⁴⁷, S. Stahl⁴⁷

S. Stanislaus⁶², E. N. Stein⁴⁷, O. Steinkamp⁴⁹, O. Stenyakin⁴², H. Stevens¹⁸, D. Strelalina⁴², Y. Su⁷, F. Suljik⁶², J. Sun³⁰, L. Sun⁷², Y. Sun⁶⁵, D. Sundfeld², W. Sutcliffe⁴⁹, P. N. Swallow⁵², F. Swystun⁵⁴, A. Szabelski⁴⁰, T. Szumlak³⁸, Y. Tan⁴, M. D. Tat⁶², A. Terentev⁴², F. Terzuoli^{33,z}, F. Teubert⁴⁷, E. Thomas⁴⁷, D. J. D. Thompson⁵², H. Tilquin⁶⁰, V. Tisserand¹¹, S. T'Jampens¹⁰, M. Tobin⁵, L. Tomassetti^{24,f}, G. Tonani^{28,47,k}, X. Tong⁶, D. Torres Machado², L. Toscano¹⁸, D. Y. Tou⁴, C. Tripll⁴³, G. Tuci²⁰, N. Tuning³⁶, L. H. Uecker²⁰, A. Ukleja³⁸, D. J. Unverzagt²⁰, E. Ursov⁴², A. Usachov³⁷, A. Ustyuzhanin⁴², U. Uwer²⁰, V. Vagnoni²³, A. Valassi⁴⁷, G. Valenti²³, N. Valls Canudas⁴⁷, H. Van Hecke⁶⁶, E. van Herwijnen⁶⁰, C. B. Van Hulse^{45,aa}, R. Van Laak⁴⁸, M. van Veghel³⁶, G. Vasquez⁴⁹, R. Vazquez Gomez⁴⁴, P. Vazquez Regueiro⁴⁵, C. Vázquez Sierra⁴⁵, S. Vecchi²⁴, J. J. Velthuis⁵³, M. Veltri^{25,bb}, A. Venkateswaran⁴⁸, M. Vesterinen⁵⁵, M. Vieites Diaz⁴⁷, X. Vilasis-Cardona⁴³, E. Vilella Figueras⁵⁹, A. Villa²³, P. Vincent¹⁵, F. C. Volle⁵², D. vom Bruch¹², N. Voropaev⁴², K. Vos⁷⁶, G. Vouters^{10,47}, C. Vrahas⁵⁷, J. Wagner¹⁸, J. Walsh³³, E. J. Walton^{1,55}, G. Wan⁶, C. Wang²⁰, G. Wang⁸, J. Wang⁶, J. Wang⁵, J. Wang⁴, J. Wang⁷², M. Wang²⁸, N. W. Wang⁷, R. Wang⁵³, X. Wang⁸, X. Wang⁷⁰, X. W. Wang⁶⁰, Z. Wang¹³, Z. Wang⁴, Z. Wang²⁸, J. A. Ward^{55,1}, M. Waterlaet⁴⁷, N. K. Watson⁵², D. Websdale⁶⁰, Y. Wei⁶, J. Wendel⁷⁸, B. D. C. Westhenry⁵³, D. J. White⁶¹, M. Whitehead⁵⁸, A. R. Wiederhold⁵⁵, D. Wiedner¹⁸, G. Wilkinson⁶², M. K. Wilkinson⁶⁴, M. Williams⁶³, M. R. J. Williams⁵⁷, R. Williams⁵⁴, F. F. Wilson⁵⁶, W. Wislicki⁴⁰, M. Witek³⁹, L. Witola²⁰, C. P. Wong⁶⁶, G. Wormser¹³, S. A. Wotton⁵⁴, H. Wu⁶⁷, J. Wu⁸, Y. Wu⁶, Z. Wu⁷, K. Wyllie⁴⁷, S. Xian⁷⁰, Z. Xiang⁵, Y. Xie⁸, A. Xu³³, J. Xu⁷, L. Xu⁴, L. Xu⁴, M. Xu⁵⁵, Z. Xu¹¹, Z. Xu⁷, Z. Xu⁵, D. Yang⁴, K. Yang⁶⁰, S. Yang⁷, X. Yang⁶, Y. Yang^{27,j}, Z. Yang⁶, Z. Yang⁶⁵, V. Yeroshenko¹³, H. Yeung⁶¹, H. Yin⁸, C. Y. Yu⁶, J. Yu⁶⁹, X. Yuan⁵, E. Zaffaroni⁴⁸, M. Zavertyaev¹⁹, M. Zdybal³⁹, C. Zeng^{5,7}, M. Zeng⁴, C. Zhang⁶, D. Zhang⁸, J. Zhang⁷, L. Zhang⁴, S. Zhang⁶⁹, S. Zhang⁶, Y. Zhang⁶, Y. Z. Zhang⁴, Y. Zhao²⁰, A. Zharkova⁴², A. Zhelezov²⁰, X. Z. Zheng⁴, Y. Zheng⁷, T. Zhou⁶, X. Zhou⁸, Y. Zhou⁷, V. Zhovkovska⁵⁵, L. Z. Zhu⁷, X. Zhu⁴, X. Zhu⁸, V. Zhukov¹⁶, J. Zhuo⁴⁶, Q. Zou^{5,7}, D. Zuliani^{31,g} and G. Zunica⁴⁸

(LHCb Collaboration)

¹School of Physics and Astronomy, Monash University, Melbourne, Australia

²Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

³Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

⁴Center for High Energy Physics, Tsinghua University, Beijing, China

⁵Institute of High Energy Physics (IHEP), Beijing, China

⁶School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁷University of Chinese Academy of Sciences, Beijing, China

⁸Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁹Consejo Nacional de Rectores (CONARE), San Jose, Costa Rica

¹⁰Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

¹¹Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

¹²Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

¹³Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

¹⁴Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

¹⁵LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

¹⁶I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁷Universität Bonn—Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany

¹⁸Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹⁹Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

²⁰Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

²¹School of Physics, University College Dublin, Dublin, Ireland

²²INFN Sezione di Bari, Bari, Italy

²³INFN Sezione di Bologna, Bologna, Italy

²⁴INFN Sezione di Ferrara, Ferrara, Italy

²⁵INFN Sezione di Firenze, Firenze, Italy

²⁶INFN Laboratori Nazionali di Frascati, Frascati, Italy

- ²⁷*INFN Sezione di Genova, Genova, Italy*
- ²⁸*INFN Sezione di Milano, Milano, Italy*
- ²⁹*INFN Sezione di Milano-Bicocca, Milano, Italy*
- ³⁰*INFN Sezione di Cagliari, Monserrato, Italy*
- ³¹*INFN Sezione di Padova, Padova, Italy*
- ³²*INFN Sezione di Perugia, Perugia, Italy*
- ³³*INFN Sezione di Pisa, Pisa, Italy*
- ³⁴*INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³⁵*INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³⁶*Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³⁷*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ³⁸*AGH—University of Krakow, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ³⁹*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ⁴⁰*National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ⁴¹*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ⁴²*Affiliated with an institute covered by a cooperation agreement with CERN*
- ⁴³*DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁴⁴*ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁵*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁶*Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia—CSIC, Valencia, Spain*
- ⁴⁷*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴⁸*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁹*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁵⁰*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁵¹*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁵²*School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*
- ⁵³*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁵⁴*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁵*Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁶*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵⁷*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁸*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁹*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁶⁰*Imperial College London, London, United Kingdom*
- ⁶¹*Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁶²*Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁶³*Massachusetts Institute of Technology, Cambridge, Massachusetts, USA*
- ⁶⁴*University of Cincinnati, Cincinnati, Ohio, USA*
- ⁶⁵*University of Maryland, College Park, Maryland, USA*
- ⁶⁶*Los Alamos National Laboratory (LANL), Los Alamos, New Mexico, USA*
- ⁶⁷*Syracuse University, Syracuse, New York, USA*
- ⁶⁸*Pontificia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil
(associated with Universidade Federal do Rio de Janeiro (UFRJ),
Rio de Janeiro, Brazil)*
- ⁶⁹*School of Physics and Electronics, Hunan University, Changsha City, China
(associated with Institute of Particle Physics, Central China Normal University,
Wuhan, Hubei, China)*
- ⁷⁰*Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of
Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China
(associated with Center for High Energy Physics, Tsinghua University, Beijing, China)*
- ⁷¹*Lanzhou University, Lanzhou, China
(associated with Institute Of High Energy Physics (IHEP), Beijing, China)*
- ⁷²*School of Physics and Technology, Wuhan University, Wuhan, China
(associated with Center for High Energy Physics, Tsinghua University, Beijing, China)*
- ⁷³*Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia
(associated with LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3,
Paris, France)*
- ⁷⁴*Eotvos Lorand University, Budapest, Hungary
(associated with European Organization for Nuclear Research (CERN), Geneva, Switzerland)*

⁷⁵*Van Swinderen Institute, University of Groningen, Groningen, Netherlands
(associated with Nikhef National Institute for Subatomic Physics,
Amsterdam, Netherlands)*

⁷⁶*Universiteit Maastricht, Maastricht, Netherlands
(associated with Nikhef National Institute for Subatomic Physics,
Amsterdam, Netherlands)*

⁷⁷*Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland
(associated with Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences,
Kraków, Poland)*

⁷⁸*Universidade da Coruña, A Coruna, Spain
(associated with DS4DS, La Salle, Universitat Ramon Llull,
Barcelona, Spain)*

⁷⁹*Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden
(associated with School of Physics and Astronomy, University of Glasgow,
Glasgow, United Kingdom)*

⁸⁰*University of Michigan, Ann Arbor, Michigan, USA
(associated with Syracuse University, Syracuse, New York, USA)*

⁸¹*Departement de Physique Nucleaire (SPhN), Gif-Sur-Yvette, France*

^aDeceased.

^bAlso at Università degli Studi di Milano-Bicocca, Milano, Italy.

^cAlso at Università di Roma Tor Vergata, Roma, Italy.

^dAlso at Università di Firenze, Firenze, Italy.

^eAlso at Scuola Normale Superiore, Pisa, Italy.

^fAlso at Università di Ferrara, Ferrara, Italy.

^gAlso at Università di Padova, Padova, Italy.

^hAlso at Facultad de Ciencias Fisicas, Madrid, Spain.

ⁱAlso at Università di Bologna, Bologna, Italy.

^jAlso at Università di Genova, Genova, Italy.

^kAlso at Università degli Studi di Milano, Milano, Italy.

^lAlso at Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras.

^mAlso at Università di Cagliari, Cagliari, Italy.

ⁿAlso at Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Rio De Janeiro, Brazil.

^oAlso at Università di Bari, Bari, Italy.

^pAlso at Università di Perugia, Perugia, Italy.

^qAlso at LIP6, Sorbonne Université, Paris, France.

^rAlso at Universidade de Brasília, Brasília, Brazil.

^sAlso at Università di Pisa, Pisa, Italy.

^tAlso at Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China.

^uAlso at School of Physics and Electronics, Henan University, Kaifeng, China.

^vAlso at Excellence Cluster ORIGINS, Munich, Germany.

^wAlso at Università di Bergamo, Bergamo, Italy.

^xAlso at Department of Physics/Division of Particle Physics, Lund, Sweden.

^yAlso at Università della Basilicata, Potenza, Italy.

^zAlso at Università di Siena, Siena, Italy.

^{aa}Also at Universidad de Alcalá, Alcalá de Henares, Spain.

^{bb}Also at Università di Urbino, Urbino, Italy.