

Serotonergic Signaling Controls Input-Specific Synaptic Plasticity at Striatal Circuits

Highlights

- 5-HT signaling controls t-LTD in striatal projection neurons of the direct pathway
- 5-HT signaling shapes dendritic Ca^{2+} signals
- 5-HT-mediated regulation of t-LTD biases thalamostriatal synapses
- Gating of t-LTD requires dampened activity of the 5-HT₄R subtype, which modulates BK channel function

Authors

Anna Cavaccini, Marta Gritti, Andrea Giorgi, ..., Alessandro Gozzi, Massimo Pasqualetti, Raffaella Tonini

Correspondence

raffaella.tonini@iit.it

In Brief

Cavaccini and Gritti et al. combine chemogenetic and optogenetic approaches to show that serotonergic signaling provides a control mechanism of synaptic plasticity at thalamic inputs to the striatum.



Serotonergic Signaling Controls Input-Specific Synaptic Plasticity at Striatal Circuits

Anna Cavaccini,^{1,8} Marta Gritti,^{1,8} Andrea Giorgi,^{2,7} Andrea Locarno,¹ Nicolas Heck,³ Sara Migliarini,² Alice Bertero,^{2,7} Maddalena Mereu,⁴ Giulia Margiani,⁵ Massimo Trusel,¹ Tiziano Catelani,⁶ Roberto Marotta,⁶ Maria Antonietta De Luca,⁵ Jocelyne Caboche,³ Alessandro Gozzi,⁷ Massimo Pasqualetti,^{2,7} and Raffaella Tonini^{1,*}

¹Neuromodulation of Cortical and Subcortical Circuits Laboratory, Neuroscience and Brain Technologies Department, Istituto Italiano di Tecnologia, Genova, Italy

²Department of Biology, Unit of Cell and Developmental Biology, University of Pisa, Pisa, Italy

³Neurosciences Paris Seine, Institut de Biologie Paris Seine, CNRS, INSERM, Sorbonne Université, Paris, France

⁴Department of Pharmaceutical and Pharmacological Sciences, University of Padua, Padua, Italy

⁵Department of Biomedical Sciences, Cittadella Universitaria di Monserrato, University of Cagliari, Cagliari, Italy

⁶Electron Microscopy Facility, Istituto Italiano di Tecnologia, Genova, Italy

⁷Functional Neuroimaging Laboratory, Center for Neuroscience and Cognitive Systems, Istituto Italiano di Tecnologia, Rovereto, Italy

⁸Co-first authors

*Correspondence: raffaella.tonini@iit.it

<https://doi.org/10.1016/j.neuron.2018.04.008>

SUMMARY

Monoaminergic modulation of cortical and thalamic inputs to the dorsal striatum (DS) is crucial for reward-based learning and action control. While dopamine has been extensively investigated in this context, the synaptic effects of serotonin (5-HT) have been largely unexplored. Here, we investigated how serotonergic signaling affects associative plasticity at glutamatergic synapses on the striatal projection neurons of the direct pathway (dSPNs). Combining chemogenetic and optogenetic approaches reveals that impeding serotonergic signaling preferentially gates spike-timing-dependent long-term depression (t-LTD) at thalamostriatal synapses. This t-LTD requires dampened activity of the 5-HT₄ receptor subtype, which we demonstrate controls dendritic Ca²⁺ signals by regulating BK channel activity, and which preferentially localizes at the dendritic shaft. The synaptic effects of 5-HT signaling at thalamostriatal inputs provide insights into how changes in serotonergic levels associated with behavioral states or pathology affect striatal-dependent processes.

INTRODUCTION

The dorsal striatum (DS) of the basal ganglia (BG) plays a critical role in voluntary movement, learning, and motivation. It also represents the primary site of dysfunction in psychomotor diseases such as Parkinson's disease (PD), drug addiction, and obsessive-compulsive disorders (OCDs). The DS receives glutamatergic excitatory inputs from the cortex and thalamus. These inputs converge on striatal projection neurons (SPNs) of the direct (dSPN) and indirect (iSPN) pathways, which exert opposing con-

trol over motor output and hedonic states. While dSPNs stimulate movement, reinforcement, and reward, iSPNs inhibit movement and mediate punishment and aversion (Kraavitz et al., 2010, 2012; Tecuapetla et al., 2016; Vicente et al., 2016).

It is increasingly clear that encoding of action selection at striatal circuits requires regulation of SPN activity and plasticity (Kraavitz et al., 2012; O'Hare et al., 2016; Sippy et al., 2015). While the neuromodulatory signaling mechanisms controlling iSPN synaptic plasticity, especially long-term-depression (LTD), have been studied extensively, identifying parallel mechanisms in dSPNs has proven more difficult (Zhai et al., 2018). One of the most important neuromodulatory inputs to the striatum is dense dopaminergic innervation from the substantia nigra (Tritsch and Sabatini, 2012). In particular, dSPNs expressing the Gs-coupled D1 dopamine (DA) receptors (D1R) are stimulated by dopaminergic signaling. In the presence of behaviorally relevant stimuli, DA is thought to reinforce motor actions that result in reward outcomes through the potentiation of corticostriatal synapses formed on dSPNs. Recent evidence indicates that DA depletion specifically weakens thalamic connections to dSPNs, but not to cortical afferents to these neurons (Parker et al., 2016). This weakening, which does not occur in iSPNs, is directly associated with motor abnormalities.

In addition to dopaminergic afferents, serotonergic inputs from the medial and dorsal raphe nucleus (DRN) modulate the activity of SPNs, affecting striatal function and consequently behaviors, including motor and motivational goal-directed behavior (Mathur and Lovinger, 2012). Serotonergic neurons of the DRN fire coincidentally with the initiation and termination of selected voluntary movements, whereas reduction in DRN activity to baseline or below is associated with pauses in motor sequences (Fornal et al., 1996; Jacobs and Fornal, 1999). The evidence that striatal serotonin (5-HT) influences action control is consistent with the dysfunction of the 5-HT system described in OCD and with the use of selective serotonin reuptake inhibitors (SSRIs) for treatment of this disorder (Goddard et al., 2008).

In the DS, 5-HT acts on diverse receptors on both presynaptic and postsynaptic circuit elements (Mathur and Lovinger, 2012).



Depending on their localization, striatal 5-HT receptors initiate transduction through several Gi/o-, Gq/11-, and Gs protein-dependent and -independent pathways (Hannon and Hoyer, 2008). The specific responses to 5-HT may also be contingent upon crosstalk with other receptor types. In dSPNs, the temporal, combinational logic of serotonergic and D1R-mediated signaling may shape the net synaptic output, depending on the relative balance of these two neuromodulators. It has been reported that increased concentrations of 5-HT in the striatum reduce glutamate release at corticostriatal afferents by activating presynaptic Gi-coupled 5-HT_{1b} receptors (Mathur et al., 2011). 5-HT can also facilitate the release of DA from dopaminergic projections (Navailles and De Deurwaerdère, 2011). Despite their potential implications for BG function and for pathological conditions characterized by hypo-serotonergic states, the synaptic effects of reduced striatal 5-HT signaling remain unexplored. Neither is it known whether there is any input specificity for 5-HT modulation at corticostriatal and thalamostriatal synapses.

In the current study, we investigated the effect of impeding 5-HT release, and thereby serotonergic signaling, on timing-dependent synaptic plasticity in dSPNs. This allowed us to describe cellular mechanisms underlying the input-specific impact of 5-HT at thalamostriatal synapses.

RESULTS

Chemogenetic Inhibition of 5-HT Release Gates t-LTD at Glutamatergic Afferents to dSPNs

To determine how plasticity is modulated at glutamatergic inputs on dSPNs of the dorsolateral striatum (DLS), we recorded excitatory postsynaptic potentials (EPSPs) following intrastriatal stimulation. Intrastriatal stimulation likely activates cortical and thalamic afferents as well as dopaminergic and serotonergic fibers (Figure 1) (Shen et al., 2015; Threlfell et al., 2012; Wu et al., 2015). dSPNs were selected through their negative resting membrane potential and firing properties (Kreitzer and Malenka, 2007; Nazzaro et al., 2012). After recording, dSPN cell identity was confirmed by immunostaining for the specific marker substance P, and the lack of immunoreactivity for the iSPN marker A2A (Figure 1A) (Nazzaro et al., 2012; Trusel et al., 2015).

We investigated long-lasting changes in EPSPs induced upon a spike-timing-dependent plasticity paradigm (STDP) (Shen et al., 2008). As previously shown (Shen et al., 2008, 2015), a post-pre STDP protocol (Figure 1B), in which the postsynaptic spike preceded the presynaptic stimulation, did not induce plasticity in this SPN subpopulation (96% ± 4% of baseline, n = 8, p > 0.05; Figure 1B). Previous studies show that this STDP protocol leads to long-term synaptic depression (t-LTD) in dSPNs when either D1R is inhibited or striatal DA is depleted (Shen et al., 2008, 2015). The effects of decreased 5-HT signaling at glutamatergic dSPN synapses during STDP remain to be determined, however.

To address the role of 5-HT, we manipulated its release chemogenetically. We used a newly generated transgenic DREADD (designer receptors exclusively activated by designer drugs) mouse line (hM4Di^{+/-}/Pet1₂₁₀-Cre) obtained by crossing the Pet1₂₁₀-Cre transgenic mice (Pelosi et al., 2014) with hM4Di

knockin mice expressing the DREADD hM4Di receptor (hM4DiR) fused in-frame with the mCherry reporter and flanked by two sets of Lox sites (i.e., LoxP/Lox2722) (Figure S1A). Upon Cre-mediated somatic recombination, the double-floxed DREADD sequence is effectively inverted, leading to transcription of the inhibitory hM4DiR in Pet1₂₁₀-expressing 5-HT neurons. In brain slices from hM4Di^{+/-}/Pet1₂₁₀-Cre mice, these neurons and their projections can be identified by the expression of the mCherry reporter (Figures S1B and S2A–S2L). We first confirmed that the hM4DiR silenced serotonergic neurons of the DRN when activated by its cognate ligand clozapine-N-oxide (CNO; Figures S1C–S1F). Bath application of CNO (5 μM) significantly decreased the firing discharge of mCherry^{positive}, but not mCherry^{negative}, neurons (Figures S1D and S1E). CNO did not affect the firing rate of DRN 5-HT neurons from hM4Di^{+/-} littermates negative for Pet1₂₁₀-Cre (Pet1₂₁₀-Cre⁰; Figure S1F). mCherry- and 5-HT co-immunoreactivity was verified in the DLS of hM4Di^{+/-}/Pet1₂₁₀-Cre mice by immunoelectron microscopy (Figures S2M–S2O). We also confirmed the chemogenetic synaptic silencing of 5-HT release by *in vivo* 5-HT microdialysis in the DLS of freely moving hM4Di^{+/-}/Pet1₂₁₀-Cre mice and their control littermates (Figure S3A). In hM4Di^{+/-}/Pet1₂₁₀-Cre mice, high K⁺ (120 mM) infusion in the DLS induced reliable 5-HT release from serotonergic terminals; the stimulatory effect of a second high K⁺ challenge was blunted by the intrastriatal infusion of CNO (5 μM; p < 0.01; Figure S3B). CNO did not affect K⁺-evoked release of 5-HT when infused in control mice (p > 0.05; Figure S3C). These controls indicate that this model can be used to effectively silence the synaptic release of 5-HT in the DLS.

We assessed whether 5-HT synaptic silencing in the DLS affects plasticity outcomes by CNO application in hM4Di^{+/-}/Pet1₂₁₀-Cre mice (Figure 1). In dSPNs from these mice, t-LTD occurred in the presence of CNO, but not in the control (5 μM; 74% ± 4% of baseline, n = 8, p < 0.05 versus 95% ± 5% of baseline, n = 5, p > 0.05; Figure 1C). There were no changes in synaptic responses when CNO was applied during a burst pattern of back propagating action potentials (bAPs) (94% ± 7% of baseline, n = 5, p > 0.05) or in the absence of the STDP protocol (93% ± 3% of baseline, n = 8, p > 0.05; Figure S4A). Further, CNO had no effect when applied during the post-pre STDP paradigm in naive mice (92% ± 6% of baseline, n = 7, p > 0.05; Figure S4B). These results indicate that the chemogenetic inhibition of 5-HT release in the DLS gates t-LTD at glutamatergic afferents to dSPNs.

Antagonism of 5-HT_{4R} Signaling Enables t-LTD in dSPNs

Our findings indicate that 5-HT, similar to DA through the Gs-coupled D1R (Shen et al., 2008), disrupts the induction of t-LTD in dSPNs. To elucidate the molecular mechanism underlying this role, we tested whether inhibiting 5-HT Gs-mediated signaling allows t-LTD to occur. The striatum contains a high density of the 5-HT_{6R} and 5-HT_{4R} subtypes, which, like D1R, are coupled to Gs and are expressed in SPNs (Di Matteo et al., 2008). Bath perfusion of the 5-HT_{6R} antagonist SB271046 (5 μM) during the post-pre protocol did not promote plasticity in dSPNs (99% ± 8% of baseline, n = 6, p > 0.05; Figure S4C). In contrast, delivering the post-pre STDP paradigm in the




Figure 1. Chemogenetic Inhibition of 5-HT Release in the DLS Promotes the Induction of t-LTD at dSPN Glutamatergic Synapses

(A) (Top) Experimental configuration in horizontal brain slices containing the dorsolateral striatum (DLS). Schematic shows glutamatergic cortical and thalamic inputs to dSPNs, as well as dopaminergic and serotonergic afferents, which are recruited following intra-striatal electrical stimulation. (Bottom) Confocal laser scanning microscopy images of triple immunofluorescence for adenosine A2A receptor (A2AR), substance P (SP), and biotin in patch-recorded neurons.

(B) (Top) The post-pre pairing protocol for the induction of t-LTD. (Bottom) In dSPNs from naive mice, t-LTD was not induced by the post-pre timing pairing (dSPN, $n = 8$, 7 mice; RM1WA $F_{7,44} = 1$, $p = 0.4$).

(C) Experimental configuration in brain slices from hM4D^{+/+}/Pet1₂₁₀-Cre mice, highlighting the 5-HT terminals (green) originating from the dorsal raphe nucleus (DRN). In DLS dSPNs from these mice, bath perfusion of CNO (5 μ M; blue bar) during STDP gated a form of t-LTD, which was absent under control conditions (dSPN_{DREADD} + CNO, $n = 8$, 7 mice; RM1WA $F_{7,44} = 8$, $p < 0.0001$, $p < 0.05$ Tukey; dSPN_{DREADD}, $n = 5$, 5 mice; RM1WA $F_{4,44} = 1.5$, $p = 0.3$; dSPN_{DREADD} + CNO versus dSPN_{DREADD}, * $p < 0.05$, Mann-Whitney test).

(B and C) In these figures, and in the analogous plots that follow throughout the manuscript, data are presented as a time course (mean \pm SEM) of normalized EPSP amplitudes and normalized Rinp. Scatter plot summarizes the ratios of synaptic responses after (a) and before (b) the STDP, at the time points indicated. Insets represent superimposed averaged recordings (ten traces) before and after the delivery of the STDP protocol (vertical bar) (See also Figures S1–S4.)

onist in the absence of the pairing protocol did not affect glutamatergic synaptic transmission (GR113808, 94% \pm 5% of baseline, $n = 5$, $p > 0.05$; RS39604, 96% \pm 4% of baseline, $n = 6$, $p > 0.05$; Figure S4D), confirming that coordinated postsynaptic and presynaptic activity is necessary for the induction of t-LTD.

Previous studies demonstrated that in dSPNs, antagonism of the Gs-coupled D1R signaling allows endocannabinoid (eCB) biosynthesis, ultimately leading to LTD mediated by activation of the presynaptic cannabinoid receptor CB1 (Shen et al., 2008, 2015). However, this was not

presence of the 5-HT₄R antagonists GR113808 (5 μ M) or RS39604 (5 μ M) revealed t-LTD in the same neuronal subpopulation (GR113808, 79% \pm 5% of baseline, $n = 8$, $p < 0.05$; RS39604, 68% \pm 4% of baseline, $n = 10$, $p < 0.05$; Figure 2A). Thus, 5-HT₄R inhibition recapitulates the synaptic effects of chemogenetic inhibition of 5-HT release. Application of either antagon-

the putative mechanism for antagonism of 5-HT₄R signaling, as the co-application of RS39604 with the CB1R antagonist AM251 (4 μ M) did not prevent t-LTD induction (RS39604 + AM251, 62% \pm 7% of baseline, $n = 5$, $p < 0.05$; Figure 2B).

We next investigated the locus of plasticity expression by recording synaptic responses generated by twin stimuli, before




Figure 2. Antagonizing the 5-HT4R Subtype Is a Permissive Condition for t-LTD to Occur

(A) t-LTD was observed at glutamatergic dSPN synapses when the post-pre STDP protocol was delivered in the presence of the 5-HT4R antagonists GR113808 or RS39604 (5 μM; GR113808, n = 8, 7 mice; RM1WA $F_{7,44} = 7$, $p = 0.003$, $p < 0.05$ Tukey; RS39604, n = 10, 9 mice; RM1WA $F_{9,44} = 14$, $p < 0.0001$, $p < 0.05$ Tukey; 1WA $F_{2,23} = 10$, $p = 0.0008$, dSPN+ GR113808 versus dSPN, * $p < 0.05$, dSPN+ RS39604 versus dSPN, *** $p < 0.001$; Tukey). The two 5-HT4R antagonists were applied throughout the entire duration of the recording. Solid black line (average) is the time course from Figure 1B, reported here for reference.

(B) t-LTD gated by RS39604 was not blocked by the CB1R antagonist AM251 (4 μM; dSPN + RS39604/AM251, n = 5, 4 mice; RM1WA $F_{4,44} = 10$, $p = 0.001$, $p < 0.05$ Tukey; dSPN + AM251/RS39604 versus dSPN + RS39604, $p = 0.6$, Mann-Whitney test). Solid green line (average) is t-LTD from (A) obtained in the presence of RS39604, reported here for reference.

(C) t-LTD induced with the post-pre pairing upon application of RS39604 (5 μM) was not associated with changes in paired-pulse ratio (PPR). (Left) Plot shows the normalized first peak (p1) and second peak (p2) amplitudes of synaptic responses generated by twin stimuli at a 50 ms interval as a function of time (dSPN + RS39604, n = 6, 5 mice; p1, RM1WA $F_{5,44} = 5$, $p < 0.0001$, $p < 0.05$ Tukey; p2, RM1WA $F_{5,44} = 6$, $p < 0.0001$, $p < 0.05$ Tukey). (Right) Boxplot diagram indicates the PPRs of baseline (black box), and after STDP (red box) at the time points indicated ($p = 0.5$, paired t test). The values are the minimum, the median (bar inside the box), and the maximum. PPR is expressed as the ratio between the amplitude of the second and first EPSPs.

(D) In the presence of RS39604, t-LTD was blocked by intracellular application of the endocytosis-disrupting peptides D15 (1.5 mM; dSPN + D15, n = 8, 8 mice; RM1WA $F_{7,44} = 1.7$, $p = 0.17$) or GluA2_{3Y} (100 μg/ml; dSPN + GluA2_{3Y}, n = 6, 5 mice; RM1WA $F_{5,44} = 1$, $p = 0.4$). The specificity of action of D15 and GluA2_{3Y} was confirmed by their respective control peptides (S15 for D15; GluA2_{3A} for GluA2_{3Y}) (S15, 1.5 mM; dSPN + S15, n = 5, 3 mice; RM1WA $F_{4,44} = 4$, $p = 0.03$, $p < 0.05$ Tukey; GluA2_{3A}, 100 μg/ml; dSPN + GluA2_{3A}, n = 6, 5 mice; RM1WA $F_{5,44} = 8$, $p = 0.0007$, $p < 0.05$ Tukey; 1WA, $F_{4,28} = 7$, $p = 0.0004$; dSPN + S15 versus dSPN,

(legend continued on next page)

and after the STDP protocol. In this set of experiments, we found that t-LTD was enabled by antagonism of 5-HT₄R (RS39604, 77% ± 5% of baseline, n = 6, p < 0.05; Figure 2C) but was not accompanied by a change in paired-pulse ratio (PPR), a hallmark of presynaptic plasticity (PPR, before_{STDP}, 1.33 ± 0.09; after_{STDP}, 1.36 ± 0.10, p > 0.05; Figure 2C). This raises the possibility that this form of LTD is expressed postsynaptically. Consistent with this, intracellular dialysis of the peptide D15 (1.5 mM)—a peptide that prevents AMPAR endocytosis by disrupting the interaction between dynamin and amphiphysin (Carroll et al., 1999; Morishita et al., 2005)—blocked t-LTD (D15/RS39604, 91% ± 5% of baseline, n = 8, p > 0.05), whereas dialysis of a scramble peptide (S15, 1.5 mM) had no effect (S15/RS39604, 63% ± 8% of baseline, n = 5, p < 0.05; Figure 2D). The application in the postsynaptic neuron of the GluA2_{3Y} peptide (100 μg/ml), which blocks regulated, but not constitutive, AMPAR endocytosis (Brebner et al., 2005; Collingridge et al., 2010), had a similar effect (GluA2_{3Y}/RS39604, 94% ± 3% of baseline, n = 6, p > 0.05; Figure 2D). Confirming this peptide's specific action, the control peptide GluA2_{3A} (100 μg/ml) did not influence plasticity (GluA2_{3A}/RS39604, 70% ± 5% of baseline, n = 6, p < 0.05; Figure 2D). Furthermore, variance analysis of synaptic responses indicated that t-LTD resulting from chemogenetic inhibition of 5-HT release (Figure 1) or application of 5-HT₄R antagonists (Figure 2) is not accompanied by changes in trial-to-trial variation in EPSP amplitude (Figure S4E). This further supports a postsynaptic locus of plasticity expression (Shen et al., 2008).

Overall, these data indicate that 5-HT controls plasticity induction in dSPNs through the activation of 5-HT₄R; this effect is not mediated by eCBs, and it relies on postsynaptic expression mechanisms.

Serotonin Modulates bAP-Induced Ca²⁺ Transients during STDP

At glutamatergic synapses on SPNs, the induction of t-LTD depends on the temporal convergence of synaptic input and postsynaptic elevations of dendritic Ca²⁺ signal triggered by (bAPs) (Plotkin et al., 2013; Shindou et al., 2011). 5-HT has been shown to regulate ionic conductance that may affect the propagation of bAPs (King et al., 2008). Modulation of the bAP-induced Ca²⁺ signal may therefore be one of the mechanisms that affect timing-dependent synaptic plasticity in dSPNs (Plotkin et al., 2013; Shindou et al., 2011; Zhou et al., 2005). To test this hypothesis, simultaneously with the chemogenetic and pharmacological inhibition of 5-HT signaling, we performed two-photon Ca²⁺ imaging and whole-cell current clamp recordings and monitored changes in intracellular Ca²⁺. In the proximal dendrites of dSPNs, a single bout of the negative STDP protocol (1b-STDP) triggered Ca²⁺ transients that were stable over time, as indicated by the lack of variation in the area under the curve (AUC) of the fluorescence transients over 10–15 min (106% ±

8% of control; n = 8, p > 0.05; Figure 3A). In the dSPNs of hM4Di^{+/−}/Pet1₂₁₀-Cre mice, bath application of CNO (5 μM) to impede 5-HT release enhanced dendritic Ca²⁺ influx in response to 1b-STDP (AUC, 130% ± 8% of control; n = 11, p < 0.01; Figure 3B). The 1b-STDP-associated dendritic Ca²⁺ signal increased upon application of the two 5-HT₄R antagonists GR113808 (5 μM; AUC, 122% ± 6% of control; n = 15, p < 0.01; Figure 3C) or RS39604 (5 μM; AUC, 130% ± 7% of control; n = 14, p < 0.01; Figure 3C). Together, these results indicate that impeding 5-HT₄R-mediated signaling boosts the bAP-induced dendritic Ca²⁺ signal.

These results appear to conflict with the notion that activation, not inhibition, of Gs-coupled receptors typically enhances voltage-gated Ca²⁺ currents triggered by APs via stimulation of protein kinase A (PKA) (Higley and Sabatini, 2010). In several neuronal subtypes, including striatal SPNs, inhibition of various types of K⁺ currents that contribute to the AP repolarization, and after hyperpolarization phases, enhances dendritic Ca²⁺ transients (Cai et al., 2004; Day et al., 2008; Trusel et al., 2015). Therefore, we hypothesized that 5-HT₄R regulates dendritic Ca²⁺ levels by instead influencing the activity of K⁺ channels. In addition to PKA, 5-HT₄R can potentially signal to Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) (Liu and Voyno-Yasenetskaya, 2005; Ponimaskin et al., 2002), which positively regulates the activity of the large-conductance Ca²⁺-activated K⁺ channel BK (Nelson et al., 2005; van Welie and du Lac, 2011). To test if this is the case in dSPNs, we applied the BK channel blocker paxilline (10 μM) to isolate the BK-mediated current (as the paxilline-sensitive component, I_{paxilline}) in response to a depolarizing voltage-ramp protocol (Figure S5A) (Deng et al., 2013). The 5-HT₄R agonist RS67333 (5 μM) significantly enhanced I_{paxilline} amplitude (p < 0.05); this effect was prevented by the postsynaptic intracellular application of a specific competitive inhibitor of CaMKII, the autocamide-2-related inhibitory peptide (AIP, 10 μM; p > 0.05; Figure S5B). We also confirmed that stimulating 5-HT₄R activates CaMKII in DLS dSPNs. On acute striatal slices from drd1a-tdTomato^{tg} mice, which express Tomato in dSPNs (Shuen et al., 2008), we monitored phosphorylation of CaMKII at Thr286 (pCaMKII). Incubation (10 min) with the 5-HT₄R agonist RS67333 (5 μM) significantly increased the number of pCaMKII⁺/Tomato⁺ cells compared to controls (p < 0.01; Figure S5C). These data indicate that 5-HT₄R can signal through CaMKII to positively modulate BK channel activity in dSPNs.

BK channels shape the duration of bAP in several neuronal populations and affect dendritic Ca²⁺ influx (Golding et al., 1999). Therefore, reduced release of 5-HT or direct inhibition of 5-HT₄R could dampen the activation of BK channels during the STDP protocol, leading to an enhanced dendritic Ca²⁺ signal in response to bAPs. To test this, we first assessed whether blocking BK channels occluded the effect of RS39604. Bath perfusion of paxilline (10 μM) significantly enhanced the

**p < 0.01; dSPN + D15 versus dSPN, p = 0.9, dSPN + D15 versus dSPN + S15, *p < 0.05; dSPN + GluA2_{3A} versus dSPN, *p < 0.05; dSPN + GluA2_{3Y} versus dSPN, p = 1; dSPN + GluA2_{3Y} versus dSPN + GluA2_{3A}, *p < 0.05; Tukey). Solid black line (average) is the time-course from Figure 1B, reported here for reference. (A–D) (Insets, left) In these figures, and throughout the manuscript, schematics depict proposed signaling pathway targeted by the defined drug, unless specified otherwise (see also Figure S4).




Figure 3. 5-HT Signaling Modulates bAP-Induced Dendritic Ca²⁺ Transients in dSPNs

(A) (Top left) Maximum-intensity projection images of a dSPN filled with Alexa 568. Blue dotted circle lines delimit the region of proximal dendrites, 30–50 μ m from the soma. (Bottom left) High magnification of a dSPN-proximal dendrite from which line scans were taken (yellow dotted line). (Center) Dendritic Ca²⁺ transients and the corresponding relative changes in fluorescence that are induced by one bout of the post-pre STDP paradigm (1b-STDP, inset) for control conditions, and after 10–15 min ($n = 8$, 4 mice; paired t test, $p = 0.4$). (B) Chemogenetic inhibition of 5-HT release in the DLS of hM4Di^{+/+}/Pet1₂₁₀-Cre mice enhanced dendritic Ca²⁺ transients in response to 1b-STDP ($n = 11$, 6 mice). (B–G) (Left) Comparison of Ca²⁺ transients in response to 1b-STDP (B–D) or to a burst pattern of bAPs (E–G), before and after a specific pharmacological manipulation, as indicated.

(legend continued on next page)

1b-STDP-associated Ca^{2+} transient (AUC, $138\% \pm 7\%$ of control; $n = 9$, $p < 0.001$; [Figure 3D](#)). As predicted, sequential co-application of paxilline with RS39604 had no further effect on the induced change in dendritic fluorescence (AUC, $139\% \pm 7\%$ of control; $p < 0.001$; $100\% \pm 3\%$ of paxilline, $p > 0.05$; $n = 9$; [Figure 3D](#)), suggesting that BK channels are targets of 5-HT4R signaling.

5-HT4R Modulates bAP-Induced Ca^{2+} Signal through CaMKII

Our 1b-STDP-stimulation protocol is likely to activate a set of glutamatergic, dopaminergic, and serotonergic inputs. Thus, it cannot isolate the impact of 5-HT4R on dendritic Ca^{2+} influx specifically induced by bAPs. To overcome this issue, we estimated the Ca^{2+} transients generated by somatically eliciting postsynaptic bAPs in the absence of presynaptic stimulation ([Figure 3E](#)). We replaced the contribution of 5-HT release on local electrical stimulation with the direct pharmacological stimulation of 5-HT4R by the agonist RS67333 ($5 \mu\text{M}$). Consistent with the hypothesis that 5-HT4R stimulates a K^+ channel conductance, the presence of RS67333 reduced the bAP-induced Ca^{2+} signal (AUC, $71\% \pm 7\%$ of control; $n = 10$, $p < 0.01$; [Figure 3E](#)). Furthermore, blocking BK-mediated current prevented the effect of 5-HT4R stimulation: RS67333 failed to reduce the bAP-induced Ca^{2+} transients upon pre-exposure to paxilline (AUC, paxilline, $142\% \pm 10\%$ of control; $p < 0.01$; paxilline + RS67333, $142\% \pm 9\%$ of control, $p < 0.05$; paxilline + RS67333, $101\% \pm 7\%$ of paxilline, $p > 0.05$; $n = 7$; [Figure 3F](#)). Further, the intracellular application of the CaMKII inhibitor AIP ($10 \mu\text{M}$) prevented RS67333 from reducing Ca^{2+} transients elicited by bAPs (AUC, $105\% \pm 5\%$ of AIP, $p > 0.05$; $n = 10$; [Figure 3G](#)). The selective cell-permeable CaMKII inhibitor KN-62 ($10 \mu\text{M}$) had a similar effect: bath application boosted bAP-induced Ca^{2+} transients ([Figure S5D](#)). This is consistent with bursts of bAPs activating CaMKII in proximal dendrites ([Lee et al., 2009](#)) and with CaMKII hindering the activation of BK channels recruited by bursts of bAPs, thus resulting in enhanced Ca^{2+} transients. KN-62 prevented the inhibitory effect of RS67333 (AUC, KN-62, $155\% \pm 15\%$ of control, $p < 0.05$; KN-62 + RS67333, $158\% \pm 22\%$ of control, $p < 0.05$; KN-62 + RS67333, $99\% \pm 7\%$ of KN-62, $p > 0.05$; $n = 7$; [Figure S5D](#)), further supporting that CaMKII acts downstream of 5-HT4R to regulate bAP-induced Ca^{2+} transients.

These results indicate that the modulatory effect of 5-HT4R activation involves CaMKII signaling. PKA may also act in the same pathway to shape bAP-induced Ca^{2+} signal. Nevertheless, when the PKA inhibitor peptide 5-24 ($20 \mu\text{M}$) was intracellularly applied, RS67333 still reduced bAP-induced Ca^{2+} signal (AUC, $83\% \pm 5\%$ of PKI; $p < 0.05$, $n = 7$; [Figure S5E](#)), suggesting PKA does not play a major role in this process.

Overall, these data indicate that regulation of BK channel activity is required for the 5-HT4R-mediated modulation of dendritic Ca^{2+} signal in dSPNs and that this modulation involves CaMKII signaling.

5-HT Signaling Controls t-LTD at Thalamic Inputs

We have demonstrated that 5-HT signaling, through the 5-HT4R subtype, influences the postsynaptic response of dSPNs to glutamatergic input activation during STDP. To dissect whether this modulation occurs at cortical and/or thalamic inputs, we probed the striatal circuitry with optogenetics. First, we used Thy1-ChR2 transgenic mice to measure postsynaptic responses evoked by light activation of corticostriatal afferents, which represent at least 50% of glutamatergic synapses on SPNs ([Doig et al., 2010](#); [Huerta-Ocampo et al., 2014](#)). In these mice, the expression of channel rhodopsin (ChR2) is high in cortical layer V neurons ([Figure 4A](#)) and appears not to extend to thalamostriatal neurons ([Plotkin et al., 2014](#); [Wu et al., 2015](#)). Short blue light pulses through the microscope objective ($0.1\text{--}1 \text{ ms}$ at 470 nm) elicits EPSPs in dSPNs (o-EPSP_{Ctx}). We investigated post-pre STDP at corticostriatal synapses by monitoring o-EPSP_{Ctx} before and after the induction protocol. During the post-pre pairing paradigm, optogenetic stimulation of cortical inputs was replaced by conventional electrical stimulation in order to recruit 5-HT fibers ([Figure 4B](#)). In this experimental setting, the STDP protocol failed to induce significant changes in o-EPSP_{Ctx} ($98\% \pm 8\%$ of baseline, $n = 7$, $p > 0.05$; [Figure 4C](#)), even when applied in the presence of RS39604 ($5 \mu\text{M}$; $92\% \pm 6\%$ of baseline, $n = 8$, $p > 0.05$; [Figure 4C](#)). To corroborate this lack of involvement of corticostriatal synapses in a second model, we injected an adeno-associated viral (AAV) vector encoding ChR2-eYFP under the control of the CaMKIIa promoter [rAAV5-CaMKIIa-ChR2(H134R)-eYFP] into the M1 motor cortex region of C57BL/6J mice ([Johnson et al., 2017](#)) ([Figure S6A](#)). As with Thy1-ChR2 mice ([Figure 4C](#)), the STDP protocols failed to induce significant changes in dSPNs o-EPSP_{Ctx}, regardless of the presence or absence of RS39604 (dSPN_{Ctx}, $97\% \pm 6\%$ of baseline, $n = 7$, $p > 0.05$; dSPN_{Ctx} + RS39604, $90\% \pm 2\%$ of baseline, $n = 6$, $p > 0.05$; dSPN_{Ctx} versus dSPN_{Ctx} + RS39604, $p = 0.7$ Mann Whitney test; [Figures S6B](#) and [S6C](#)). Thus, corticostriatal synapses are not the primary target of the 5-HT4R-mediated effects at non-dissected glutamatergic inputs to dSPNs ([Figure 2A](#)).

These results also suggest that gating of t-LTD due to decreased 5-HT signaling is in fact occurring at thalamostriatal inputs. To test this, we virally expressed ChR2 in the thalamus of C57BL/6J mice ([Figure 4D](#)) ([Ellender et al., 2013](#); [Johnson et al., 2017](#)) and measured light-evoked EPSPs (o-EPSP_{Thal}) at thalamostriatal connections. To induce plasticity, we applied the same post-pre STDP paradigm described for corticostriatal synapses ([Figure 4B](#)). As predicted, the STDP protocol did not evoke plasticity of o-EPSP_{Thal} in dSPNs under control conditions ($103\% \pm 10\%$ of baseline, $n = 6$, $p > 0.05$; [Figure 4E](#)), whereas it triggered a significant t-LTD of o-EPSP_{Thal} in the presence of RS39604 ($5 \mu\text{M}$; $71\% \pm 8\%$ of baseline, $n = 8$, $p < 0.05$, [Figure 4E](#)). We also controlled for the cell-type specificity of this effect. In iSPNs, the post-pre STDP paradigm already induced t-LTD of

(A–G) (Right) Plot summary of modulation of dendritic Ca^{2+} measured as area under the curve (AUC) of the fluorescence transients, in control conditions and with pharmacological manipulations, as indicated (A, B, and C, GR113808, $n = 15$, 10 mice; RS39604, $n = 14$, 11 mice; E, $n = 10$, 4 mice; G, $n = 10$, 4 mice; ** $p < 0.01$, paired t test; D, $n = 9$, 5 mice; RM1WA, $F_{8,2} = 20$ $p < 0.0001$, paxilline versus control, *** $p < 0.001$; paxilline + RS39604 versus control, *** $p < 0.001$; paxilline versus paxilline + RS39604, $p = 1$; Tukey; F, $n = 7$, 6 mice; RM1WA, $F_{6,2} = 9$ $p = 0.005$, paxilline versus control, ** $p < 0.01$; paxilline + RS67333 versus control, * $p = 0.01$, Tukey; paxilline versus paxilline + RS67333, $p = 1$; Tukey). (See also [Figure S5](#).)




Figure 4. Decreased 5-HT Signaling Gates t-LTD at Thalamo-dSPN Synapses

(A) Confocal images of horizontal sections of Thy1-ChR2 transgenic mice; scale bar, 500 μ m; Ctx, cortex; Str, striatum; Hip, hippocampus; Thal, thalamus. Inset shows ChR2-eYFP expressing axon terminals in the DLS; scale bar, 10 μ m.

(B) Schematic of recording configuration in dSPNs. (Top) Corticostriatal stimulation evoked by 470 nm LED pulses through the microscope objective was performed before and after the STDP protocol. (Bottom) During the post-pre STDP paradigm, light stimulation of cortical terminals was replaced by local electrical stimulation.

(C) The post-pre STDP protocol failed to induce plasticity of o-EPSP_{Ctx} either under control conditions (dSPN_{Ctx}, n = 7, 6 mice; RM1WA $F_{7,44} = 1.1$, p = 0.4) or upon RS39604 application (dSPN_{Ctx} + RS39604, n = 8, 8 mice; RM1WA $F_{7,44} = 0.8$, p = 0.5).

(D) (Left and top right) rAAV5-CaMKIIa-ChR2(H134R)-eYFP expression targeting the thalamus (Thal); scale bar, 500 μ m; Ctx, cortex; Str, striatum; Hip, hippocampus. (Inset) ChR2-eYFP⁺ axons in the DLS; scale bar, 10 μ m. (Right) Schematic of recording configuration of o-EPSP_{Thal} evoked by 470 nm LED pulses (center right), and during the delivery of the STDP paradigm (bottom right).

(E) t-LTD is induced at thalamo-dSPN synapses upon 5-HT4R inhibition (dSPN_{Thal}, n = 6, 6 mice; RM1WA $F_{5,44} = 1.5$, p = 0.3; dSPN_{Thal} + RS39604, n = 8, 4 mice; RM1WA $F_{7,44} = 10$, p = 0.0002, p < 0.05 Tukey; dSPN_{Thal} + RS39604 versus dSPN_{Thal}, **p < 0.01, Mann-Whitney test).

(F) In hM4Di^{-/-}/Pet1₂₁₀-Cre mice injected with the rAAV5-CaMKIIa-ChR2(H134R)-eYFP virus in the thalamus, bath application of CNO (5 μ M) gated t-LTD of o-EPSP_{Thal} in response to STDP (dSPN_{DREADD_Thal}, n = 7, 4 mice; RM1WA $F_{6,44} = 1.8$, p = 0.1; dSPN_{DREADD_Thal} + CNO, n = 7, 4 mice; RM1WA $F_{6,44} = 10$, p = 0.0002, p < 0.05 Tukey; dSPN_{DREADD_Thal} + CNO versus dSPN_{DREADD_Thal}, ***p < 0.001, Mann-Whitney test). (See also Figures S6–S8.)

o-EPSP_{Thal} in control conditions (o-EPSP_{Thal}, 67% \pm 5% of baseline, n = 7, p < 0.05; Figures S7A and S7B); the application of RS39604 did not influence the plasticity outcome (o-EPSP_{Thal} + RS39604, 68% \pm 8% of baseline, n = 5, p < 0.05; Figure S7B).

We next assessed the link between the synaptic effects of 5-HT at thalamic inputs with the regulation of BK channel activity (Figure 3). In our model, decreased 5-HT signaling dampens the

activation of BK channels during STDP and boosts the Ca²⁺ signal induced by bAP (Figure 3). This mechanism could underlie the permissive role of 5-HT4R inhibition on t-LTD induction (Figures 2A and 4E). If this is the case, we would expect the pharmacological inhibition of BK channels during conventional STDP to be sufficient to gate synaptic depression at thalamic inputs recruited by optogenetic activation. Reminiscent of 5-HT4R

antagonism and chemogenetic inhibition of 5-HT release, in the presence of the BK channel blocker paxilline (10 μ M), the STDP paradigm resulted in t-LTD of o-EPSP_{Thal} (75% \pm 4% of baseline, $n = 7$, $p < 0.05$; [Figure S8A](#)).

The hm4Di^{+/-}/Pet1₂₁₀-Cre animals expressing ChR2 in the thalamus showed effects ([Figure 4F](#)) similar to those described above. The chemogenetic inhibition of 5-HT release by bath application of CNO (5 μ M) during the post-pre STDP paradigm enabled t-LTD at thalamo-dSPN synapses (dSPN_{DREADD_Thal}, 93% \pm 1% of baseline, $n = 7$, $p > 0.05$; dSPN_{DREADD_Thal}, + CNO, 61% \pm 5% of baseline, $n = 7$, $p < 0.05$; [Figure 4F](#)). Together, these results indicate that impeding serotonergic signaling, and in turn the activation of the 5-HT4R subtype, is sufficient to gate t-LTD at thalamostriatal synapses.

In addition to bAP-induced Ca²⁺ transients, NMDA receptor activation is a possible means by which dendritic Ca²⁺ signals could be enhanced to enable t-LTD at thalamostriatal synapses on dSPNs ([Ellender et al., 2013](#); [Wu et al., 2015](#)). However, the NMDAR antagonist R-CPP (10 μ M) had no effect on o-EPSP_{Thal} t-LTD gated by inhibition of BK channels during the post-pre STDP paradigm (dSPN_{Thal} + Paxilline + R-CPP, 78% \pm 6% of baseline, $n = 7$, $p < 0.05$; [Figure S8A](#)) or by chemogenetic inhibition of 5-HT release in hm4Di^{+/-}/Pet1₂₁₀-Cre mice (dSPN_{DREADD_Thal}, + CNO + R-CPP, 60% \pm 7% of baseline, $n = 5$, $p < 0.05$; [Figure S8B](#)). This indicates that NMDARs are not involved in this form of synaptic depression.

Lastly, we asked whether activating 5-HT4R signaling during the post-pre STDP paradigm is sufficient to disrupt t-LTD at thalamic inputs to dSPNs. We first tested the assumption that t-LTD at thalamo-dSPN synapses is suppressed by neuromodulatory signals (including 5-HT) released upon conventional electrical stimulation during STDP, as previously proposed ([Wu et al., 2015](#)). We re-examined plasticity of o-EPSP_{Thal} upon the delivery of an optogenetic post-pre pairing STDP protocol (o-STDP), which exclusively engages thalamic afferents. In this experimental setting, we observed a form of t-LTD at thalamo-dSPN synapses (72% \pm 6% of baseline, $n = 6$, $p < 0.05$, [Figure 5A](#)), consistent with previous observations ([Ellender et al., 2013](#); [Wu et al., 2015](#)). The 5-HT4R agonist RS67333 (5 μ M) prevented t-LTD of o-EPSP_{Thal} when applied during the o-STDP paradigm (91% \pm 3% of baseline, $n = 8$, $p < 0.05$; [Figure 5A](#)). We also injected the rAAV5-EF1a-DIO-hChR2(H134R)-eYFP virus into the DRN of Pet1₂₁₀-Cre mice to exert an optogenetic control of 5-HT release during plasticity induction ([Figure 5B](#)). In oblique horizontal slices from these mice, we electrically elicited thalamic synaptic responses in the DLS (e-EPSP_{Thal}) by placing the stimulating electrode in the thalamus, close to the border of the thalamic reticular nucleus ([Figure 5C](#)) ([Ding et al., 2008, 2010](#)). Consistent with the response to the o-STDP paradigm ([Figure 5A](#)), e-EPSP_{Thal} showed t-LTD upon a post-pre STDP paradigm in which thalamic axons were stimulated (Thal-STDP) (61% \pm 7% of baseline, $n = 6$, $p < 0.05$; [Figures 5C and 5D](#)). Optogenetically evoking 5-HT release at the time of presynaptic stimulation during pairing prevented t-LTD of e-EPSP_{Thal} (93% \pm 5% of baseline, $n = 8$, $p > 0.05$); plasticity was rescued by the application of RS39604 (5 μ M; 44% \pm 7% of baseline, $n = 5$, $p < 0.05$; [Figures 5C and 5D](#)). This also indicates that

t-LTD suppression is not due to light-induced release of glutamate from DRN terminals.

Collectively, these results further support the involvement of the 5-HT4R subtype in the 5-HT-mediated modulation of t-LTD at thalamic inputs on dSPNs.

5-HT4R Preferentially Localizes at Thalamostriatal Synapses

To increase certainty of the input specificity of the 5-HT4R mediated effects, we combined immunohistochemistry with three-dimensional (3D) segmentation and morphometric analysis of glutamatergic DLS synapses ([Gilles et al., 2017](#)) ([Figures 6, S9A, and S9B](#)). We initially monitored the expression of 5-HT4R in two different mouse lines: the VGLUT1-Venus knockin line, which expresses the VGLUT1 transporter fused to the YFP-variant Venus and which allows direct visualization of axonal boutons from cortical afferents ([Heck et al., 2015](#); [Herzog et al., 2011](#); [Smith et al., 2014](#)), and the VGLUT2-Cre transgenic mice injected in the thalamus with the AAVrh9-pCAG-flex-eGFP-WRPE virus (AAV-FLEX-eGFP), to visualize thalamic inputs to the striatum. In this second mouse line, thalamic boutons were identified by immunolabeling for the presynaptic marker Bassoon ([Südhof, 2012](#)). We measured the 3D edge-to-edge minimal distance from VGLUT1-positive cortical boutons and VGLUT2-positive thalamic boutons to the closest 5-HT4R. Although there was a higher density of VGLUT1 boutons, the distance between VGLUT1 boutons and 5-HT4R was significantly greater than between VGLUT2 boutons and 5-HT4R (VGLUT1-5-HT4R, 608 \pm 26 nm; VGLUT2-5-HT4R, 364 \pm 19 nm, $p < 0.0001$; with 19.34% \pm 1.23% of VGLUT2 co-localizing with 5-HT4R and 9.81% \pm 0.74% of VGLUT1 co-localizing with 5-HT4R; [Figure 6A](#)). These morphological data support our functional observations showing that thalamostriatal synapses are the primary site of 5-HT4R-mediated effects ([Figure 4](#)).

We then investigated the postsynaptic subcellular localization of 5-HT4R in dSPNs. The viral construct AAVrh9-PPTA-Cre (AAV-PPTA-Cre), which contains an expression cassette consisting of Cre recombinase driven by the dSPN-specific promoter of the PPTA gene (substance P), and the AAVrh9-pCAG-flex-tdTomato-WRPE (AAV-FLEX-tdTOMATO) construct expressing “flexed” tdTomato ([Dos Santos et al., 2017](#); [Hikida et al., 2010](#); [Yagishita et al., 2014](#)) were co-injected into the DLS. In dSPNs, we measured the number of co-localizing 5-HT4Rs in 3D-reconstructed dendrites and segmented headspines. 5-HT4R localization is predominant in the shaft, with 3.2 \pm 0.5 5-HT4R per 10 μ m of dendritic length ([Figure 6B](#)). To selectively visualize thalamic terminals on dSPNs, we analyzed VGLUT2-Cre mice co-injected with the AAV-FLEX-eGFP virus in the thalamus and with the AAV-PPTA-Cre and the AAV-FLEX-tdTOMATO viruses in the DLS. This analysis confirmed the postsynaptic subcellular localization of 5-HT4R in both axo-spinous and axo-dendritic thalamostriatal synapses ([Figure 6C](#)).

DISCUSSION

The results presented here reveal a previously unknown role of 5-HT at striatal circuits. This is the first demonstration that signaling by 5-HT receptors provides a control mechanism of




Figure 5. Activation of 5-HT Signaling Blunts t-LTD Induction at Thalamo-dSPN Synapses

(A) (Left) Schematic of recording configuration of o-EPSP_{Thal} evoked by 470 nm LED pulses, and during the delivery of the optogenetic STDP paradigm. (Right) The o-STDP protocol induced t-LTD at thalamostriatal synapses (n = 6, 2 mice; RM1WA $F_{5,44} = 4$, $p < 0.05$ Tukey); plasticity was blocked by application of the 5-HT4R agonist RS67333 (n = 8, 3 mice; RM1WA $F_{7,44} = 2$, $p = 0.13$; dSPN_{Thal} + RS67333 versus dSPN_{Thal}, * $p < 0.05$, Mann-Whitney test). (B) (Left) Schematic of bilateral virus injection [rAAV5-EF1a-DIO-hChR2(H134R)-eYFP] in the DRN of Pet1₂₁₀-Cre mice. (Right) (1–3) Confocal images show the expression of 5-HT (red), YFP (green) and merge in the DRN. DAPI nuclear staining (inset in 2) identifies the two injection sites. Scale bar, 200 μm. (4–9) High-resolution confocal images showing SERT (red) and YFP (green) expression, and their co-localization (merge) in serotonergic afferents to the DLS. Scale bars 20 μm (4–6) and 10 μm (7–9).


(C) (Top) Experimental configuration in para-horizontal slices, showing the DLS, lateral globus pallidus (LGP), thalamic reticular nucleus (Rt), and the DRN. (Bottom) The electric post-pre pairing protocol (Thal-STDP) for the induction of t-LTD and timing of 5-HT_{opto} application.

(D) (Left) Recording configuration. (Right) In dSPNs, t-LTD of e-EPSP_{Thal} (dSPN_{e-Thal}, n = 6, 6 mice; RM1WA $F_{5,44} = 4$, $p < 0.0001$, $p < 0.05$ Tukey) was suppressed by light-induced 5-HT release (dSPN_{e-Thal} + 5-HT_{opto}, n = 8, 6 mice; RM1WA $F_{7,44} = 0.8$, $p = 0.9$), and rescued by application of the 5-HT4R antagonist RS39604 (5 μM, dSPN_{e-Thal} + 5-HT_{opto} + RS39604, n = 5, 4 mice; RM1WA $F_{4,44} = 18$, $p = 0.0003$, $p < 0.05$ Tukey; 1WA, $F_{2,16} = 16$, $p = 0.0002$; dSPN_{e-Thal} versus dSPN_{e-Thal} + 5-HT_{opto}, ** $p < 0.01$; dSPN_{e-Thal} versus dSPN_{e-Thal} + 5-HT_{opto} + RS39604, $p = 0.2$; dSPN_{e-Thal} + 5-HT_{opto} versus dSPN_{e-Thal} + 5-HT_{opto} + RS39604, *** $p < 0.001$; Tukey).

Hebbian plasticity (t-LTD) at thalamo-SPN synapses of the direct pathway. These synaptic effects of 5-HT are mediated by the postsynaptic 5-HT4R subtype, which shapes bAP-evoked dendritic Ca²⁺ signals through the regulation of BK channel

activity and which is preferentially localized at axo-dendritic thalamostriatal synapses.

Our findings help elucidate the full complement of 5-HT actions in the DS. Previous observations have shown that



spine head (Bottom) from dSPNs (red). Only 5-HT4Rs co-localizing with thalamic axonal bouton and dendrite shaft or spine are shown. Thalamic axons were labeled by injecting AAV-FLEX-eGFP in the thalamus of VGLUT2-Cre mice. dSPNs were labeled by injecting AAV-PPTA-Cre and AAV-FLEX-tdTOMATO in the DLS. 5-HT4R was detected by immunolabeling. (A–C) Scale bars, 1 μ m. (See also Figure S9.)

increasing 5-HT levels in the DLS reduces presynaptic glutamate release and depresses neurotransmission at corticostriatal inputs through the activation of the presynaptic 5-HT_{1b} receptor (Mathur et al., 2011). Notably, 5-HT_{1b}-mRNA is detectable in layer V cortical neurons, but not in the thalamic intralaminar nuclei (Allen Brain Atlas; <http://www.brain-map.org/>). In the current study, we find that impeding serotonergic signaling, thereby dampening the activation of the 5-HT_{4R} subtype, affects postsynaptic dendritic Ca²⁺ signals and gates spike-timing-dependent long-term depression (t-LTD) preferentially at thalamic afferents to dSPNs.

Results presented in this study and by Mathur et al. (2011) highlight the complexity of striatal 5-HT signaling, which acts through diverse receptors at both presynaptic and postsynaptic sites (Mathur and Lovinger, 2012) and which may show specific subcellular compartmentalization. Here, we provide

evidence that 5-HT_{4R}s are enriched at thalamostriatal synapses compared to cortical ones, and primarily localize at the dendritic shaft. Exogenously applied 5-HT or inhibition of 5-HT reuptake depresses corticostriatal afferents, whereas decreased synaptic release of 5-HT during STDP gates t-LTD at thalamostriatal inputs, suggesting that the net synaptic effect of 5-HT likely depends on its spatiotemporal dynamic in response to different patterns of stimulation. This may ultimately lead to spatiotemporal specificity of synaptic plasticity at defined inputs (i.e., cortical and thalamic inputs).

Upon impeding 5-HT signaling, t-LTD at undissected glutamatergic inputs is of an only slightly lower magnitude than that at thalamostriatal inputs recruited by optogenetic activation. These results appear to be at odds with the notion that at least 50% of excitatory inputs to SPNs are from the cortex (Doig et al., 2010; Huerta-Ocampo et al., 2014) and our current finding that

Figure 6. Postsynaptic Subcellular Localization of 5-HT_{4R}

(A) (Left) 3D volume rendering of confocal image stack showing 5-HT_{4R} and glutamatergic boutons in the DLS. Cortical VGLUT1 positive boutons were imaged in VGLUT1-Venus knockin mice. Thalamic VGLUT2 boutons were imaged in VGLUT2-Cre mice injected in the thalamus with the AAV-FLEX-eGFP virus. Co-localization with immunostained presynaptic active zone protein Bassoon allowed the selection of thalamic boutons from the segmented images (see Figure S9). (Right) Measurements of 3D border-to-border minimal distance from VGLUT1-positive boutons and VGLUT2-positive boutons to the closest 5-HT_{4R}. 18 to 24 image stacks from 2 sections from 3 mice were analyzed. Each image stack corresponded to a field of view of 60 \times 60 μ m and 5 μ m depth, and contained on average 4149 \pm 118 VGLUT1-positive boutons and 658 \pm 69 VGLUT2-positive boutons. The distance from VGLUT2 boutons to 5-HT_{4R} was shorter than the distance between VGLUT1 boutons and 5-HT_{4R} (*****p* < 0.0001, *t* test).

(B) 3D volume rendering of a dSPN dendrite labeled after injection in the DLS of a mix of AAV9-PPTA-CRE and AAV9-FLEX-tdTOMATO viruses with immunostained 5-HT_{4R}. The dendritic shaft was reconstructed in 3D by using Neuronstudio software, the dendrite model then imported into ImageJ and the headspines were segmented. Eight dendritic segments, from 3 mice, for a total of 423 μ m dendrite length with more than 1000 spines were analyzed. (Bottom left) 3D volume rendering of extracted dendritic shaft (red) and segmented spine head (a specific color is attributed to each spine head) with co-localizing 5-HT_{4R} (white). Arrowheads indicate examples of 5-HT_{4R} in shaft; arrows indicate 5-HT_{4R} on spines. (Right) Percentage of 5-HT_{4R} co-localizing with either dendritic shaft or spine head.

(C) 3D volume rendering showing 5-HT_{4R} (white) found in the contact between boutons from thalamic axons (green) and dendrite shaft (Top) or

inhibiting 5-HT₄R does not gate plasticity at corticostriatal synapses. Indeed, one might predict that the magnitude of LTD recorded during electrical stimulation experiments would be algebraically diluted by unaffected cortical synapses. This prediction would require the plasticity at cortical and thalamic inputs to be independent. However, we cannot exclude the possibility that the Ca²⁺ influx at cortical synapses that respond to electrical stimulation might influence thalamic synapses during the post-pre STDP. Although cortical synapses do not experience t-LTD upon 5-HT₄R inhibition, Ca²⁺-activated signaling molecules might diffuse from cortical spines to the dendritic shaft (Tang and Yasuda, 2017), where 5-HT₄R strongly co-localize with thalamic inputs (this study). This might affect local dendritic spikes and facilitate cooperative plasticity at thalamostriatal synapses (Golding et al., 2002; Popescu et al., 2007). An alternative scenario is that electrical stimulation may activate unidentified inputs contributing to the EPSP that also experience LTD upon inhibition of 5-HT signaling; this form of plasticity might be independent of that occurring at thalamostriatal inputs.

Our findings add to the limited knowledge of the intracellular signaling pathways regulating the thalamostriatal system, in particular the molecular mechanisms underlying synaptic depression of thalamo-dSPN synapses (Atwood et al., 2014; Ellender et al., 2011; Johnson et al., 2017; Parker et al., 2016; Wu et al., 2015). t-LTD at thalamostriatal inputs has been shown to be independent of eCBs released from the postsynaptic SPN (Wu et al., 2015). Consistent with this and with the evidence that thalamic afferents to the striatum are devoid of presynaptic CB1R (Wu et al., 2015), we found that t-LTD gated by 5-HT₄R antagonism is insensitive to CB1R inhibition and relies on postsynaptic expression mechanisms. In the DS, 5-HT₄R are localized mainly postsynaptically (Di Matteo et al., 2008). Activation of this G_s-coupled receptor subtype typically increases PKA activity and leads to the inhibition of K⁺ channels (King et al., 2008). Our findings point to an alternative signaling mechanism which involves CaMKII and activation of large-conductance BK channels. Indeed, 5-HT₄R can couple not only with G_s proteins, but also with G_α13, which signals to CaMKII (Liu and Voyno-Yasenetskaya, 2005; Nelson et al., 2005; Ponimaskin et al., 2002). In brain regions other than the striatum, CaMKII positively regulates BK channels (Nelson et al., 2005; Smith et al., 2002). We found that the paxilline-sensitive, BK channel-mediated current was significantly increased upon 5-HT₄R activation, and this enhancement was prevented by inhibition of CaMKII. The most parsimonious interpretation of these results is that 5-HT₄R activation leads to CaMKII activation, which, in turn, increases the amplitude of the BK channel-mediated current.

The post-pre STDP protocol used in this study (intra-striatal stimulation) may be reminiscent of behavioral states in which the striatal release of 5-HT (and other neuromodulators) occurs concomitantly with coordinated post- and presynaptic activity. In post-pre STDP, the EPSP coincides with the afterhyperpolarization (AHP) that follows the bAP (Feldman, 2012). Ultimately affecting bAP-induced Ca²⁺ transients, AHP is shaped by the activity of Ca²⁺-activated K⁺ conductances, including BK channels (Day et al., 2008). We propose a model in which 5-HT positively modulates BK channel activity via stimulation of 5-HT₄R during the post-pre STDP paradigm, ultimately reducing dendritic

bAP-induced Ca²⁺ influx (Figure 7). This may hinder the induction of t-LTD at thalamo-dSPN synapses, possibly by interfering with signaling pathways involved in the internalization of AMPAR (Grueter et al., 2010; Lüscher et al., 2000).

Conversely in this model, a decrease in 5-HT signaling during conventional post-pre STDP would dampen activation of BK channels and enhance bAP-induced Ca²⁺ signaling, thus promoting AMPAR internalization and enabling t-LTD (Grueter et al., 2010; Lüscher et al., 2000) (Figure 7). This could potentially mimic the circumstances *in vivo*, e.g., when post-pre STDP is uncorrelated to 5-HT release but occurs concomitantly with dopamine release or when there are pathological conditions characterized by hypofunctioning of the serotonergic system. Although our findings do not establish the precise identity of the thalamic afferents that are modulated by 5-HT, only thalamic inputs from the parafascicular thalamic (Pf) nucleus, and not from the central lateral (CL) nucleus, have been shown to express t-LTD (Ellender et al., 2013). Notably, Pf nucleus projections are the primary source for axo-dendritic synapses on SPNs (Smith et al., 2014), where we found 5-HT₄R to primarily localize, at least on dSPNs.

The 5-HT-mediated regulation of dendritic Ca²⁺ signaling during STDP represents a means by which changes in serotonergic levels may shape the thalamostriatal system. Projections of the intralaminar nuclei of the thalamus to the striatum are critical to interrupt ongoing motor behavior and redirect attention toward salient stimuli (Thorn and Graybiel, 2010), or to encoding A-O contingencies (Bradfield et al., 2013a, 2013b). Therefore, gating mechanisms of t-LTD at thalamic inputs that depend on serotonergic levels may affect the BG network and striatal information processing. Investigating the behavioral impact of 5-HT signaling specifically through the 5-HT₄R subtype, as well as establishing synaptic causalities with modulation of thalamostriatal inputs, is certainly an area for future studies.

Our results indicate that 5-HT at thalamic inputs, via 5-HT₄R, exerts a similar function as DA at cortical inputs, via D1R (Shen et al., 2008); both signaling pathways contribute to the activation of dSPNs by increasing the strength of glutamatergic inputs. The fact that the activation of D1R by DA released during intra-striatal stimulation (Shen et al., 2015; Threlfell et al., 2012; Wu et al., 2015) cannot compensate for the decrease in G_s signaling when 5-HT₄R are blocked suggests that D1R and 5-HT₄R may be localized in the different functional microdomains. Further studies are required to assess whether D1Rs segregate at cortical spines. Regardless, morphologically, 5-HT₄R are predominantly localized at axo-dendritic rather than axo-spinous synapses. Since only thalamic inputs, but not cortical afferents, form axo-dendritic synapses, this subcellular postsynaptic localization of 5-HT₄R (Figure 6) is consistent with the effects of 5-HT being specific to thalamic inputs. Building on existing models of striatal plasticity, we may now recognize excitatory DA and 5-HT as two complementary neuromodulatory systems controlling timing-dependent plasticity at two different glutamatergic inputs to dSPNs. This might allow for greater selectivity in tuning the activity of the direct pathway, which supports reinforcement and reward (Fisher et al., 2017; Kravitz et al., 2012).

Overall, the findings from this study not only provide mechanistic insights into the thalamic control of striatal functions, but




Figure 7. Proposed Model for 5-HT-Mediated Regulation of Postsynaptic Dendritic Ca²⁺ Signals during Timing-Dependent Plasticity

(A) Stimulation of 5-HT₄R during the post-pre STDP paradigm promotes phosphorylation of CaMKII at Thr286, thus boosting CaMKII activation, which in turn increases the function of BK channels during bursts of bAPs, ultimately reducing dendritic bAP-induced Ca²⁺ influx. This reduction would tip the balance in favor of t-LTD suppression.

(B) From the opposite view, in the presence of neuromodulatory afferent stimulation, it is sufficient to inhibit 5-HT release, or to prevent the activation of the 5-HT₄R subtype, to limit the activity of BK channels during the post-pre pairing, thus enhancing Ca²⁺ influx. This would promote the induction of t-LTD, which involves internalization of AMPAR.

they also identify synaptic substrates for maladaptive plasticity in BG disorders that involve dysfunction of the serotonergic system.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENTS AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
 - Animals
- METHOD DETAILS
 - Generation of the hM4Di knockin mouse line
 - *In vivo* brain microdialysis
 - Viral injections for optogenetic experiments
 - Electrophysiology
 - Imaging

- Immunofluorescence
- Postsynaptic localization of 5-HT₄R in dSPNs
- Immuno-EM analysis
- Activation of CaMKII in acute slices
- QUANTIFICATION AND STATISTICAL ANALYSIS
- DATA AND SOFTWARE AVAILABILITY

SUPPLEMENTAL INFORMATION

Supplemental Information includes nine figures and can be found with this article at <https://doi.org/10.1016/j.neuron.2018.04.008>.

ACKNOWLEDGMENTS

We are grateful to Mattia Pesce and Alice Gino for technical assistance during imaging experiments and immunohistochemistry, and to Alexis Bemelmans (Molecular Imaging Research Center, Commissariat pour l'Energie Atomique, France) for producing the adeno-associated virus PPTA-Cre. Funding: J.C. and N.H. acknowledge the Centre National pour la Recherche Scientifique (CNRS), Sorbonne University, the Fondation pour la Recherche Médicale

(# DEQ20150734352 to J.C.), and the Labex Bio-Psy Cluster of Excellence; A. Gozzi acknowledges the Brain and Behavior Foundation (2017 NARSAD independent Investigator Grant) and the Simons Foundation (SFARI 400101). M.P. acknowledges the Italian Ministry of Education and Progetti di Ricerca di Ateneo (PRA 2016) from University of Pisa. This research was primarily supported by the Fondazione Istituto Italiano di Tecnologia (to R.T.) and by grants provided by Compagnia di San Paolo (# 2013 0942 to R.T.) and Fondazione Cariplo (# 2013 0871 to R.T.).

AUTHOR CONTRIBUTIONS

A.C. and M.G. designed, performed and analyzed the electrophysiological experiments. M.G. conducted and analyzed the imaging experiments. A. Giorgi, M.P., and A. Gozzi designed and generated the *hM4Di^{+/+}/Pet1₂₁₀-Cre* mouse line. A.L. conducted and analyzed some electrophysiological recordings. M.G., A. Giorgi, M.T., A.L., and A.B. performed intracerebral viral injections. A.C., M.G., and M.T. performed immunostaining to identify dSPNs and iSPNs. S.M. and A.B. carried out the immunofluorescence analyses. M.M., G.M., and M.D.L. performed microdialysis in mice. N.H. carried out the 3D morphometric analysis with the help of M.G., and N.H. and J.C. conducted the immunohistochemical analysis on acute brain slices. T.C. and R.M. performed the electron microscopy analysis. A.C., M.G., A.L., and M.P. contributed to critical reading of the manuscript. M.P. supervised A. Giorgi, S.M., and A.B.; R.T. conceived the study, supervised the project, directed the experiments, and wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing financial interests.

Received: February 2, 2017

Revised: February 15, 2018

Accepted: April 5, 2018

Published: April 26, 2018

REFERENCES

- Atwood, B.K., Kupferschmidt, D.A., and Lovinger, D.M. (2014). Opioids induce dissociable forms of long-term depression of excitatory inputs to the dorsal striatum. *Nat. Neurosci.* *17*, 540–548.
- Bradfield, L.A., Bertran-Gonzalez, J., Chieng, B., and Balleine, B.W. (2013a). The thalamostriatal pathway and cholinergic control of goal-directed action: interlacing new with existing learning in the striatum. *Neuron* *79*, 153–166.
- Bradfield, L.A., Hart, G., and Balleine, B.W. (2013b). The role of the anterior, mediodorsal, and parafascicular thalamus in instrumental conditioning. *Front. Syst. Neurosci.* *7*, 51.
- Brebner, K., Wong, T.P., Liu, L., Liu, Y., Campsall, P., Gray, S., Phelps, L., Phillips, A.G., and Wang, Y.T. (2005). Nucleus accumbens long-term depression and the expression of behavioral sensitization. *Science* *310*, 1340–1343.
- Cai, X., Liang, C.W., Muralidharan, S., Kao, J.P., Tang, C.M., and Thompson, S.M. (2004). Unique roles of SK and Kv4.2 potassium channels in dendritic integration. *Neuron* *44*, 351–364.
- Carroll, R.C., Beattie, E.C., Xia, H., Lüscher, C., Altschuler, Y., Nicoll, R.A., Malenka, R.C., and von Zastrow, M. (1999). Dynamin-dependent endocytosis of ionotropic glutamate receptors. *Proc. Natl. Acad. Sci. USA* *96*, 14112–14117.
- Collingridge, G.L., Peineau, S., Howland, J.G., and Wang, Y.T. (2010). Long-term depression in the CNS. *Nat. Rev. Neurosci.* *11*, 459–473.
- Day, M., Wokosin, D., Plotkin, J.L., Tian, X., and Surmeier, D.J. (2008). Differential excitability and modulation of striatal medium spiny neuron dendrites. *J. Neurosci.* *28*, 11603–11614.
- Deng, P.Y., Rotman, Z., Blundon, J.A., Cho, Y., Cui, J., Cavalli, V., Zakharenko, S.S., and Klyachko, V.A. (2013). FMRP regulates neurotransmitter release and synaptic information transmission by modulating action potential duration via BK channels. *Neuron* *77*, 696–711.
- Di Matteo, V., Pierucci, M., Esposito, E., Crescimanno, G., Benigno, A., and Di Giovanni, G. (2008). Serotonin modulation of the basal ganglia circuitry: therapeutic implication for Parkinson's disease and other motor disorders. *Prog. Brain Res.* *172*, 423–463.
- Ding, J., Peterson, J.D., and Surmeier, D.J. (2008). Corticostriatal and thalamostriatal synapses have distinctive properties. *J. Neurosci.* *28*, 6483–6492.
- Ding, J.B., Guzman, J.N., Peterson, J.D., Goldberg, J.A., and Surmeier, D.J. (2010). Thalamic gating of corticostriatal signaling by cholinergic interneurons. *Neuron* *67*, 294–307.
- Doig, N.M., Moss, J., and Bolam, J.P. (2010). Cortical and thalamic innervation of direct and indirect pathway medium-sized spiny neurons in mouse striatum. *J. Neurosci.* *30*, 14610–14618.
- Dos Santos, M., Cahill, E.N., Bo, G.D., Vanhoutte, P., Caboche, J., Giros, B., and Heck, N. (2017). Cocaine increases dopaminergic connectivity in the nucleus accumbens. *Brain Struct. Funct.* *223*, 913–923.
- Ellender, T.J., Huerta-Ocampo, I., Deisseroth, K., Capogna, M., and Bolam, J.P. (2011). Differential modulation of excitatory and inhibitory striatal synaptic transmission by histamine. *J. Neurosci.* *31*, 15340–15351.
- Ellender, T.J., Harwood, J., Kosillo, P., Capogna, M., and Bolam, J.P. (2013). Heterogeneous properties of central lateral and parafascicular thalamic synapses in the striatum. *J. Physiol.* *591*, 257–272.
- Feldman, D.E. (2012). The spike-timing dependence of plasticity. *Neuron* *75*, 556–571.
- Fisher, S.D., Robertson, P.B., Black, M.J., Redgrave, P., Sagar, M.A., Abraham, W.C., and Reynolds, J.N.J. (2017). Reinforcement determines the timing dependence of corticostriatal synaptic plasticity in vivo. *Nat. Commun.* *8*, 334.
- Fornal, C.A., Metzler, C.W., Marrosu, F., Ribiero-do-Valle, L.E., and Jacobs, B.L. (1996). A subgroup of dorsal raphe serotonergic neurons in the cat is strongly activated during oral-buccal movements. *Brain Res.* *716*, 123–133.
- Giacomini, C., Mahajani, S., Ruffilli, R., Marotta, R., and Gasparini, L. (2016). Lamin B1 protein is required for dendrite development in primary mouse cortical neurons. *Mol. Biol. Cell* *27*, 35–47.
- Gilles, J.F., Dos Santos, M., Boudier, T., Bolte, S., and Heck, N. (2017). DiAna, an ImageJ tool for object-based 3D co-localization and distance analysis. *Methods* *115*, 55–64.
- Giorgi, A., Migliarini, S., Galbusera, A., Maddaloni, G., Mereu, M., Margiani, G., Gritti, M., Landi, S., Trovato, F., Bertozzi, S.M., et al. (2017). Brain-wide mapping of endogenous serotonergic transmission via chemogenetic fMRI. *Cell Rep.* *21*, 910–918.
- Goddard, A.W., Shekhar, A., Whiteman, A.F., and McDougle, C.J. (2008). Serotonergic mechanisms in the treatment of obsessive-compulsive disorder. *Drug Discov. Today* *13*, 325–332.
- Golding, N.L., Jung, H.Y., Mickus, T., and Spruston, N. (1999). Dendritic calcium spike initiation and repolarization are controlled by distinct potassium channel subtypes in CA1 pyramidal neurons. *J. Neurosci.* *19*, 8789–8798.
- Golding, N.L., Staff, N.P., and Spruston, N. (2002). Dendritic spikes as a mechanism for cooperative long-term potentiation. *Nature* *418*, 326–331.
- Grueter, B.A., Brasnjo, G., and Malenka, R.C. (2010). Postsynaptic TRPV1 triggers cell type-specific long-term depression in the nucleus accumbens. *Nat. Neurosci.* *13*, 1519–1525.
- Hannon, J., and Hoyer, D. (2008). Molecular biology of 5-HT receptors. *Behav. Brain Res.* *195*, 198–213.
- Hebert-Chatelain, E., Reguero, L., Puente, N., Lutz, B., Chaouloff, F., Rossignol, R., Piazza, P.V., Benard, G., Grandes, P., and Marsicano, G. (2014). Cannabinoid control of brain bioenergetics: Exploring the subcellular localization of the CB1 receptor. *Mol. Metab.* *3*, 495–504.
- Heck, N., Dos Santos, M., Amairi, B., Sallery, M., Besnard, A., Herzog, E., Boudier, T., Vanhoutte, P., and Caboche, J. (2015). A new automated 3D detection of synaptic contacts reveals the formation of cortico-striatal synapses upon cocaine treatment in vivo. *Brain Struct. Funct.* *220*, 2953–2966.

- Herzog, E., Nadrigny, F., Silm, K., Biesemann, C., Helling, I., Bersot, T., Steffens, H., Schwartzmann, R., Nägerl, U.V., El Mestikawy, S., et al. (2011). In vivo imaging of intersynaptic vesicle exchange using VGLUT1 Venus knock-in mice. *J. Neurosci.* *31*, 15544–15559.
- Higley, M.J., and Sabatini, B.L. (2010). Competitive regulation of synaptic Ca²⁺ influx by D2 dopamine and A2A adenosine receptors. *Nat. Neurosci.* *13*, 958–966.
- Hikida, T., Kimura, K., Wada, N., Funabiki, K., and Nakanishi, S. (2010). Distinct roles of synaptic transmission in direct and indirect striatal pathways to reward and aversive behavior. *Neuron* *66*, 896–907.
- Huerta-Ocampo, I., Mena-Segovia, J., and Bolam, J.P. (2014). Convergence of cortical and thalamic input to direct and indirect pathway medium spiny neurons in the striatum. *Brain Struct. Funct.* *219*, 1787–1800.
- Jacobs, B.L., and Fornal, C.A. (1999). Activity of serotonergic neurons in behaving animals. *Neuropsychopharmacology* *21* (2, Suppl), 9S–15S.
- Johnson, K.A., Mateo, Y., and Lovinger, D.M. (2017). Metabotropic glutamate receptor 2 inhibits thalamically-driven glutamate and dopamine release in the dorsal striatum. *Neuropharmacology* *117*, 114–123.
- Kilkenny, C., Browne, W.J., Cuthill, I.C., Emerson, M., and Altman, D.G. (2010). Improving bioscience research reporting: The ARRIVE guidelines for reporting animal research. *J. Pharmacol. Pharmacother.* *1*, 94–99.
- King, M.V., Marsden, C.A., and Fone, K.C. (2008). A role for the 5-HT_{1A}, 5-HT₄ and 5-HT₆ receptors in learning and memory. *Trends Pharmacol. Sci.* *29*, 482–492.
- Kravitz, A.V., Freeze, B.S., Parker, P.R., Kay, K., Thwin, M.T., Deisseroth, K., and Kreitzer, A.C. (2010). Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature* *466*, 622–626.
- Kravitz, A.V., Tye, L.D., and Kreitzer, A.C. (2012). Distinct roles for direct and indirect pathway striatal neurons in reinforcement. *Nat. Neurosci.* *15*, 816–818.
- Kreitzer, A.C., and Malenka, R.C. (2007). Endocannabinoid-mediated rescue of striatal LTD and motor deficits in Parkinson's disease models. *Nature* *445*, 643–647.
- Lee, S.J., Escobedo-Lozoya, Y., Szatmari, E.M., and Yasuda, R. (2009). Activation of CaMKII in single dendritic spines during long-term potentiation. *Nature* *458*, 299–304.
- Liu, G., and Voyno-Yasenetskaya, T.A. (2005). Radixin stimulates Rac1 and Ca²⁺/calmodulin-dependent kinase, CaMKII: cross-talk with Gα₁₃ signaling. *J. Biol. Chem.* *280*, 39042–39049.
- Lüscher, C., Nicoll, R.A., Malenka, R.C., and Muller, D. (2000). Synaptic plasticity and dynamic modulation of the postsynaptic membrane. *Nat. Neurosci.* *3*, 545–550.
- Mathur, B.N., and Lovinger, D.M. (2012). Serotonergic action on dorsal striatal function. *Parkinsonism Relat. Disord.* *18* (Suppl 1), S129–S131.
- Mathur, B.N., Capik, N.A., Alvarez, V.A., and Lovinger, D.M. (2011). Serotonin induces long-term depression at corticostriatal synapses. *J. Neurosci.* *31*, 7402–7411.
- Mereu, M., Tronci, V., Chun, L.E., Thomas, A.M., Green, J.L., Katz, J.L., and Tanda, G. (2015). Cocaine-induced endocannabinoid release modulates behavioral and neurochemical sensitization in mice. *Addict. Biol.* *20*, 91–103.
- Mereu, M., Contarini, G., Buonaguro, E.F., Latte, G., Managò, F., Iasevoli, F., de Bartolomeis, A., and Papaleo, F. (2017). Dopamine transporter (DAT) genetic hypofunction in mice produces alterations consistent with ADHD but not schizophrenia or bipolar disorder. *Neuropharmacology* *121*, 179–194.
- Migliarini, S., Pacini, G., Pelosi, B., Lunardi, G., and Pasqualetti, M. (2013). Lack of brain serotonin affects postnatal development and serotonergic neuronal circuitry formation. *Mol. Psychiatry* *18*, 1106–1118.
- Morishita, W., Marie, H., and Malenka, R.C. (2005). Distinct triggering and expression mechanisms underlie LTD of AMPA and NMDA synaptic responses. *Nat. Neurosci.* *8*, 1043–1050.
- Navailles, S., and De Deurwaerdère, P. (2011). Presynaptic control of serotonin on striatal dopamine function. *Psychopharmacology (Berl.)* *213*, 213–242.
- Nazzaro, C., Greco, B., Cerovic, M., Baxter, P., Rubino, T., Trusel, M., Parolaro, D., Tkatch, T., Benfenati, F., Pedarzi, P., and Tonini, R. (2012). SK channel modulation rescues striatal plasticity and control over habit in cannabinoid tolerance. *Nat. Neurosci.* *15*, 284–293.
- Nelson, A.B., Gittis, A.H., and du Lac, S. (2005). Decreases in CaMKII activity trigger persistent potentiation of intrinsic excitability in spontaneously firing vestibular nucleus neurons. *Neuron* *46*, 623–631.
- O'Hare, J.K., Ade, K.K., Sukharnikova, T., Van Hooser, S.D., Palmeri, M.L., Yin, H.H., and Calakos, N. (2016). Pathway-Specific Striatal Substrates for Habitual Behavior. *Neuron* *89*, 472–479.
- Parker, P.R., Lalive, A.L., and Kreitzer, A.C. (2016). Pathway-Specific Remodeling of Thalamostriatal Synapses in Parkinsonian Mice. *Neuron* *89*, 734–740.
- Pasqualetti, M., Ren, S.Y., Poulet, M., LeMour, M., Dierich, A., and Rijli, F.M. (2002). A Hoxa2 knockin allele that expresses EGFP upon conditional Cre-mediated recombination. *Genesis* *32*, 109–111.
- Pelosi, B., Migliarini, S., Pacini, G., Pratelli, M., and Pasqualetti, M. (2014). Generation of Pet1210-Cre transgenic mouse line reveals non-serotonergic expression domains of Pet1 both in CNS and periphery. *PLoS ONE* *9*, e104318.
- Plotkin, J.L., Shen, W., Rafalovich, I., Sebel, L.E., Day, M., Chan, C.S., and Surmeier, D.J. (2013). Regulation of dendritic calcium release in striatal spiny projection neurons. *J. Neurophysiol.* *110*, 2325–2336.
- Plotkin, J.L., Day, M., Peterson, J.D., Xie, Z., Kress, G.J., Rafalovich, I., Kondapalli, J., Gertler, T.S., Flajolet, M., Greengard, P., et al. (2014). Impaired TrkB receptor signaling underlies corticostriatal dysfunction in Huntington's disease. *Neuron* *83*, 178–188.
- Ponimaskin, E.G., Profirovic, J., Vaiskunaitė, R., Richter, D.W., and Voyno-Yasenetskaya, T.A. (2002). 5-Hydroxytryptamine 4(a) receptor is coupled to the Gα₁₃ subunit of heterotrimeric G13 protein. *J. Biol. Chem.* *277*, 20812–20819.
- Popescu, A.T., Saghyian, A.A., and Paré, D. (2007). NMDA-dependent facilitation of corticostriatal plasticity by the amygdala. *Proc. Natl. Acad. Sci. USA* *104*, 341–346.
- Shen, W., Flajolet, M., Greengard, P., and Surmeier, D.J. (2008). Dichotomous dopaminergic control of striatal synaptic plasticity. *Science* *321*, 848–851.
- Shen, W., Plotkin, J.L., Francardo, V., Ko, W.K., Xie, Z., Li, Q., Fieblinger, T., Wess, J., Neubig, R.R., Lindsley, C.W., et al. (2015). M4 Muscarinic Receptor Signaling Ameliorates Striatal Plasticity Deficits in Models of L-DOPA-Induced Dyskinesia. *Neuron* *88*, 762–773.
- Shindou, T., Ochi-Shindou, M., and Wickens, J.R. (2011). A Ca²⁺ threshold for induction of spike-timing-dependent depression in the mouse striatum. *J. Neurosci.* *31*, 13015–13022.
- Shuen, J.A., Chen, M., Gloss, B., and Calakos, N. (2008). Drd1a-tdTomato BAC transgenic mice for simultaneous visualization of medium spiny neurons in the direct and indirect pathways of the basal ganglia. *J. Neurosci.* *28*, 2681–2685.
- Sippy, T., Lapray, D., Crochet, S., and Petersen, C.C. (2015). Cell-type-specific sensorimotor processing in striatal projection neurons during goal-directed behavior. *Neuron* *88*, 298–305.
- Smith, M.R., Nelson, A.B., and Du Lac, S. (2002). Regulation of firing response gain by calcium-dependent mechanisms in vestibular nucleus neurons. *J. Neurophysiol.* *87*, 2031–2042.
- Smith, Y., Galvan, A., Ellender, T.J., Doig, N., Villalba, R.M., Huerta-Ocampo, I., Wichmann, T., and Bolam, J.P. (2014). The thalamostriatal system in normal and diseased states. *Front. Syst. Neurosci.* *8*, 5.
- Südhof, T.C. (2012). The presynaptic active zone. *Neuron* *75*, 11–25.
- Tang, S., and Yasuda, R. (2017). Imaging ERK and PKA activation in single dendritic spines during structural plasticity. *Neuron* *93*, 1315–1324.
- Tecuapetla, F., Jin, X., Lima, S.Q., and Costa, R.M. (2016). Complementary contributions of striatal projection pathways to action initiation and execution. *Cell* *166*, 703–715.

- Thorn, C.A., and Graybiel, A.M. (2010). Pausing to regroup: thalamic gating of cortico-basal ganglia networks. *Neuron* 67, 175–178.
- Threlfell, S., Lalic, T., Platt, N.J., Jennings, K.A., Deisseroth, K., and Cragg, S.J. (2012). Striatal dopamine release is triggered by synchronized activity in cholinergic interneurons. *Neuron* 75, 58–64.
- Tritsch, N.X., and Sabatini, B.L. (2012). Dopaminergic modulation of synaptic transmission in cortex and striatum. *Neuron* 76, 33–50.
- Trusel, M., Cavaccini, A., Gritti, M., Greco, B., Saintot, P.P., Nazzaro, C., Cerovic, M., Morella, I., Brambilla, R., and Tonini, R. (2015). Coordinated Regulation of Synaptic Plasticity at Striatopallidal and Striatonigral Neurons Orchestrates Motor Control. *Cell Rep.* 13, 1353–1365.
- van Welie, I., and du Lac, S. (2011). Bidirectional control of BK channel open probability by CAMKII and PKC in medial vestibular nucleus neurons. *J. Neurophysiol.* 105, 1651–1659.
- Vicente, A.M., Galvão-Ferreira, P., Tecuapetla, F., and Costa, R.M. (2016). Direct and indirect dorsolateral striatum pathways reinforce different action strategies. *Curr. Biol.* 26, R267–R269.
- Wu, Y.W., Kim, J.I., Tawfik, V.L., Lalchandani, R.R., Scherrer, G., and Ding, J.B. (2015). Input- and cell-type-specific endocannabinoid-dependent LTD in the striatum. *Cell Rep.* 10, 75–87.
- Yagishita, S., Hayashi-Takagi, A., Ellis-Davies, G.C., Urakubo, H., Ishii, S., and Kasai, H. (2014). A critical time window for dopamine actions on the structural plasticity of dendritic spines. *Science* 345, 1616–1620.
- Zhai, S., Tanimura, A., Graves, S.M., Shen, W., and Surmeier, D.J. (2018). Striatal synapses, circuits, and Parkinson's disease. *Curr. Opin. Neurobiol.* 48, 9–16.
- Zhou, Y.D., Acker, C.D., Netoff, T.I., Sen, K., and White, J.A. (2005). Increasing Ca²⁺ transients by broadening postsynaptic action potentials enhances timing-dependent synaptic depression. *Proc. Natl. Acad. Sci. USA* 102, 19121–19125.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit polyclonal antibody to A2A	Enzo Life Sciences	Cat# BML-SA654; RRID: AB_2226507
Rat monoclonal antibody to substance P	Millipore	Cat# MAB356; RRID: AB_94639
Alexa 488 anti-rabbit	ThermoFisher Scientific	Cat# A-11034; RRID: AB_2576217
Alexa 647 anti-rat	ThermoFisher Scientific	Cat# A-21247; RRID: AB_141778
Rabbit anti-RFP antibody	AbCam	Cat# ab62341; RRID: AB_945213
Chicken polyclonal anti-eGFP	AbCam	Cat# ab13970; RRID: AB_300798
Rat monoclonal anti-5HT	Millipore	Cat# MAB352 RRID: AB_94865
Alexa Fluor 488 goat anti-chicken IgG	Molecular Probes	Cat# A-11039; RRID: AB_142924
VectaFluor Excel Amplified DyLight 594 Anti-Rabbit IgG Kit	Vector Laboratories	Cat# DK-1594; RRID: AB_2336778
Alexa Fluor 488 donkey anti-rat IgG	Molecular Probes	Cat# A-21208; RRID: AB_141709
Rabbit anti-5-HT4R	AbCam	Cat# ab60359; RRID: AB_2122438
Mouse anti-bassoon	AbCam	Cat# ab82958; RRID: AB_1860018
Goat anti-rabbit Cy3-conjugated	Jackson	Cat# 111-165-003; RRID: AB_2338000
Goat anti-rabbit Cy5-conjugated	Jackson	Cat# 111-175-144; RRID: AB_2338013
Goat anti-mouse Cy5-conjugated	Jackson	Cat# 115-175-205; RRID: AB_2338715
Rat monoclonal anti-mCherry	ThermoFisher Scientific	Cat# M11217; RRID: AB_2536611
Rabbit polyclonal anti-5HT	ThermoFisher Scientific	Cat# PA1-36157; RRID: AB_1087251
Rabbit polyclonal anti-SERT	Millipore	Cat# AB9726; RRID: AB_612176
Goat anti-rabbit biotinylated secondary antibody	Vector Laboratories	Cat# BA-1000; RRID: AB_2313606
1.4 nm gold-labeled goat anti-rat Immunoglobulin-G (Fab' fragment)	Nanoprobes	Cat# 2008; RRID N/A
Alexa 568-conjugated streptavidin	ThermoFisher Scientific	Cat# S-11226; RRID: AB_2315774
Rabbit anti-pCaMKII (Thr286)	Cell Signaling Technology	Cat# 3361; RRID: AB_10015209

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
rAAV5-CaMKIIa-ChR2(H134R)-eYFP	UNC Vector Core	AV43160
rAAV5-EF1a-DIO-hChR2(H134R)-eYFP	UNC Vector Core	AV4313X
AAV9.CAG.Flex.eGFP.WPRE.bGH	Upenn, Dr. Hongkui Zeng	AV-9-ALL854
AAVrh9-PPTA-cre.WPRE	Laboratory of Dr. J. Caboche	N/A
AAV9.CAG.Flex.tdTomato.WPRE.bGH	Upenn, Dr. Hongkui Zeng	AV-9-ALL864
Biological Samples		
Chemicals, Peptides, and Recombinant Proteins		
Fluo-4	ThermoFisher Scientific	Cat# F14200
Alexa Fluor 568 Hydrazide	ThermoFisher Scientific	Cat# A10437
Neurobiotin	Vector Laboratories	Cat# SP-1120; RRID: AB_2313575
ProLong Gold AntiFade reagent	ThermoFisher Scientific	Cat# P36935
SR95531	Tocris Bioscience	Cat# 1262; CAS Number: 104104-50-9
SB271046	Tocris Bioscience	Cat# 3368; CAS Number: 209481-24-3
GR113808	Tocris Bioscience	Cat# 1322; CAS Number: 144625-51-4
RS39604	Tocris Bioscience	Cat# 0991; CAS Number: 167710-87-4
AM251	Tocris Bioscience	Cat# 1117; CAS Number: 183232-66-8
Paxilline	Tocris Bioscience	Cat# 2006; CAS Number: 57186-25-1
RS67333	Tocris Bioscience	Cat# 0989; CAS Number: 168986-60-5
AIP	Tocris Bioscience	Cat# 1688; CAS Number: 167114-91-2
PKI 5-24	Tocris Bioscience	Cat# 6221; CAS Number: 99534-03-9
KN-62	Tocris Bioscience	Cat# 1277; CAS Number: 127191-97-3
R-CPP	Tocris Bioscience	Cat# 2334; CAS Number: 126453-07-4
D15	Tocris Bioscience	Cat# 2334; CAS Number: 251939-41-0; Sequence: PPPQVPSRPNRAPPG
S15	TermoFisher Scientific	Sequence: ANVRRGPPPPQPSP
GluA23Y	TermoFisher Scientific	Sequence: YKEGYNVYV
GluA23Y Scramble (GluA23A)	TermoFisher Scientific	Sequence: AKEGANVAG
CNO (Clozapine-N-Oxide)	Sigma Aldrich	C0832; CAS Number: 34233-69-7
Critical Commercial Assays		
HQ Silver kit	Nanoprobes	Cat# 2012
avidin-biotin complex	Vector Laboratories	Cat# PK-6100; RRID: AB_2336819
Deposited Data		
Analyzed Data	This paper	N/A
Experimental Models: Cell Lines		
Mouse: ESCs	BayGenomics	E14Tg2a.4
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	Charles River	C57BL/6J; RRID: IMSR_JAX:000664
Mouse: hM4Di ^{+/-} /Pet1 ₂₁₀ -Cre	Laboratory of Prof. Massimo Pasqualetti	N/A
Mouse: hM4Di ^{+/-} /Pet1 ₂₁₀ -Cre ⁰	Laboratory of Prof. Massimo Pasqualetti	N/A
Mouse: Pet1 ₂₁₀ -Cre	Laboratory of Prof. Massimo Pasqualetti	N/A
Mouse: B6.Cg-Tg(Thy1-COP4/EYFP)18Gfng/J	The Jackson Laboratory	Cat# JAX:007612; RRID: IMSR_JAX:007612
Mouse: hM4Di ^{+/-} /Pet1 ₂₁₀ -Cre/Tph2 ^{GFP+/-}	Laboratory of Prof. Massimo Pasqualetti	N/A
Mouse: <i>Drd1a</i> -td Tomato BAC transgenic mouse line 5, and backcrossed to C57BL/6J	Laboratory of Dr. N.Calagos	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Mouse: VGLUT1 ^{Venus} knock-in	Laboratory of Dr. Etienne Herzog; Laboratory of Dr. J. Caboche	N/A
Mouse: Slc17a6 ^{tm2(cre)Lowl} /J (VGLUT2-Cre)	The Jackson Laboratory	Cat# 016963; RRID: IMSR_JAX:016963
Oligonucleotides		
Primer: wild-type <i>ROSA26</i> allele forward 5'-GAGGGGAGTGTTCGAATACC-3'	BMR Genomics	N/A
Primer: wild-type <i>ROSA26</i> allele reverse 5'-AGTCTAACTCGCGACTGTGA-3'	BMR Genomics	N/A
Primer: reverse 5'-GTCCCTATTGGCGTTACTATG-3'	BMR Genomics	N/A
Recombinant DNA		
Plasmid: pCX-CAG-eGFP	Pasqualetti et al., 2002	https://doi.org/10.1002/gene.10053
Plasmid: pROSA26-1 ^{Sor}	Soriano P. Nat Genet. 1999	AddGene: #21714; https://doi.org/10.1038/5007
Plasmid: pPSVKeoFRT	Pasqualetti et al., 2002	https://doi.org/10.1002/gene.10053
Vector: pROSA26-1 ^{Sor} -CAG-hM4Di/mCherry	This paper	N/A
Plasmid: pAAV-hSyn-DIO-hM4Di/mCherry	Krashes et al., J Clin Invest. 2011	AddGene: #44362; https://doi.org/10.1172/JCI46229 .
Software and Algorithms		
GraphPad Prism 6 software	GraphPad Software, Inc.	https://www.graphpad.com/ ; RRID: SCR_002798
pClamp 10	Molecular Devices	http://mdc.custhelp.com/ ; RRID: SCR_011323
ImageJ	NIH	https://imagej.nih.gov/ij/ ; RRID: SCR_003070
Neuronstudio software	Computational Neurobiology and Imaging Center	http://research.mssm.edu/cnic/tools-ns.html ; RRID: SCR_013798
Huygens software (Maximum Likelihood Estimation algorithm)	Scientific Volume Imaging	https://svi.nl/HuygensSoftware ; RRID: SCR_014237
Origin 9.1	OriginLab Corporation	https://store.originlab.com/store/Default.aspx? CategoryID=0 ; RRID: SCR_014212
Adobe Photoshop CS6	Adobe Systems	https://www.adobe.com/ ; RRID: SCR_014199
Other		
AN69 dialyzing membranes	Hospal Dasco	Nephral ST 500

CONTACT FOR REAGENTS AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Dr. Raffaella Tonini (raffaella.tonini@iit.it). Distribution of the Drd1a-td Tomato BAC transgenic mouse line 5 is restricted by MTA.

EXPERIMENTAL MODEL AND SUBJECT DETAILS**Animals**

All procedures involving animals were carried out in accordance with the Italian Ministry of Health's directives (D.lgs 26/2014) regulating animal research. Animal experiments were designed in accordance with the ARRIVE (Animal Research: Reporting of *In Vivo* Experiments) guidelines (Kilkenny et al., 2010), with a commitment to refinement, reduction, and replacement, minimizing the number of mice. Drug, and test naive male mice at postnatal day 35-80 were housed 4-5 per cage in a temperature- and humidity-controlled room under a 12:12 h light/dark cycle with lights on at 07:00, and fed regular laboratory chow with *ad libitum* access to water. The following mouse strains were used in this study: C57BL/6J (Charles River Laboratories; RRID: IMSR_JAX:000664), hM4Di^{+/-}/Pet1₂₁₀-Cre, hM4Di^{+/-}/Pet1₂₁₀-Cre⁰, Pet1₂₁₀-Cre, hM4Di^{+/-}/Pet1₂₁₀-Cre/Tph2^{GFP+/-} (Laboratory of Prof. Massimo Pasqualetti); B6.Cg-Tg(Thy1-COP4/EYFP)18Gfng/J (The Jackson Laboratory; RRID: IMSR_JAX:007612), Slc17a6^{tm2(cre)Lowl}/J (VGLUT2-Cre) (The Jackson Laboratory; RRID: IMSR_JAX:016963); Drd1a-td Tomato BAC transgenic mouse line 5 (Laboratory of Dr. N.Calakos) backcrossed to C57BL/6J; VGLUT1^{Venus} knockin (Laboratory of Dr. Etienne Herzog and Laboratory of Dr. J. Caboche). For all strains, male mice were routinely bred with C57BL/6J females, with the exception of VGLUT2-Cre and VGLUT1^{Venus}, which were maintained in homozygosis. For production of hM4Di^{+/-}/Pet1₂₁₀-Cre animals, hM4Di^{+/-} adult males were mated to Pet1₂₁₀-Cre transgenic females; progeny was genotyped for the presence of both hM4Di in the ROSA26 locus and Cre. For production of

hM4Di^{+/-}/Pet1₂₁₀-Cre/Tph2^{GFP+/-} mice, hM4Di^{+/-}/Pet1₂₁₀-Cre males were mated to Tph2^{GFP+/-} females. Progeny was genotyped for the presence of hM4Di in the ROSA26 locus, Cre, and GFP within the Tph2 locus.

METHOD DETAILS

Generation of the hM4Di knockin mouse line

We generated the hM4Di^{DIO} conditional knockin mouse line taking advantage of a gene targeting strategy allowing the insertion of the construct via homologous recombination within the ROSA26 genomic locus. The in-frame fusion protein resulting from the hM4Di DREADD receptor and the mCherry reporter coding sequences was flanked by two couples of Lox sites (LoxP/Lox2722). The resulting cassette was isolated from the pAAV-hSyn-DIO-hM4Di/mCherry vector and cloned in the 3'-5' orientation within the pCX-CAG-eGFP plasmid downstream of the CAG promoter by replacing the eGFP sequence. Subsequently, the CAG-hM4Di/mCherry fragment was isolated from pCX-CAG-hM4Di/mCherry via SpeI/XbaI double digestion and cloned in the XbaI site of the pROSA26-1^{Sor} plasmid that had been previously modified by the addition of a PvuI recognition site downstream to the *pgk-DTa* cassette in place of the unique XhoI site. Finally, the SpeI/NheI fragment containing the pgk-Neo/Kana cassette was obtained from the pPSVKeoFRT plasmid and cloned into the unique XbaI site of the pROSA26-1^{Sor}-CAG-hM4Di/mCherry vector. PvuI-linearized targeting vector was electroporated into E14Tg2a.4 ES cells and G418 selection was maintained for 7 days as described (Giorgi et al., 2017). Among 192 G418-resistant ES cell colonies screened, 32 were positive recombinant clones as assessed by Southern blot analysis. Following karyotype analysis, recombinant ES cells were microinjected into C57BL/6J host blastocysts that were later reimplanted into pseudopregnant CD1 females. Germline transmission was achieved for the 60% of the 29 chimeras obtained. hM4Di^{DIO} mice were born at normal Mendelian ratios and lived a normal lifespan with no obvious defects.

Animals were genotyped using DNA primers as follows: wild-type ROSA26 allele forward 5'-GAGGGGAGTGTTCGAATACC-3' and reverse 5'-AGTCTAACTCGCGACTGTA-3'. The hM4Di^{DIO} knockin allele was genotyped using the same forward primer as above in combination with the reverse primer 5'-GTCCCTATTGGCGTTACTATG-3'.

In vivo brain microdialysis

hM4Di^{+/-}/Pet1₂₁₀-Cre mice were anesthetized with Avertin (1.5 mg/kg) and placed in a stereotaxic frame (2B, 2biological Instruments) for probe implantation. Concentric dialysis probes were implanted either in the right or left dorsolateral striatum (DLS) (uncorrected coordinates +0.38 AP, ± 2.0 ML, -3.8 DV). After surgery, mice were allowed to recover overnight in square Plexiglas cages (Allentown). Concentric dialysis probes were prepared using AN69 dialyzing membranes (Hospal Dasco, Bologna, Italy) as previously described (Mereu et al., 2017; Mereu et al., 2015). The exposed dialyzing surface of the membrane was limited to the lowest 2.00 mm portion of the probes, which were less than 18 mm in total length. Experiments were performed in freely moving animals in the same cage in which they recovered from surgery. Microdialysis test sessions started at 9 AM, approximately 22-24 hours after surgical procedures. Probes were connected to PE50 tubes and perfused with Ringer's solution (147.0 mM NaCl, 2.2 mM CaCl₂ and 4.0 mM KCl) delivered through the dialysis probes by a 1.0 ml syringe, operated by a BAS BeeSyndrome Pump Controller (BAS West Lafayette, IN, USA), at a constant flow rate of 1 µl/min. Collection of dialysate samples (20 µl) started after about 30 min, and extracellular serotonin levels were quantified using an HPLC apparatus equipped with a reverse-phase column (C83.5 µm, Waters, Milford, USA) and a coulometric electrochemical detector (ESA Coulochem II, Bedford, USA). At the end of each experiment, mice were sacrificed and their brains were removed and stored in formalin (4%) before histological analysis. Brains were cut in serial coronal slices (100 µm) to locate the placement of the microdialysis probe.

Viral injections for optogenetic experiments

Five- to six-week-old mice were anesthetized (2% isoflurane) and placed in a stereotaxic frame. To measure light-induced EPSPs at cortico- and thalamostriatal connections on dSPNs and iSPNs, C57BL/6J and hM4Di^{+/-}/Pet1₂₁₀-Cre mice were injected with rAAV5-CaMKIIa-ChR2(H134R)-eYFP virus (UNC Vector Core, Chapel Hill, North Carolina). The motor cortex M1 (+1.1 AP; ± 1.6 ML; -0.8 DV from brain surface) or the thalamus was targeted (-2.3 AP; ± 0.75 ML; -3.5 DV). A volume of 0.5 µl/site of virus was injected at a rate of 0.1 µl/min. To measure the effect of the release of serotonin on electrically-induced EPSPs at thalamostriatal connections on dSPNs, Pet1₂₁₀-Cre mice were injected with rAAV5-EF1a-DIO-hChR2(H134R)-eYFP virus (UNC Vector Core, Chapel Hill, North Carolina). The dorsal raphe nucleus (DRN) was targeted (-4.36 AP; ± 0.5 ML; -3, -4 DV). A volume of 0.8 µl/site of virus was injected at a rate of 0.1 µl/min. Mice were used for electrophysiological recordings 4-6 weeks after the injection.

Electrophysiology

Slice preparation

Mice were anesthetized with isoflurane and decapitated, and their brains were transferred to ice-cold dissecting artificial cerebrospinal fluid (aCSF) containing 110 mM choline chloride, 2.5 mM KCl, 1.25 mM NaH₂PO₄, 7 mM MgSO₄, 0.5 mM CaCl₂, 25 mM NaHCO₃, 25 mM D-glucose, 11.6 mM sodium-L-ascorbate and 3.1 mM sodium pyruvate, saturated with 95% O₂ and 5% CO₂. Coronal sections (250 µm thick), horizontal slices (270 µm thick) and para-horizontal slices (20° oblique) were cut using a Vibratome 1000S (Leica, Wetzlar, Germany), then transferred to aCSF containing 115 mM NaCl, 3.5 mM KCl, 1.2 mM NaH₂PO₄, 1.3 mM MgCl₂, 2 mM CaCl₂, 25 mM NaHCO₃ and 25 mM D-glucose and aerated with 95% O₂ and 5% CO₂. Following 20 min of incubation

at 32°C, slices were kept at 22–24°C. During experiments, slices were continuously superfused with aCSF at a rate of 2 ml/min at 28°C.

Electrophysiological recordings

Whole-cell patch-clamp recordings were made on striatal projection neurons (SPNs) of the dorsolateral striatum (DLS) in horizontal or para-horizontal slices, or in neurons of the dorsal raphe nucleus (DRN) in coronal slices. Intracellular solution contained 130 mM KMeSO₄, 5 mM KCl, 5 mM NaCl, 10 mM HEPES, 0.1 mM EGTA, 2 mM MgCl₂, 0.05 mM CaCl₂, 2 mM Na₂-ATP and 0.4 mM Na₃-GTP (pH 7.2–7.3, 280–290 mOsm/kg).

In the DLS, excitatory postsynaptic potentials (EPSPs) were induced at 0.1 Hz in the presence of the GABA_A receptor antagonist gabazine (10 μM) by intra-striatal electrical stimulation using a concentric bipolar electrode (40–100 μA; 40–80 μs; FHC, Bowdoin, Maine) or by optogenetic stimulation of cortical or thalamic afferents using short blue light pulses (range 0.1–1 ms at 470 nm) of LED light delivered through the microscope objective (CoolLED, pE-100, Andover, Hampshire, United Kingdom). STDP-LTD was induced using a protocol consisting of 20 bouts of EPSPs paired with bAPs, delivered 10 s apart. Each bout consisted of five bursts (120 ms apart), each composed of three bAPs at 50 Hz followed by one EPSP (negative timing). The onset of the EPSPs followed the peak of the last postsynaptic action potential in the burst by 10 ms ($\Delta t = 10$ ms). During plasticity induction, postsynaptic neurons were depolarized from –80 mV to –70 mV. Data were excluded when the input resistance (R_{in}) changed > 20%. In *Pet1₂₁₀-Cre* mice expressing ChR2 in the DRN, thalamic afferents were stimulated by placing the stimulation electrode in the thalamus close to the border with the thalamic reticular nucleus. To induce the release of serotonin from DRN afferents in the DLS, blue light pulses (10 ms duration at 470 nm) of LED light were delivered through the microscope objective (CoolLED, pE-100, Andover, Hampshire, United Kingdom) at the time of presynaptic stimulation evoked during the STDP protocol. BK channel-mediated current was recorded in voltage-clamp configuration in response to a depolarizing voltage ramp from –80 to 110 mV (rate 0.1 mV/ms). Data are reported without corrections for liquid junction potentials. Data were acquired using a Multiclamp 700B amplifier controlled by pClamp 10 software (Molecular Device), filtered at 2.5 kHz and sampled at 10 kHz (voltage clamp), or filtered at 10 kHz and sampled at 20 kHz (current-clamp) with a (Digidata 1322, Molecular Device). For statistical purposes (RM2WA analysis; Figure S5B), data were filtered at 0.1 kHz.

Imaging

Ca²⁺ transients were imaged with 100 μM Fluo-4 (ThermoFisher Scientific, Carlsbad, California) dissolved in the intracellular recording solution. Alexa 568 (10 μM, ThermoFisher Scientific, Carlsbad, California) was used for visualization of cell bodies and dendrites. SPNs were filled with Alexa 568 and Fluo-4 via the patch electrode for 20 min before imaging to allow dye equilibration in the proximal dendrites. Whole-cell maximum-projection images of the soma and dendrites were acquired with 0.4 μm² pixels with 15 μs pixel dwell time; ~120 images were taken with 0.4 μm focal steps. Drugs were bath applied for 10 min.

Two-photon Ca²⁺ imaging was performed with a Leica SP5 AOBS upright DM6000 CFS microscope coupled with a 2P laser Chameleon Ultra Coherent and equipped with a Leica 25x NA 0.95 water immersion objective. The fluorescence emission was collected with an external non-descanned photomultiplier detector equipped with a 525/50 emission filter. Green fluorescence line-scan signals were acquired at 6 ms and 256 pixels per line, with 0.1 μm pixels and 20 μs pixel dwell time. The laser-scanned images were acquired with 800 nm light pulsed at 90 MHz (pulse duration: ~250 fs). The line scan was used to record the fluorescence elicited by one bout of the STDP protocol. Images were collected with LAS AF Leica software and analyzed using ImageJ software (version 1.45, <https://imagej.nih.gov/ij/>, NIH, Maryland) and Origin 9.1 (OriginLab Corporation, Northampton, Massachusetts). For each recording, background fluorescence was determined from a cell-free area of comparable size to that of the line-scan image. After subtracting the averaged background signal, fluorescence values were recorded for 200 ms before the triggering the STDP protocol and averaged to give the basal fluorescence (F_{basal}). The amplitude of the fluorescence transients at the recording sites was expressed as the fractional change in basal fluorescence, $(F - F_{\text{basal}})/F_{\text{basal}}$, ($\Delta F/F$), which is approximately proportional to the changes in intracellular Ca²⁺. Over the course of the experiment, F_{basal} remained within the standard error of F_{basal} measured under control conditions. For data analysis, transients were digitally filtered offline (adjacent-averaging routine, smoothing factor $n = 5$; Origin 9.1).

Immunofluorescence

Identification of dSPNs and iSPNs

To identify SPNs of the direct (dSPN) and indirect (iSPN) pathways, neurons were filled with Neurobiotin (0.5 mg/ml, Vector Laboratories, Peterborough, United Kingdom) during recordings and subsequently processed for immunostaining of the A2A receptor (marker of iSPNs) and substance P (marker of dSPNs) (Nazzaro et al., 2012; Trusel et al., 2015). After recording, slices were fixed with 4% paraformaldehyde in 0.1 M phosphate buffer (0.1 M PB; pH 7.4) overnight at 4°C and then incubated in primary antibodies. Rabbit polyclonal antibody to A2A (1:250, Enzo Life Sciences, Farmingdale, New York) and rat monoclonal antibody to substance P (1:200, Millipore, Billerica, Massachusetts) were diluted in 0.1 M PB containing 0.3% Triton X-100. Sections were subsequently incubated with Alexa 647- or Alexa 488-conjugated secondary antibodies (1:200) and Alexa 568-conjugated streptavidin (1:1000) (ThermoFisher Scientific, Carlsbad, California), mounted on glass slides, and coverslipped with ProLong Gold antifade reagent (ThermoFisher Scientific, Carlsbad, California). Images were acquired with an inverted Leica TCS SP5 AOBS TANDEM confocal microscope.

5-HT, SERT and mCherry expression

hM4D^{+/−}/Pet1₂₁₀-Cre/Tph2^{GFP+/−} or Pet1₂₁₀-Cre mice were perfused with 4% paraformaldehyde (w/v) in phosphate buffer (0.1 M PB; pH 7.4) and vibratome sections (50 μm) were processed for immunofluorescence (Migliarini et al., 2013). After blocking and permeabilization, slices were incubated in primary antibody diluted in PB/Triton solution (0.1 M PB, 0.5% Triton X-100, 5% horse serum) [rabbit anti-RFP (1:500, Abcam, Cambridge, United Kingdom); chicken polyclonal anti-eGFP (1:1000, Abcam, Cambridge, United Kingdom); rat monoclonal anti-5-HT (1:200, Merck Millipore, Darmstadt, Germany)] overnight at 4°C. Rabbit anti-SERT (1:500, Millipore, Darmstadt, Germany) primary antibody was diluted in modified PB/Triton (0.1 M PB, 0.3% Triton X-100, 10% horse serum) and slices incubated for 36 h at 4°C. Slices were then incubated in secondary antibody [Alexa Fluor 488 goat anti-chicken IgG (1:500, Molecular Probes, Eugene, Oregon); VectaFluor Excel Amplified DyLight® 594 Anti-Rabbit IgG Kit; Alexa Fluor 488 donkey anti-rat IgG (1:500, Molecular Probes, Eugene, Oregon)] for 2 h at room temperature and mounted onto slides for imaging. Images were acquired with a MacroFluo microscope (Leica, Wetzlar, Germany) with a DS-SMc digital camera (Nikon, Tokyo, Japan) or with an A1 confocal microscope (Nikon, Tokyo, Japan).

Postsynaptic localization of 5-HT4R in dSPNs

Labeling of cortical and thalamic afferents

To identify cortical afferents, VGLUT1-Venus knockin mice were used and no injections were required (Heck et al., 2015; Herzog et al., 2011). To label thalamic afferents, AAV9.CAG.Flex.eGFP.WPRE.bGH (UPenn) was injected in the thalamus of VGLUT2-Cre mice (−2.3 AP; ± 0.75 LM; −3.5 DV). A volume of 0.5 μl/site of virus (diluted 1:10 in PBS) was injected at a rate of 0.1 μl/min. Mice were perfused 3–4 weeks after the injection. To label dSPNs, the viral construct AAVrh9-PPTA-CRE, which contains an expression cassette consisting of Cre recombinase driven by the promoter of the PPTA gene (substance P), which is specifically expressed in dSPNs, was co-injected with AAV9.CAG.Flex.tdTomato.WPRE.bGH (UPenn) expressing “flexed” tdTomato under the CMV/actin hybrid promoter (CAG). The DLS was targeted (+0.5 AP; ± 2.6 LM; −2.9 DV) and a volume of 2 μl/site of the mix was injected at a rate of 0.1 μl/min. Mice were perfused 2–3 weeks after the injection (Dos Santos et al., 2017; Hikida et al., 2010). In VGLUT2-Cre mice, double injections of the AAVrh9-PPTA-CRE/AAV9.CAG.Flex.tdTomato.WPRE.bGH mix in the DLS and of AAV9.CAG.Flex.eGFP.WPRE.bGH in the thalamus were also performed. Mice were perfused 3 weeks after the injection. The tdTomato- and eGFP-expressing viruses were made available by Dr. Hongkui Zeng of the Allen Institute for Brain Science.

Mouse perfusion and preparation of vibratome sections

Mice were deeply anesthetized and intracardially perfused with 4% PFA in 0.1 M phosphate buffer, pH 7.4 at 4°C. The brain was removed from the skull and placed in 4% PFA overnight at 4°C. Sections of 30 or 50 μm were cut in the frontal plane using a vibratome 1000S (Leica, Wetzlar, Germany), and stored in PBS.

Immunocytochemical labeling

Sections were permeabilized for 15 min in 0.5% Triton X-100, blocked for 1 h at room temperature in PBS containing 0.01% Triton X-100, 3% BSA and 3% NGS. The sections were incubated 48 hours at 4°C with primary antibodies (rabbit anti-5-HT4R: 1:500, Abcam; mouse anti-bassoon: 1:250, Abcam) in PBS with 0.01% Triton X-100, 3% BSA and 3% NGS. After washing, sections were incubated for 90 min at room temperature with secondary antibodies (goat anti-rabbit Cy3-conjugated or Cy5-conjugated; goat anti-mouse Cy5-conjugated; 1:500, Jackson) in PBS with 0.01% Triton X-100, 3% BSA and 3% NGS. Sections were washed before mounting in Prolong Gold.

Confocal image acquisition and deconvolution

Images stacks were acquired with a Confocal Laser Scanning Microscope (SP5, Leica, Wetzlar, Germany), through an 63x oil immersion objective with numerical aperture of 1.4 (Leica, Wetzlar, Germany) using 700Hz scanning frequency and pinhole aperture set to 1 Airy Unit. Stacks of 25 images (1024x1024 pixels) were acquired with a pixel size of 60 nm and z-step of 200 nm. Excitation (ex.) wavelength and emission (em.) range were set as follows: Venus, ex. 514, em. 545–540; GFP, ex. 488, em. 500–550; tdTomato and Cy3, ex. 561, em. 570–625; Cy5, ex. 633, em. 650–750 nm. Images were acquired in sequential mode with a low-noise Hybrid detector combining avalanche gain and GaAsP detection (HyD, Leica Wetzlar, Germany). Metrology measurements were performed on fluorescent beads to confirm proper alignment of lasers and field homogeneity, using the ImageJ-based MetroloJ plugin. Deconvolution with the experimental point spread of function from fluorescent beads using the Maximum Likelihood Estimation algorithm was performed with Huygens software (Scientific Volume Imaging).

Image analysis

Segmentation varicosities, distance measurements and co-localization analysis were performed in three dimensions (3D) using the procedures implemented in the ImageJ-based DiAna plugin, extensively described in Gilles (Gilles et al., 2017). The 3D intensity radial distribution centered on the local maximum of each object was used to segment the objects. GFP in thalamic afferents was localized in the axons as well as in boutons. Thus, axonal boutons were selected according to their co-localization with the immunolabeled presynaptic active zone protein Bassoon. Following segmentation, 3D distances between the borders of VGLUT objects and the closest 5-HT4R were computed and co-localizing objects counted. For co-localization analysis of 5-HT4R on spine heads and shafts, the dendritic shaft was first reconstructed with Neuronstudio software, then imported into ImageJ while spine heads were segmented in ImageJ (Dos Santos et al., 2017; Heck et al., 2015). 5-HT4R co-localization analysis was then performed with shafts and spine heads.

Immuno-EM analysis

Pre-embedding immuno-gold analyses were performed as described (Hebert-Chatelain et al., 2014). In short, after a pre-incubation step in blocking solution [(10% BSA, 0.1% sodium azide and 0.02% saponin in Tris-buffered saline (TBS)], coronal sections were incubated with rat monoclonal anti-mCherry (1:50, Sigma Aldrich Italia, Milano, Italy) and rabbit polyclonal anti-5-HT (1:200, ThermoFisher Scientific, Waltham, Massachusetts, USA) primary antibodies in blocking solution with 0.004% saponin. After several washes (1% BSA in TBS), sections were incubated with a goat anti-rabbit biotinylated secondary antibody (1:150, Vector Laboratories, Peterborough, United Kingdom) and with secondary 1.4 nm gold-labeled goat anti-rat Immunoglobulin-G (Fab' fragment, 1:100, Nanoprobes Inc., Yaphank, New York, USA). Sections were then processed in avidin-biotin complex (1:50, Vector Laboratories, Peterborough, United Kingdom), silver-intensified with the HQ Silver kit (Nanoprobes Inc., Yaphank, New York, USA), and incubated in a solution of 0.05% DAB and 0.01% hydrogen peroxide (Vector Laboratories, Peterborough, United Kingdom). After post-fixation with 1% glutaraldehyde, the immunolabeled sections were analyzed by electron microscopy, as described (Giacomini et al., 2016). Scanning transmission electron microscopy (STEM) images were collected with a Schottky field-emission gun FEI Tecnai G2 F20 (FEI, Hillsboro, Oregon) transmission electron microscope equipped with a high angular annular dark field (HAADF) detector and operating at an acceleration voltage of 200 kV.

Activation of CaMKII in acute slices

Acute slices from *drd1a-tdTomato^{tg}* mice ($n = 3$) were prepared as described (Trusel et al., 2015). After 1 h of recovery in aCSF at 32°C, slices were incubated 10 min at 32°C with or without 5 μ M of the 5-HT₄R agonist RS67333; after 5 min washout at 32°C in aCSF, slices were fixed in 4% PFA + 50 mM NaF for 4 h at 4°C. Slices were then rinsed three times (5 min each) in PBS and stored at -20°C in a cryoprotective solution (PBS + 30% glycerol + 30% PEG) until immunofluorescence was performed. After several washes in PBS, slices were permeabilized in PBS + 0.5% Triton X-100 for 4 h at room temperature, blocked with PBS + 0.5% Triton X-100 + 3% BSA for 1 h at room temperature and incubated with the primary antibody [rabbit anti-pCaMKII (Thr286) 1:400, Cell Signaling] in PBS + 0.1% Triton X-100 + 3% BSA overnight at 4°C. After several washes in PBS, slices were incubated with the secondary antibody (Alexa 488 anti-rabbit, 1:500, ThermoFisher Scientific) in PBS + 0.1% Triton X-100 + 3% BSA for 2 h at room temperature. After several washes in PBS, slices were mounted in ProLong Gold. Images were acquired with an inverted confocal microscope (Leica TCS SP5 AOBs TANDEM) with a 20X objective, acquiring 10 μ m thick stacks with a 2 μ m z-step, starting from 30 μ m below the surface of the slice. For each experimental condition, 3 slices were acquired, and 2 images per slice were analyzed, counting the total number of pCaMKII and tdTomato double-positive cells.

QUANTIFICATION AND STATISTICAL ANALYSIS

Appropriate parametric statistic were utilized to test our hypothesis, unless data did not meet the assumptions of the intended parametric test (normality test). In this case, appropriate non-parametric test were used. Power analysis assumptions were: power = 0.9; alpha = 0.5; two-tailed and expected difference 50% greater than the observed standard deviation. Data were analyzed by two-way repeated-measure (RM2WA) or one-way repeated-measures ANOVA (RM1WA) for comparisons within a group, and one-way ANOVA (1WA) for between-group comparisons (GraphPad Prism 6 software). Post hoc analysis (Tukey; Home-Sidak, as indicated) was performed only when ANOVA yielded a significant main effect. Two groups were tested for statistical significance using the two-population t test and Mann-Whitney U-nonparametric test (GraphPad Prism 6 software). Statistical details of experiments are shown in results, figures and figure legends. Data are reported as mean \pm s.e.m., unless stated otherwise. In the electrophysiological experiments, measurement of EPSPs were made using pClamp software (Molecular Devices). The occurrence and magnitude of synaptic plasticity was evaluated by comparing the normalized EPSP amplitudes from the last 5 min of baseline recordings with the corresponding values at 25-35 min after STDP (as indicated in Figure 1B). LTD plots were generated by averaging the peak amplitude of individual EPSPs in 1 min bins. Coefficients of variation (CV) for EPSPs were calculated by the ratio of the standard deviation (sd) and the mean EPSP amplitude (Shen et al., 2008). In imaging experiments, the magnitude of the Ca²⁺ signal was measured as the area under the curve (AUC) in the transient region. n values for electrophysiological and imaging experiments represent the number of recorded cells. No statistical methods were used to pre-determine sample sizes, which were, however, similar to those reported previously (Nazzaro et al., 2012; Trusel et al., 2015). For immunohistochemistry, images were processed with Adobe Photoshop (Adobe Systems) using only linear adjustments of contrast and color.

DATA AND SOFTWARE AVAILABILITY

Mouse strains: *Pet1_{2/10}-Cre*; MGI:5620670 [<http://www.informatics.jax.org/allele/key/854058>]; mouse *Tph2GFP^{+/-}*; MGI:5442736 [<http://www.informatics.jax.org/allele/key/821477>]; a custom macro in ImageJ using algorithms from the Plugin 3DImageSuite (http://imagejdocu.tudor.lu/doku.php?id=plugin:stacks:3d_ij_suite:start) is available upon request (nicolas.heck@upmc.fr) since it requires to be adapted according to image names and calibration.