

Logics of left variable inclusion and Płonka sums of matrices

S. Bonzio, T. Moraschini & M. Pra Baldi

Archive for Mathematical Logic

ISSN 0933-5846

Volume 60

Combined 1-2

Arch. Math. Logic (2021) 60:49-76

DOI 10.1007/s00153-020-00727-6

Your article is protected by copyright and all rights are held exclusively by Springer-Verlag GmbH Germany, part of Springer Nature. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".



Logics of left variable inclusion and Płonka sums of matrices

S. Bonzio¹ · T. Moraschini² · M. Pra Baldi³Received: 18 December 2018 / Accepted: 18 March 2020 / Published online: 13 April 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Abstract

The paper aims at studying, in full generality, logics defined by imposing a variable inclusion condition on a given logic \vdash . We prove that the description of the algebraic counterpart of the left variable inclusion companion of a given logic \vdash is related to the construction of Płonka sums of the matrix models of \vdash . This observation allows to obtain a Hilbert-style axiomatization of the logics of left variable inclusion, to describe the structure of their reduced models, and to locate them in the Leibniz hierarchy.

Keywords Płonka sums · Kleene logics · Abstract algebraic logic · Regular varieties

Mathematics Subject Classification Primary 03G27; Secondary 03G25

1 Introduction

It is always possible to associate with an arbitrary propositional logic \vdash , two new substitution-invariant consequence relations \vdash^l and \vdash^r , which satisfy respectively a *left* and a *right variable inclusion principle*, as follows:

$$\Gamma \vdash^l \varphi \iff \text{there is } \Delta \subseteq \Gamma \text{ s.t. } \text{Var}(\Delta) \subseteq \text{Var}(\varphi) \text{ and } \Delta \vdash \varphi,$$

✉ S. Bonzio
stefano.bonzio@gmail.com

T. Moraschini
tommaso.moraschini@ub.edu

M. Pra Baldi
m.prabaldi@gmail.com

¹ Department of Biomedical Sciences and Public Health, Polytechnic University of the Marche, Ancona, Italy

² Department of Philosophy, University of Barcelona, Carrer Montalegre 6, 08001 Barcelona, Spain

³ Department of Pedagogy, Psychology and Philosophy, University of Cagliari, Cagliari, Italy

and

$$\Gamma \vdash^r \varphi \iff \begin{cases} \Gamma \vdash \varphi \text{ and } \text{Var}(\varphi) \subseteq \text{Var}(\Gamma), \text{ or} \\ \Sigma \subseteq \Gamma, \text{ with } \Sigma \text{ a set of inconsistency terms for } \vdash . \end{cases}$$

Accordingly, we say that the logics \vdash^l and \vdash^r are, respectively, the *left* and the *right variable inclusion companions* of \vdash , sometimes also referred to as *containment logics*.

Prototypical examples of variable inclusion companions are found in the realm of three-valued logics. For instance, the left and the right variable inclusion companions of classical (propositional) logic are respectively *paraconsistent weak Kleene logic* (PWK for short) [33,40], and *Bochvar logic* [7]. The fact that these logics coincide with the variable inclusion companions of classical logic was shown in [20,62]. Remarkably, both PWK and Bochvar logic feature the presence of a non-sensical, infectious truth value [21,61], which made them a valuable tool in modeling reasonings with non-existing objects [55], computer-programs affected by errors [27] as well as recent developments in the theory of truth [23] and philosophy of logic [8].

Recent work [10] linked PWK to the algebraic theory of regular varieties, i.e. equational classes axiomatized by equations $\varphi \approx \psi$ such that $\text{Var}(\varphi) = \text{Var}(\psi)$. The representation theory of regular varieties is largely due to the pioneering work of Płonka [51], and is tightly related to a special class-operator $\mathcal{P}_l(\cdot)$ nowadays called *Płonka sums*. Over the years, regular varieties have been studied in depth both from a purely algebraic perspective [34,35,39,52] and in connection to their topological duals [9,11,32,46,60]. The machinery of Płonka sums has also found useful applications in the study of the constraint satisfaction problem [2] and database semantics [47,56] and in the application of algebraic methods in computer science [13].

One of the main results of [10] states that the algebraic counterpart of PWK is the class of Płonka sums of Boolean algebras. This observation led us to investigate the relations between left variable inclusion companions and Płonka sums in full generality.¹ Our study is carried on in the conceptual framework of abstract algebraic logic [22,28,29].

We begin by generalizing the construction of Płonka sums from algebras to logical matrices (Sect. 3). This allows us to condense the connection between left variable inclusion principles and Płonka sums in the following slogan: The left variable inclusion companion \vdash^l of a logic \vdash is complete w.r.t. the class of Płonka sums of matrix models of \vdash (Corollary 15).

As a matter of fact, left variable inclusion companions \vdash^l are especially well-behaved in case the original logic \vdash has a *partition function* [59], a feature shared by the vast majority of non-pathological logics in the literature. The importance of partition functions is reflected both at a syntactic and at a semantic level. Accordingly, on the one hand we present a general method to transform every Hilbert-style calculus for a finitary logic \vdash with a partition function into a Hilbert-style calculus for \vdash^l (Theorem 24). On the other hand, partition functions can be exploited to tame the structure of the matrix semantics $\text{Mod}^{\text{Su}}(\vdash^l)$ of \vdash^l , given by the so-called Suszko reduced models of \vdash^l . In particular, we obtain a full description of $\text{Mod}^{\text{Su}}(\vdash^l)$ in case

¹ A similar investigation of right variable inclusion companions is developed in [12].

\vdash is a finitary equivalential logic with a partition function (Theorems 31, 38). We close our investigation by determining the location of \vdash^l in the Leibniz hierarchy (Section 6).

2 Preliminaries

Abstract algebraic logic

For standard background on universal algebras and abstract algebraic logic we refer the reader respectively to [1, 4–6, 14, 22, 24, 28–30, 48, 63]. In this paper, algebraic languages are assumed not to contain constant symbols. Moreover, unless stated otherwise, we work within a fixed but arbitrary algebraic language. We denote algebras by $\mathbf{A}, \mathbf{B}, \mathbf{C} \dots$ respectively with universes $A, B, C \dots$. A class of algebras is a *variety* if is axiomatized by equations. Given a class of algebras \mathbf{K} , we denote by $\mathbb{V}(\mathbf{K})$ the variety generated by \mathbf{K} . Let \mathbf{Fm} be the algebra of formulas built up over a countably infinite set Var of variables. Given a formula $\varphi \in Fm$, we denote by $Var(\varphi)$ the set of variables really occurring in φ . Similarly, given $\Gamma \subseteq Fm$, we set

$$Var(\Gamma) = \bigcup \{Var(\gamma) : \gamma \in \Gamma\}.$$

A *logic* is a substitution invariant consequence relation $\vdash \subseteq \mathcal{P}(Fm) \times Fm$, namely for every substitution $\sigma : \mathbf{Fm} \rightarrow \mathbf{Fm}$,

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi).$$

Given $\varphi, \psi \in Fm$, we write $\varphi \dashv\vdash \psi$ as a shorthand for $\varphi \vdash \psi$ and $\psi \vdash \varphi$. Moreover, we denote by $Cn_{\vdash} : \mathcal{P}(Fm) \rightarrow \mathcal{P}(Fm)$ the closure operator associated with \vdash . A logic \vdash is *finitary* when the following holds for all $\Gamma \cup \{\varphi\} \subseteq Fm$:

$$\Gamma \vdash \varphi \iff \exists \Delta \subseteq \Gamma \text{ s.t. } \Delta \text{ is finite and } \Delta \vdash \varphi.$$

A *matrix* is a pair $\langle \mathbf{A}, F \rangle$ where \mathbf{A} is an algebra and $F \subseteq A$. In this case, \mathbf{A} is called the *algebraic reduct* of the matrix $\langle \mathbf{A}, F \rangle$. We denote by $\mathbb{I}, \mathbb{S}, \mathbb{P}$ and \mathbb{P}_{SD} respectively the class operators of isomorphic copies, substructures, direct products and subdirect products, which apply both to classes of algebras and classes of matrices.

Every class of matrices \mathbf{M} induces a logic as follows:

$$\Gamma \vdash_{\mathbf{M}} \varphi \iff \text{for every } \langle \mathbf{A}, F \rangle \in \mathbf{M} \text{ and homomorphism } h : \mathbf{Fm} \rightarrow \mathbf{A}, \\ \text{if } h[\Gamma] \subseteq F, \text{ then } h(\varphi) \in F.$$

A logic \vdash is *complete* w.r.t. a class of matrices \mathbf{M} when it coincides with $\vdash_{\mathbf{M}}$.

A matrix $\langle \mathbf{A}, F \rangle$ is a *model* of a logic \vdash when

$$\begin{aligned} &\text{if } \Gamma \vdash \varphi, \text{ then for every homomorphism } h: \mathbf{Fm} \rightarrow \mathbf{A}, \\ &\text{if } h[\Gamma] \subseteq F, \text{ then } h(\varphi) \in F. \end{aligned}$$

A set $F \subseteq A$ is a (deductive) *filter* of \vdash on \mathbf{A} , or simply a \vdash -*filter*, when the matrix $\langle \mathbf{A}, F \rangle$ is a model of \vdash . We denote by $\mathcal{F}_{i\vdash}\mathbf{A}$ the set of all filters of \vdash on \mathbf{A} , which turns out to be a closure system. Moreover, we denote by $\text{Fg}_{\vdash}^{\mathbf{A}}(\cdot)$ the closure operator of \vdash -filter generation on \mathbf{A} .

Let \mathbf{A} be an algebra and $F \subseteq A$. A congruence θ of \mathbf{A} is *compatible* with F when for every $a, b \in A$,

$$\text{if } a \in F \text{ and } \langle a, b \rangle \in \theta, \text{ then } b \in F.$$

It turns out that there exists the largest congruence of \mathbf{A} which is compatible with F . This congruence is called the *Leibniz congruence* of F on \mathbf{A} , and it is denoted by $\Omega^{\mathbf{A}}F$.

Let \mathbf{A} be an algebra, $F \subseteq A$ and \vdash be a logic. The *Suszko congruence* of F on \mathbf{A} , is defined as

$$\tilde{\Omega}_{\vdash}^{\mathbf{A}}F := \bigcap \{ \Omega^{\mathbf{A}}G : F \subseteq G \text{ and } G \in \mathcal{F}_{i\vdash}\mathbf{A} \}.$$

Let \mathbf{A} be an algebra. A function $p: A^n \rightarrow A$ is a *polynomial function* of \mathbf{A} if there are a natural number m , a formula $\varphi(x_1, \dots, x_{n+m})$, and elements $b_1, \dots, b_m \in A$ such that

$$p(a_1, \dots, a_n) = \varphi^{\mathbf{A}}(a_1, \dots, a_n, b_1, \dots, b_m)$$

for every $a_1, \dots, a_n \in A$.

Lemma 1 [28, Theorem 4.23] *Let \mathbf{A} be an algebra, $F \subseteq A$ and $a, b \in A$.*

$$\langle a, b \rangle \in \Omega^{\mathbf{A}}F \iff \text{for every unary pol. function } p: A \rightarrow A, \\ p(a) \in F \text{ if and only if } p(b) \in F.$$

Lemma 2 [28, Theorem 5.32] *Let \vdash be a logic, \mathbf{A} be an algebra, $F \subseteq A$ and $a, b \in A$.*

$$\langle a, b \rangle \in \tilde{\Omega}_{\vdash}^{\mathbf{A}}F \iff \text{for every unary pol. function } p: A \rightarrow A, \\ \text{Fg}_{\vdash}^{\mathbf{A}}(F \cup \{p(a)\}) = \text{Fg}_{\vdash}^{\mathbf{A}}(F \cup \{p(b)\}).$$

The Leibniz and Suszko congruences allow to associate two distinguished classes of models to logics. More precisely, given a logic \vdash , we set

$$\begin{aligned} \text{Mod}(\vdash) &:= \{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash\}; \\ \text{Mod}^*(\vdash) &:= \{\langle \mathbf{A}, F \rangle \in \text{Mod}(\vdash) : \Omega^{\mathbf{A}}F \text{ is the identity}\}; \\ \text{Mod}^{\text{Su}}(\vdash) &:= \{\langle \mathbf{A}, F \rangle \in \text{Mod}(\vdash) : \tilde{\Omega}_{\vdash}^{\mathbf{A}}F \text{ is the identity}\}. \end{aligned}$$

The above classes of matrices are called, respectively, the classes of models, *Leibniz reduced models*, and *Suszko reduced models* of \vdash . It turns out that $\text{Mod}^{\text{Su}}(\vdash) = \mathbb{P}_{\text{SD}} \text{Mod}^*(\vdash)$.

Trivial matrices will play a useful role in the whole paper. More precisely, a matrix $\langle \mathbf{A}, F \rangle$ is *trivial* if $F = A$. We denote by $\langle \mathbf{1}, \{1\} \rangle$ the trivial matrix, where $\mathbf{1}$ is the trivial algebra. Observe that the latter matrix is a model (resp. Leibniz and Suszko reduced model) of every logic. Moreover, if \vdash is a logic and $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\text{Su}}(\vdash)$ is a trivial matrix, then $\langle \mathbf{A}, F \rangle = \langle \mathbf{1}, \{1\} \rangle$.

Given a logic \vdash , we set

$$\text{Alg}(\vdash) = \{\mathbf{A} : \text{there is } F \subseteq A \text{ s.t. } \langle \mathbf{A}, F \rangle \in \text{Mod}^{\text{Su}}(\vdash)\}.$$

In other words, $\text{Alg}(\vdash)$ is the class of algebraic reducts of matrices in $\text{Mod}^{\text{Su}}(\vdash)$. The class $\text{Alg}(\vdash)$ is called the *algebraic counterpart* of \vdash . For the vast majority of logics \vdash , the class $\text{Alg}(\vdash)$ is the class of algebras intuitively associated with \vdash .

Lemma 3 [28, Lemma 5.78] *Let \vdash be a logic defined by a class of matrices M . Then $\text{Alg}(\vdash) \subseteq \mathbb{V}(K)$, where K is the class of algebraic reducts of M .*

Lemma 4 *Let \vdash be a logic and $\epsilon, \delta \in \text{Fm}$. The following are equivalent:*

- (i) $\text{Alg}(\vdash) \models \epsilon \approx \delta$;
- (ii) $\varphi(\epsilon, \vec{z}) \dashv\vdash \varphi(\delta, \vec{z})$, for every formula $\varphi(v, \vec{z})$.

Proof See [28, Lemma 5.74(1)] and [28, Theorem 5.76]. □

Now, we turn our attention to a fundamental topic in abstract algebraic logic, that is the so-called *Leibniz hierarchy*, see for example [28,36–38,57]. We review only the material which is necessary for the present purpose. A logic \vdash is *protoalgebraic* if there is a set of formulas $\Delta(x, y)$ such that

$$\emptyset \vdash \Delta(x, x) \text{ and } x, \Delta(x, y) \vdash y.$$

Remarkably, \vdash is protoalgebraic if and only if $\text{Mod}^*(\vdash) = \text{Mod}^{\text{Su}}(\vdash)$.

A logic \vdash is *equivalential* if there is a set of formulas $\Delta(x, y)$ such that for every $\langle \mathbf{A}, F \rangle \in \text{Mod}(\vdash)$,

$$\langle a, b \rangle \in \Omega^{\mathbf{A}}F \iff \Delta^{\mathbf{A}}(a, b) \subseteq F, \text{ for all } a, b \in A.$$

In this case, $\Delta(x, y)$ is said to be a set of *congruence formulas* for \vdash . Remarkably, \vdash is equivalential if and only if $\text{Mod}^*(\vdash)$ is closed under \mathbb{S} and \mathbb{P} . Consequently, every equivalential logic is protoalgebraic.

A logic \vdash is *truth-equational* if there is a set of equations $\tau(x)$ such that for all $\langle \mathbf{A}, F \rangle \in \text{Mod}^*(\vdash)$,

$$a \in F \iff \mathbf{A} \models \tau(a), \text{ for all } a \in A.$$

In this case, $\tau(x)$ is said to be a set of *defining equations* for \vdash .

Finally, a logic \vdash is *algebraizable* when it is both equivalential and truth-equational. In this case, $\text{Alg}(\vdash)$ is called the *equivalent algebraic semantics* of \vdash .

Ponka sums

For standard information on Płonka sums we refer the reader to [51–53,59]. A *semilattice* is an algebra $\mathbf{A} = \langle A, \vee \rangle$, where \vee is a binary commutative, associative and idempotent operation. Given a semilattice \mathbf{A} and $a, b \in A$, we set

$$a \leq b \iff a \vee b = b.$$

It is easy to see that \leq is a partial order on A .

Definition 5 *A directed system of algebras consists of:*

- (i) *a semilattice $I = \langle I, \vee \rangle$;*
- (ii) *a family of algebras $\{\mathbf{A}_i : i \in I\}$ with disjoint universes;*
- (iii) *a homomorphism $f_{ij} : \mathbf{A}_i \rightarrow \mathbf{A}_j$, for every $i, j \in I$ such that $i \leq j$;*

moreover, f_{ii} is the identity map for every $i \in I$, and if $i \leq j \leq k$, then $f_{ik} = f_{jk} \circ f_{ij}$.

Let X be a directed system of algebras as above. The *Płonka sum* of X , in symbols $\mathcal{P}_t(X)$ or $\mathcal{P}_t(\mathbf{A}_i)_{i \in I}$, is the algebra defined as follows. The universe of $\mathcal{P}_t(\mathbf{A}_i)_{i \in I}$ is the union $\bigcup_{i \in I} A_i$. Moreover, for every n -ary basic operation f and $a_1, \dots, a_n \in \bigcup_{i \in I} A_i$, we set

$$f^{\mathcal{P}_t(\mathbf{A}_i)_{i \in I}}(a_1, \dots, a_n) := f^{\mathbf{A}_j}(f_{i_1 j}(a_1), \dots, f_{i_n j}(a_n))$$

where $a_1 \in A_{i_1}, \dots, a_n \in A_{i_n}$ and $j = i_1 \vee \dots \vee i_n$.

Observe that if in the above display we replace f by any complex formula φ in n -variables, we still have that

$$\varphi^{\mathcal{P}_t(\mathbf{A}_i)_{i \in I}}(a_1, \dots, a_n) = \varphi^{\mathbf{A}_j}(f_{i_1 j}(a_1), \dots, f_{i_n j}(a_n)).$$

Notation: Given a formula φ , we will often write $\varphi^{\mathcal{P}_t}$ instead of $\varphi^{\mathcal{P}_t(\mathbf{A}_i)_{i \in I}}$ when no confusion shall occur.

The theory of Płonka sums is strictly related with a special kind of operation:

Definition 6 Let \mathbf{A} be an algebra of type ν . A function $\cdot : A^2 \rightarrow A$ is a partition function in \mathbf{A} if the following conditions are satisfied for all $a, b, c \in A$, $a_1, \dots, a_n \in A^n$ and for any operation $g \in \nu$ of arity $n \geq 1$.

- P₁.** $a \cdot a = a$
- P₂.** $a \cdot (b \cdot c) = (a \cdot b) \cdot c$
- P₃.** $a \cdot (b \cdot c) = a \cdot (c \cdot b)$
- P₄.** $g(a_1, \dots, a_n) \cdot b = g(a_1 \cdot b, \dots, a_n \cdot b)$
- P₅.** $b \cdot g(a_1, \dots, a_n) = b \cdot a_1 \dots \cdot a_n$

The next result makes explicit the relation between Plonka sums and partition functions:

Theorem 7 [51, Theorem II] Let \mathbf{A} be an algebra of type ν with a partition function \cdot . The following conditions hold:

1. A can be partitioned into $\{A_i : i \in I\}$ where any two elements $a, b \in A$ belong to the same component A_i exactly when

$$a = a \cdot b \text{ and } b = b \cdot a.$$

Moreover, every A_i is the universe of a subalgebra \mathbf{A}_i of \mathbf{A} .

2. The relation \leq on I given by the rule

$$i \leq j \iff \text{there exist } a \in A_i, b \in A_j \text{ s.t. } b \cdot a = b$$

is a partial order and $\langle I, \leq \rangle$ is a semilattice.

3. For all $i, j \in I$ such that $i \leq j$ and $b \in A_j$, the map $f_{ij} : \mathbf{A}_i \rightarrow \mathbf{A}_j$, defined by the rule $f_{ij}(x) = x \cdot b$ is a homomorphism. The definition of f_{ij} is independent from the choice of b , since $a \cdot b = a \cdot c$, for all $a \in A_i$ and $c \in A_j$.
4. $Y = \langle \langle I, \leq \rangle, \{\mathbf{A}_i\}_{i \in I}, \{f_{ij} : i \leq j\} \rangle$ is a directed system of algebras such that $\mathcal{P}_I(Y) = \mathbf{A}$.

It is worth remarking that the construction of Plonka sums preserves the validity of the so-called *regular identities*, i.e. identities of the form $\varphi \approx \psi$ such that $\text{Var}(\varphi) = \text{Var}(\psi)$ (for details, see [51,53]).

3 The left variable inclusion companion of a logic

The definition of *directed system* can be extended, as follows, to logical matrices:

Definition 8 A directed system of matrices consists of:

- (i) a semilattice $I = \langle I, \vee \rangle$;
- (ii) a family of matrices $\{\langle \mathbf{A}_i, F_i \rangle\}_{i \in I}$ with disjoint universes;
- (iii) a homomorphism $f_{ij} : \mathbf{A}_i \rightarrow \mathbf{A}_j$ such that $f_{ij}[F_i] \subseteq F_j$, for every $i, j \in I$ such that $i \leq j$;

moreover, f_{ii} is the identity map for every $i \in I$, and if $i \leq j \leq k$, then $f_{ik} = f_{jk} \circ f_{ij}$.

Given directed system of matrices X as above, we set

$$\mathcal{P}_I(X) := \langle \mathcal{P}_I(\mathbf{A}_i)_{i \in I}, \bigcup_{i \in I} F_i \rangle.$$

The matrix $\mathcal{P}_I(X)$ is the *Plonka sum* of the directed system of matrices X . Given a class \mathbf{M} of matrices, we denote by $\mathcal{P}_I(\mathbf{M})$ the class of all Plonka sums of directed systems of matrices in \mathbf{M} . The following observation is a routine computation:

Lemma 9 $\mathcal{SP}_I(\mathbf{M}) \subseteq \mathcal{P}_I(\mathcal{S}(\mathbf{M}))$ and $\mathbb{P}\mathcal{P}_I(\mathbf{M}) \subseteq \mathcal{P}_I(\mathbb{P}(\mathbf{M}))$, for every class of matrices \mathbf{M} .

Definition 10 Let \vdash be a logic. The left variable inclusion companion of \vdash is the relation $\vdash^l \subseteq \mathcal{P}(Fm) \times Fm$ defined for every $\Gamma \cup \{\varphi\} \subseteq Fm$ as

$$\Gamma \vdash^l \varphi \iff \text{there is } \Gamma' \subseteq \Gamma \text{ s.t. } \text{Var}(\Gamma') \subseteq \text{Var}(\varphi) \text{ and } \Gamma' \vdash \varphi.$$

It is immediate to check that \vdash^l is indeed a logic and that $\vdash^l \subseteq \vdash$. We will often refer to the left variable inclusion of a logic simply as its variable inclusion companion.

Example 11 Let \vdash be propositional classical logic. Then \vdash^l is the logic known as Paraconsistent Weak Kleene, PWK for short, originally introduced in [40]. This logic is equivalently defined, *syntactically*, by imposing the variable inclusion constrain, as in Definition 10, to classical logic or, *semantically* via the so-called *weak Kleene tables* with two of the three truth values as designated (see [10,20]).

Example 12 The left variable inclusion companions of Strong Kleene logic and of the logic of Paradox (introduced in [54]) have been introduced and discussed in [61]. They are semantically defined, by adding a nonsensical truth value to the (single) matrix inducing Strong Kleene and the logic of Paradox, respectively.

In [10], it is shown that an algebraic semantics for PWK is obtained via Plonka sums of Boolean algebras. We shall show that this idea can be generalized to the variable inclusion companion of any logic \vdash .

Lemma 13 Let \vdash be a logic and X be a directed system of models of \vdash . Then $\mathcal{P}_I(X)$ is a model of \vdash^l .

Proof Suppose that $\Gamma \vdash^l \varphi$ and consider a homomorphism $v: \mathbf{Fm} \rightarrow \mathcal{P}_I(\mathbf{A}_i)_{i \in I}$ such that $v[\Gamma] \subseteq \bigcup_{i \in I} F_i$. By the definition of \vdash^l , there exists $\Delta \subseteq \Gamma$ such that $\text{Var}(\Delta) \subseteq \text{Var}(\varphi)$ and $\Delta \vdash \varphi$. Consider an enumeration $\text{Var}(\varphi) = \{x_1, \dots, x_n\}$. There are $i_1, \dots, i_n \in I$ such that $v(x_1) \in A_{i_1}, \dots, v(x_n) \in A_{i_n}$. We set $j := i_1 \vee \dots \vee i_n$.

Now, consider a homomorphism $g: \mathbf{Fm} \rightarrow \mathbf{A}_j$ such that

$$g(x_m) = f_{imj}(v(x_m)), \quad \text{for every } m \leq n.$$

We claim that $g[\Delta] \subseteq F_j$. To prove this, consider an arbitrary formula $\delta \in \Delta$. Since $Var(\Delta) \subseteq \{x_1, \dots, x_n\}$, we can assume that $Var(\delta) = \{x_{m_1}, \dots, x_{m_k}\} \subseteq \{x_1, \dots, x_n\}$ for some $k \leq n$. Set $l := i_{m_1} \vee \dots \vee i_{m_k}$. From the definition of $\mathcal{P}_l(X)$ we have that

$$v(\delta) = \delta^{\mathcal{P}_l}(v(x_{m_1}), \dots, v(x_{m_k})) = \delta^{\mathbf{A}^l}(f_{i_{m_1}l}(v(x_{m_1})), \dots, f_{i_{m_k}l}(v(x_{m_k}))).$$

Since $v(\delta) \in \bigcup_{i \in I} F_i$, this implies that

$$\delta^{\mathbf{A}^l}(f_{i_{m_1}l}(v(x_{m_1})), \dots, f_{i_{m_k}l}(v(x_{m_k}))) \in F_l. \tag{1}$$

Now observe that $l \leq j$. Therefore there is a homomorphism $f_{lj} : \mathbf{A}^l \rightarrow \mathbf{A}^j$ such that $f_{lj}[F_l] \subseteq F_j$. Together with (1), this implies that

$$\begin{aligned} g(\delta) &= \delta^{\mathbf{A}^j}(f_{i_{m_1}j}(v(x_{m_1})), \dots, f_{i_{m_k}j}(v(x_{m_k}))) \\ &= \delta^{\mathbf{A}^j}(f_{lj} \circ f_{i_{m_1}l}(v(x_{m_1})), \dots, f_{lj} \circ f_{i_{m_k}l}(v(x_{m_k}))) \\ &= f_{lj} \delta^{\mathbf{A}^l}(f_{i_{m_1}l}(v(x_{m_1})), \dots, f_{i_{m_k}l}(v(x_{m_k}))) \\ &\in f_{lj}[F_l] \subseteq F_j. \end{aligned}$$

This establishes our claim.

Recall that $\Delta \vdash \varphi$. Since $\langle \mathbf{A}_j, F_j \rangle$ is a model of \vdash and by the claim $g[\Delta] \subseteq F_j$, we conclude that $g(\varphi) \in F_j$. But this means that

$$\begin{aligned} v(\varphi) &= \varphi^{\mathcal{P}_l}(v(x_1), \dots, v(x_n)) \\ &= \varphi^{\mathbf{A}^j}(f_{i_1j}(v(x_1)), \dots, f_{i_nj}(v(x_n))) \\ &= g(\varphi) \in F_j \subseteq \bigcup_{i \in I} F_i. \end{aligned}$$

Hence we conclude that $\mathcal{P}_l(X)$ is a model of \vdash^l as desired. □

Recall that $\mathbf{1}$ is the trivial algebra. The following construction originates in [44]. Given an algebra \mathbf{A} , there is always a directed system of algebras given by \mathbf{A} and $\mathbf{1}$ equipped with the identity endomorphisms and the unique homomorphism $f : \mathbf{A} \rightarrow \mathbf{1}$. We denote by $\mathbf{A} \oplus \mathbf{1}$ the Płonka sum of this directed system. Observe that $\mathbf{A} \oplus \mathbf{1}$ is the algebra with universe $A \cup \{1\}$ and basic operations f defined as follows:

$$f^{\mathbf{A} \oplus \mathbf{1}}(a_1, \dots, a_n) := \begin{cases} f^{\mathbf{A}}(a_1, \dots, a_n) & \text{if } a_1, \dots, a_n \in A \\ 1 & \text{otherwise.} \end{cases}$$

Observe that the above construction can be lifted to matrices. More precisely, given an arbitrary matrix $\langle \mathbf{A}, F \rangle$, there is always a directed system of matrices given by $\langle \mathbf{A}, F \rangle$ and $\langle \mathbf{1}, \{1\} \rangle$ equipped with the identity endomorphisms and the unique homomorphism $f : \mathbf{A} \rightarrow \mathbf{1}$. The Płonka sum of this system is the matrix $\langle \mathbf{A} \oplus \mathbf{1}, F \cup \{1\} \rangle$.

Theorem 14 *Let \vdash be a logic and M be a class of matrices containing $\langle \mathbf{1}, \{1\} \rangle$. If \vdash is complete w.r.t. M , then \vdash^l is complete w.r.t. $\mathcal{P}_l(M)$.*

Proof In the light of Lemma 13 it will be enough to show that if $\Gamma \not\vdash^l \varphi$, then $\Gamma \not\vdash_{\mathcal{P}_l(M)} \varphi$. To this end, suppose that $\Gamma \not\vdash^l \varphi$. Define

$$\begin{aligned} \Gamma^+ &:= \{\gamma \in \Gamma : \text{Var}(\gamma) \subseteq \text{Var}(\varphi)\} \\ \Gamma^- &:= \{\gamma \in \Gamma : \text{Var}(\gamma) \not\subseteq \text{Var}(\varphi)\}. \end{aligned}$$

Clearly $\Gamma = \Gamma^+ \cup \Gamma^-$. Since $\Gamma \not\vdash^l \varphi$, we know that $\Gamma^+ \not\vdash \varphi$. Together with the fact that \vdash is complete w.r.t. M , this implies that there exists a matrix $\langle \mathbf{A}, F \rangle \in M$ and a homomorphism $v: \mathbf{Fm} \rightarrow \mathbf{A}$ such that $v[\Gamma^+] \subseteq F$ and $v(\varphi) \notin F$.

Since $\langle \mathbf{A}, F \rangle, \langle \mathbf{1}, \{1\} \rangle \in M$, we have that $\langle \mathbf{A} \oplus \mathbf{1}, F \cup \{1\} \rangle \in \mathcal{P}_l(M)$. Now, consider the homomorphism $g: \mathbf{Fm} \rightarrow \mathbf{A} \oplus \mathbf{1}$ defined for every variable $x \in \text{Var}$ as follows:

$$g(x) := \begin{cases} v(x) & \text{if } x \in \text{Var}(\varphi) \\ 1 & \text{otherwise.} \end{cases}$$

From the definition of $\mathbf{A} \oplus \mathbf{1}$ it follows that

$$\begin{aligned} g[\Gamma^-] &\subseteq \{1\} \subseteq F \cup \{1\} \\ g(\gamma) &= v(\gamma) \text{ for every } \gamma \in \Gamma^+ \cup \{\varphi\}. \end{aligned}$$

Together with the fact that $v[\Gamma^+] \subseteq F$ and $v(\varphi) \notin F$, this implies that

$$g[\Gamma] = g[\Gamma^+ \cup \Gamma^-] \subseteq F \cup \{1\} \text{ and } g(\varphi) \notin F \cup \{1\}.$$

Hence we conclude that $\Gamma \not\vdash_{\mathcal{P}_l(M)} \varphi$ as desired. □

Corollary 15 *Let \vdash be a logic. The variable inclusion companion \vdash^l is complete w.r.t. any of the following classes of matrices:*

$$\mathcal{P}_l(\text{Mod}(\vdash)), \quad \mathcal{P}_l(\text{Mod}^*(\vdash)), \quad \mathcal{P}_l(\text{Mod}^{\text{Su}}(\vdash)).$$

Proof Observe that \vdash is complete w.r.t. any of the classes $\text{Mod}(\vdash)$, $\text{Mod}^*(\vdash)$, $\text{Mod}^{\text{Su}}(\vdash)$. Moreover any of these classes contains the (trivial) matrix $\langle \mathbf{1}, \{1\} \rangle$. Thus we can apply Theorem 14. □

4 Logics with a partition function and axiomatizations

Definition 16 *A logic \vdash has a partition function if there is a formula $x \cdot y$, in which the variables x and y really occur, such that $x \vdash x \cdot y$ and the operation $\cdot^{\mathbf{A}}$ is a partition function for every $\mathbf{A} \in \text{Alg}(\vdash)$. In this case, $x \cdot y$ is a partition function for \vdash .*

Remark 17 By Lemma 4, the above definition can be rephrased in purely logical terms, by requiring that $x \vdash x \cdot y$ and that

$$\varphi(\epsilon, \vec{z}) \dashv\vdash \varphi(\delta, \vec{z}) \text{ for every formula } \varphi(v, \vec{z}),$$

for every identity of the form $\epsilon \approx \delta$ in **P1.**, . . . , **P5.** □

Example 18 Logics with a partition function abound in the literature. Indeed, it is easy to check that the term $x \cdot y := x \wedge (x \vee y)$ is a partition function for every logic \vdash such that every algebra in $\text{Alg}(\vdash)$ has a lattice reduct. Such examples include all modal and substructural logics. On the other hand, $x \cdot y := (y \rightarrow x) \rightarrow x$ is a partition function for all logics \vdash such that $\text{Alg}(\vdash)$ has a Hilbert algebra reduct [25]. □

Remarkably, the presence of a partition function is inherited by the variable inclusion companion of a logic.

Lemma 19 *Let \vdash be a logic. The operation \cdot is a partition function for \vdash if and only if it is a partition function for \vdash^l .*

Proof From Remark 17 the fact that \cdot is a partition function for \vdash is witnessed by the validity of some inferences $\varphi \vdash \psi$ such that $\text{Var}(\varphi) \subseteq \text{Var}(\psi)$. Hence these inferences also hold in \vdash^l . With another application of Remark 17 we conclude that \cdot is a partition function for \vdash^l .

The other direction follows from the inclusion $\vdash^l \subseteq \vdash$. □

The following result is the generalization of Theorem 7 to the setting of logical matrices.

Theorem 20 *Let \vdash be a logic with a partition function \cdot , and $\langle \mathbf{A}, F \rangle$ be a model of \vdash such that $\mathbf{A} \in \text{Alg}(\vdash)$. Conditions (1–4) of Theorem 7 hold. Moreover, setting $F_i := F \cap A_i$ for every $i \in I$, the triple*

$$X = \langle \langle I, \leq \rangle, \{ \langle \mathbf{A}_i, F_i \rangle \}_{i \in I}, \{ f_{ij} : i \leq j \} \rangle$$

is a directed system of matrices such that $\mathcal{P}_l(X) = \langle \mathbf{A}, F \rangle$.

Proof In the light of Theorem 7, it will be enough to show that $f_{ij}[F_i] \subseteq F_j$ for every $i, j \in I$ such that $i \leq j$. To this end, consider $a \in F_i$ and $b \in A_j$ with $i \leq j$. Since \cdot is a partition function for \vdash , we have $x \vdash x \cdot y$. Together with the fact that $\langle \mathbf{A}, F \rangle \in \text{Mod}(\vdash)$ and $a \in F$, this implies that $a \cdot^{\mathbf{A}} b \in F$. Observe that $a \cdot^{\mathbf{A}} b \in A_j$ by (2) in Theorem 7 and, therefore, that $a \cdot^{\mathbf{A}} b \in F_j$. Hence, by (3), we have that $f_{ij}(a) = a \cdot^{\mathbf{A}} b \in F_j$. □

Definition 21 *Let \vdash be a logic with a partition function \cdot , and $\langle \mathbf{A}, F \rangle$ be a model of \vdash such that $\mathbf{A} \in \text{Alg}(\vdash)$. The Płonka fibers of $\langle \mathbf{A}, F \rangle$ are the matrices $\{ \langle \mathbf{A}_i, F_i \rangle \}_{i \in I}$ given in the above result.*

Lemma 22 *Let \vdash be a finitary logic with partition function \cdot and $\langle \mathbf{A}, F \rangle \in \text{Mod}(\vdash^l)$, with $\mathbf{A} \in \text{Alg}(\vdash^l)$. Then, the Płonka fibers of $\langle \mathbf{A}, F \rangle$ are models of \vdash .*

Proof Let $\langle \mathbf{A}_i, F_i \rangle$ be a Płonka fiber of $\langle \mathbf{A}, F \rangle$ and $\Gamma \vdash \varphi$, with Γ a finite set. Then consider a homomorphism $v: \mathbf{Fm} \rightarrow \mathbf{A}_i$ such that $v[\Gamma] \subseteq F_i$. Then, there are cases: either Γ is empty or not. First, suppose $\Gamma = \emptyset$. Then clearly $\emptyset \vdash^l \varphi$. Since \mathbf{A}_i is a subalgebra of \mathbf{A} and $\langle \mathbf{A}, F \rangle$ is a model of \vdash^l , this implies that $v(\varphi) \in F \cap A_i = F_i$. Then consider the case where Γ is non-void. Then there are $\gamma_1, \dots, \gamma_n \in Fm$ such that $\Gamma = \{\gamma_1, \dots, \gamma_n\}$. Since \cdot is a partition function, we have $x \vdash x \cdot y$. In particular, this implies that $\varphi \vdash \varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots))$. Then $\Gamma \vdash \varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots))$. Since the variable inclusion constraint holds for this inference, we obtain that

$$\Gamma \vdash^l \varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots)).$$

Since \mathbf{A}_i is a subalgebra of \mathbf{A} and $\langle \mathbf{A}, F \rangle$ is a model of \vdash^l , this implies that

$$v(\varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots))) \in A_i \cap F = F_i.$$

Since $v(\varphi)$ and $v(\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots))$ belong to A_i , this implies that

$$\begin{aligned} v(\varphi) &= v(\varphi) \cdot v(\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots)) \\ &= v(\varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots))) \end{aligned}$$

and, therefore, that $v(\varphi) \in F_i$, as desired. □

By a *Hilbert-style calculus with finite rules* we understand a (possibly infinite) set of Hilbert-style rules, each of which has finitely many premises.

Definition 23 Let \mathcal{H} be a Hilbert-style calculus with finite rules that determines a logic \vdash with a partition function \cdot . Let \mathcal{H}^l be the Hilbert-style calculus given by the following rules:

$$\emptyset \triangleright \psi \tag{H1}$$

$$\gamma_1, \dots, \gamma_n \triangleright \varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots)) \tag{H2}$$

$$x \triangleright x \cdot y \tag{H3}$$

$$\chi(\epsilon, \vec{z}) \triangleleft \triangleright \chi(\delta, \vec{z}) \tag{H4}$$

for every

- (i) $\emptyset \triangleright \psi$ rule in \mathcal{H} ;
- (ii) $\gamma_1, \dots, \gamma_n \triangleright \varphi$ rule in \mathcal{H} ;
- (iii) equation $\epsilon \approx \delta$ in the definition of partition function, and formula $\chi(v, \vec{z})$.

Theorem 24 *Let \vdash be a logic with partition function \cdot defined by a Hilbert-style calculus with finite rules \mathcal{H} . Then \mathcal{H}^l is a complete Hilbert-style calculus for \vdash^l .*

Proof Let $\vdash_{\mathcal{H}^l}$ be the logic determined by \mathcal{H}^l . We begin by showing that $\vdash_{\mathcal{H}^l} \subseteq \vdash^l$. It will be sufficient to show that every rule in \mathcal{H}^l holds in \vdash^l . This is clear for (H1). Moreover, the rules (H3, H4) are valid in \vdash^l , because \cdot is a partition function for \vdash^l by Lemma 19. It only remains to prove that (H2) holds in \vdash^l . To this end, consider a rule $\gamma_1, \dots, \gamma_n \triangleright \varphi$ in \mathcal{H} . Clearly we have that $\gamma_1, \dots, \gamma_n \vdash \varphi$. Since \cdot is a partition function for \vdash , we have $x \vdash x \cdot y$. In particular, $\varphi \vdash \varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots))$. Hence we conclude that

$$\gamma_1, \dots, \gamma_n \vdash^l \varphi \cdot (\gamma_1 \cdot (\gamma_2 \cdot \dots (\gamma_{n-1} \cdot \gamma_n) \dots)),$$

as desired.

To prove $\vdash^l \subseteq \vdash_{\mathcal{H}^l}$, we reason as follows. Consider $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\text{Su}}(\vdash_{\mathcal{H}^l})$. Observe that clearly $\mathbf{A} \in \text{Alg}(\vdash_{\mathcal{H}^l})$. Moreover, \cdot is a partition function in $\vdash_{\mathcal{H}^l}$ by Remark 17 and (H3, H4). Hence we can apply Theorem 20, obtaining that $\langle \mathbf{A}, F \rangle = \mathcal{P}_I(X)$, where X is the directed system of matrices $\langle I, \{\langle \mathbf{A}_i, F_i \rangle\}_{i \in I}, \{f_{ij} : i \leq j\} \rangle$ given in the statement of Theorem 20. Thanks to the rules of \mathcal{H}^l we can replicate the construction in the proof of Lemma 22 obtaining that each fiber $\langle \mathbf{A}_i, F_i \rangle$ is a model of \vdash . This observation, together with the fact that $\langle \mathbf{A}, F \rangle = \mathcal{P}_I(X)$ and Corollary 15, implies that $\langle \mathbf{A}, F \rangle$ is a model of \vdash^l . Hence we conclude that $\text{Mod}^{\text{Su}}(\vdash_{\mathcal{H}^l}) \subseteq \text{Mod}(\vdash^l)$. This implies that $\vdash^l \subseteq \vdash_{\mathcal{H}^l}$. \square

The proof of the above result establishes the following:

Corollary 25 *If \vdash is a finitary logic with a partition function, then $\text{Mod}^{\text{Su}}(\vdash^l) \subseteq \mathcal{P}_I(\text{Mod}(\vdash))$.*

Example 26 A Hilbert-style calculus for PWK is axiomatized, following Definition 23, as follows ($\varphi \rightarrow \psi$ is a shorthand for $\neg\varphi \vee \psi$):

$$\emptyset \triangleright (\varphi \vee \varphi) \rightarrow \varphi \tag{A1}$$

$$\emptyset \triangleright \varphi \rightarrow (\varphi \vee \psi) \tag{A2}$$

$$\emptyset \triangleright (\varphi \vee \psi) \rightarrow (\psi \vee \varphi) \tag{A3}$$

$$\emptyset \triangleright (\varphi \rightarrow \psi) \rightarrow ((\gamma \vee \varphi) \rightarrow (\gamma \vee \psi)) \tag{A4}$$

$$\emptyset \triangleright (\varphi \wedge \psi) \rightarrow \neg(\neg\varphi \vee \neg\psi) \tag{A5}$$

$$\emptyset \triangleright \neg(\neg\varphi \vee \neg\psi) \rightarrow (\varphi \wedge \psi) \tag{A6}$$

$$\varphi, \varphi \rightarrow \psi \triangleright \psi \wedge (\psi \vee (\varphi \wedge (\varphi \vee (\varphi \rightarrow \psi)))) \tag{R1}$$

$$\varphi \triangleright \varphi \wedge (\varphi \vee \psi) \tag{R2}$$

$$\chi(\epsilon, \vec{z}) \triangleleft \triangleright \chi(\delta, \vec{z}) \tag{R^*}$$

Notice that Axioms (A1)–(A6), together with the rule of *Modus Ponens*, provide a Hilbert-style calculus for propositional classical logic. (R1) and (R2) are obtained by setting $x \cdot y := x \wedge (x \vee y)$ as partition function for classical logic. Note, moreover, that (R*) is in fact a rule scheme, summarizing an infinity of rules. \square

5 Suszko reduced models of \vdash^I

In this section we investigate the structure of the Suszko reduced models $\text{Mod}^{\text{Su}}(\vdash^I)$ of the variable inclusion companion \vdash^I of a logic \vdash (with partition function). To this end, we rely on the following technical observation:

Lemma 27 *Let \vdash be a logic with a partition function \cdot , and $X = \langle\langle I, \leq \rangle, \{\langle \mathbf{A}_i, F_i \rangle\}_{i \in I}, \{f_{ij} : i \leq j\}\rangle$ a directed system of models of \vdash . Given an upset $J \subseteq I$, we define for every $i \in I$,*

$$G_i := \begin{cases} A_i & \text{if } i \in J \\ F_i & \text{otherwise.} \end{cases}$$

Then $\bigcup_{i \in I} G_i$ is a \vdash^I -filter on $\mathcal{P}_I(\mathbf{A}_i)_{i \in I}$.

Proof It is clear that the matrices $\{\langle \mathbf{A}_i, G_i \rangle : i \in I\}$ give naturally rise to a directed system of matrices, when equipped with the homomorphisms in X . Moreover, by assumption each $\langle \mathbf{A}_i, G_i \rangle$ is a model of \vdash . Thus $\bigcup_{i \in I} G_i$ is a \vdash^I -filter on $\mathcal{P}_I(\mathbf{A}_i)_{i \in I}$ by Lemma 13. \square

The following result identifies the Plonka sums of matrices in $\text{Mod}^{\text{Su}}(\vdash)$ that belong to $\text{Mod}^{\text{Su}}(\vdash^I)$.

Theorem 28 *Let \vdash be a logic with a partition function \cdot , and let $X = \langle\langle I, \leq \rangle, \{\langle \mathbf{A}_i, F_i \rangle\}_{i \in I}, \{f_{ij} : i \leq j\}\rangle$ be a directed system of matrices in $\text{Mod}^{\text{Su}}(\vdash)$. The following conditions are equivalent:*

- (i) $\mathcal{P}_I(X) \in \text{Mod}^{\text{Su}}(\vdash^I)$.
- (ii) For every $n, i \in I$ such that $\langle \mathbf{A}_n, F_n \rangle$ is trivial and $n < i$, there exists $j \in I$ s.t. $n \leq j, i \not\leq j$ and \mathbf{A}_j is non-trivial.

Proof (i) \Rightarrow (ii): Suppose that $\mathcal{P}_I(X) \in \text{Mod}^{\text{Su}}(\vdash^I)$, and consider $n, i \in I$ such that $\langle \mathbf{A}_n, F_n \rangle$ is trivial and $n < i$. The fact that $\langle \mathbf{A}_n, F_n \rangle$ is both trivial and belongs to $\text{Mod}^{\text{Su}}(\vdash)$ implies that \mathbf{A}_n is the trivial algebra. Then $\langle \mathbf{A}_n, F_n \rangle = \langle \mathbf{1}, \{1\} \rangle$. Moreover,

set $a := f_{ni}(1)$. Since $n < i$, we know that $a \neq 1$. Together with the fact that $\mathcal{P}_i(X) \in \text{Mod}^{\text{Su}}(\dashv^l)$, this implies that there is a \dashv^l -filter G of $\mathcal{P}_i(\mathbf{A}_i)_{i \in I}$ such that $\bigcup_{i \in I} F_i \subseteq G$ and $\langle a, 1 \rangle \notin \Omega^{\mathcal{P}_i(\mathbf{A}_i)_{i \in I}} G$. Thus, by Lemma 2, there is a formula $\varphi(x, \vec{z})$ and elements $\vec{c} \in \bigcup_{i \in I} A_i$ such that

$$\varphi^{\mathcal{P}_i}(a, \vec{c}) \in G \iff \varphi^{\mathcal{P}_i}(1, \vec{c}) \notin G. \tag{2}$$

We can assume w.l.o.g. that all the elements in the sequence \vec{c} belong to the same component A_k of the Płonka sum $\mathcal{P}_i(\mathbf{A}_i)_{i \in I}$.²

We claim that indeed $\varphi^{\mathcal{P}_i}(1, \vec{c}) \notin G$. Suppose the contrary towards a contradiction. Then $\varphi^{\mathcal{P}_i}(1, \vec{c}) \in G$. First observe that

$$\varphi^{\mathcal{P}_i}(a, \vec{c}) = \varphi^{\mathbf{A}_{i \vee k}}(f_{i, i \vee k}(a), f_{k, i \vee k}(\vec{c})) \tag{3}$$

$$= \varphi^{\mathbf{A}_{i \vee k}}(f_{i, i \vee k} \circ f_{n, i}(1), f_{k, i \vee k}(\vec{c})) \tag{4}$$

$$= \varphi^{\mathbf{A}_{i \vee k}}(f_{n \vee k, i \vee k} \circ f_{n, n \vee k}(1), f_{n \vee k, i \vee k} \circ f_{k, n \vee k}(\vec{c})) \tag{5}$$

$$= f_{n \vee k, i \vee k} \varphi^{\mathbf{A}_{n \vee k}}(f_{n, n \vee k}(1), f_{k, n \vee k}(\vec{c})) \tag{6}$$

$$= f_{n \vee k, i \vee k} \varphi^{\mathcal{P}_i}(1, \vec{c}) \tag{7}$$

$$= f_{n \vee k, i \vee k}(\varphi^{\mathcal{P}_i}(1, \vec{c})) \cdot^{\mathbf{A}_{i \vee k}} f_{i, i \vee k}(a) \tag{8}$$

$$= \varphi^{\mathcal{P}_i}(1, \vec{c}) \cdot^{\mathcal{P}_i} a \tag{9}$$

$$\in G. \tag{10}$$

The above equalities are justified as follows: (5) is a consequence of the fact that X is a directed system of matrices and that $n \vee k \leq i \vee k$ (since $n \leq i$), (8) follows from the fact that $x \cdot^{\mathcal{P}_i} y$ is the projection on the first component on the algebra $\mathbf{A}_{i \vee k}$. Condition (10) follows from the fact that $\varphi^{\mathcal{P}_i}(1, \vec{c}) \in G$, G is a \dashv^l -filter and, by Lemma 19 \cdot is a partition function for \dashv^l , hence $x \dashv^l x \cdot y$. Hence we have that $\varphi^{\mathcal{P}_i}(a, \vec{c}), \varphi^{\mathcal{P}_i}(1, \vec{c}) \in G$, which contradicts (2), establishing the claim.

From the claim and (2) we get that $\varphi^{\mathcal{P}_i}(a, \vec{c}) \in G$ and $\varphi^{\mathcal{P}_i}(1, \vec{c}) \notin G$. Set $j := n \vee k$ and $m := k \vee i$. We claim that j is such that: (A) $n \leq j$, (B) \mathbf{A}_j is non trivial and (C) $i \not\leq j$. We proceed to prove (A, B, C).

(A): Since $j = n \vee k$, we have that $n \leq j$.

(B): Observe that

$$\varphi^{\mathcal{P}_i}(1, \vec{c}) = \varphi^{\mathbf{A}_j}(f_{nj}(1), f_{kj}(\vec{c})) \in A_j.$$

Together with $\varphi^{\mathcal{P}_i}(1, \vec{c}) \notin G$, this implies that $\varphi^{\mathcal{P}_i}(1, \vec{c}) \in A_j \setminus G$.

On the other hand, since $F_n = A_n$, we have that

$$f_{nj}(1) \in f_{nj}[F_n] \subseteq F_j \subseteq A_j \cap G.$$

² More precisely, if $\vec{c} = c_1, \dots, c_m$ and $c_1 \in A_{p_1}, \dots, c_m \in A_{p_m}$, then we set $k := p_1 \vee \dots \vee p_m$ and replace c_i by $f_{p_i k}(c_i)$.

Thus both $A_j \cap G$ and $A_j \setminus G$ are non-empty. We conclude that \mathbf{A}_j is non-trivial.

(C): Suppose, by contradiction, that $i \leq j$. In particular, this implies that $m = j$ (indeed, $i \leq j = n \vee k$, thus $i \vee k \leq n \vee k$, i.e. $m \leq j$; on the other hand, since $n < i$ then $n \vee k \leq i \vee j$, i.e. $j \leq m$). Therefore we have that

$$\varphi^{\mathcal{P}_i}(1, \vec{c}) = \varphi^{\mathbf{A}_j}(f_{nj}(1), f_{kj}(\vec{c})) \tag{11}$$

$$= \varphi^{\mathbf{A}_j}(f_{ij} \circ f_{ni}(1), f_{kj}(\vec{c})) \tag{12}$$

$$= \varphi^{\mathbf{A}_j}(f_{ij}(a), f_{kj}(\vec{c})) \tag{13}$$

$$= \varphi^{\mathbf{A}_m}(f_{im}(a), f_{km}(\vec{c})) \tag{14}$$

$$= \varphi^{\mathcal{P}_i}(a, \vec{c}) \in G. \tag{15}$$

The above equalities are justified as follows. (12) follows from the fact that $i \leq m = j$. (13) is a consequence of $a = f_{ni}(1)$. (14) from $j = m$ and (15) from $m = i \vee k$. This establishes the above equalities, yielding that $\varphi^{\mathcal{P}_i}(1, \vec{c}) \in G$. But this contradicts the fact that $\varphi^{\mathcal{P}_i}(1, \vec{c}) \notin G$.

Hence (A), (B) and (C) hold establishing our claim. In particular, this implies that $j \in I$ satisfies the condition in the statement.

(ii) \Rightarrow (i): By Lemma 13 we know that $\mathcal{P}_i(X)$ is a model of \vdash^I . It only remains to prove that it is Suszko reduced. To this end, let θ be the Suszko congruence of $\mathcal{P}_i(X)$.

Observe that, in order to prove that θ is the identity, it will be enough to show that it does not identify distinct elements in components of the Płonka sum which are comparable with respect to the order \leq . To prove this, suppose indeed that θ does not identify different elements in components of the Płonka sum which are comparable. Then consider two different elements $a, b \in A = \bigcup_{j \in I} A_j$. There exist $i, j \in I$ such that $a \in A_i$ and $b \in A_j$. If i and j are comparable, then by assumption $\langle a, b \rangle \notin \theta$. Then consider the case where i and j are incomparable. Set $k := i \vee j$. Clearly we have that $i, j < k$. In particular, we have that $b \cdot b = b \in A_j$ and $a \cdot b \in A_k$ and, therefore, $b \cdot^{\mathcal{P}_i} b \neq a \cdot^{\mathcal{P}_i} b$. Since j and k are comparable, this implies that $\langle b \cdot^{\mathcal{P}_i} b, a \cdot^{\mathcal{P}_i} b \rangle \notin \theta$. In particular, this means that $\langle a, b \rangle \notin \theta$ as well. As a consequence we conclude that θ is the identity.

By the above observation, to prove that θ is the identity, it will be enough to show that it does not identify elements in components of the Płonka sum $\mathcal{P}_i(X)$ which are comparable with respect to \leq . To this end, consider two different elements $a, b \in A$ such that $a \in A_i$ and $b \in A_j$ with $i \leq j$. We have two cases: either $i = j$ or $i < j$.

First consider the case where $i = j$, that is $a, b \in A_i$. By assumption, we have that $\langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\text{Su}}(\vdash)$. Therefore we can assume w.l.o.g. that there is a \vdash -filter G_i on \mathbf{A}_i such that $F_i \subseteq G_i$, some elements $\vec{c} \in A_i$, and a formula $\varphi(x, \vec{z})$ such that $\varphi^{\mathbf{A}_i}(a, \vec{c}) \in G_i$ and $\varphi^{\mathbf{A}_i}(b, \vec{c}) \notin G_i$. For every $l \neq i$, define

$$G_l := \begin{cases} A_l & \text{if } i \leq l \\ F_l & \text{otherwise.} \end{cases}$$

An analogous argument to the one described in the proof Lemma 27 shows that $G := \bigcup_{i \in I} G_i$ is a \vdash^l -filter on $\mathcal{P}_l(\mathbf{A}_i)_{i \in I}$. Moreover, observe that

$$\begin{aligned} \varphi^{\mathcal{P}_l}(a, \vec{c}) &= \varphi^{\mathbf{A}_i}(a, \vec{c}) \in G \\ \varphi^{\mathcal{P}_l}(b, \vec{c}) &= \varphi^{\mathbf{A}_i}(b, \vec{c}) \notin G. \end{aligned}$$

We conclude that $\langle a, b \rangle \notin \theta$.

Then we consider the case where $i < j$. We have cases: either \mathbf{A}_i is trivial or not. If \mathbf{A}_i is non-trivial, then $F_i \neq A_i$ as $\langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\text{Su}}(\vdash)$. Then for every $l \in I$, we define

$$G_l := \begin{cases} A_l & \text{if } i < l \\ F_l & \text{otherwise.} \end{cases}$$

By Lemma 27 we know that $G := \bigcup_{i \in I} G_i$ is a \vdash^l -filter on $\mathcal{P}_l(\mathbf{A}_i)_{i \in I}$. Then choose an element $c \in A_i \setminus F_i$. We have that

$$c \cdot^{\mathcal{P}_l} a = c \cdot^{\mathbf{A}_i} a = c \in A_i \setminus F_i = A_i \setminus G_i$$

and $c \cdot^{\mathcal{P}_l} b \in A_j = G_j$. Therefore, $c \cdot^{\mathcal{P}_l} a \notin G$ and $c \cdot^{\mathcal{P}_l} b \in G$. Hence we conclude that $\langle a, b \rangle \notin \theta$, as desired.

Then we consider the case where \mathbf{A}_i is trivial. We have cases: either $F_i = \emptyset$ or $F_i = A_i$. First suppose that $F_i = \emptyset$. Iterating the argument in the previous paragraph (taking $c := a$) we obtain that $\langle a, b \rangle \notin \theta$. Then consider the case where $F_i = A_i$. Observe that in this case $\langle \mathbf{A}_i, F_i \rangle$ is a trivial matrix. Therefore we can apply the assumption, obtaining an element $k \in I$ such that \mathbf{A}_k is non-trivial, $i < k$ and $j \not\leq k$. Then for every $l \in I$ we define

$$G_l := \begin{cases} A_l & \text{if } k \vee j \leq l \\ F_l & \text{otherwise.} \end{cases}$$

By Lemma 27 we know that $G := \bigcup_{i \in I} G_i$ is a \vdash^l -filter on $\mathcal{P}_l(\mathbf{A}_i)_{i \in I}$. Since \mathbf{A}_k is non-trivial and $\langle \mathbf{A}_k, F_k \rangle \in \text{Mod}^{\text{Su}}(\vdash)$, there is $c \in A_k \setminus F_k$. Since $k < k \vee j$, we have that

$$\begin{aligned} c \cdot^{\mathcal{P}_l} a &= c \cdot^{\mathbf{A}_k} f_{ik}(a) = c \in A_k \setminus F_k = A_k \setminus G_k \\ c \cdot^{\mathcal{P}_l} b &\in A_{j \vee k} = G_{j \vee k}. \end{aligned}$$

Hence we conclude that $c \cdot^{\mathcal{P}_l} a \notin G$ and $c \cdot^{\mathcal{P}_l} b \in G$. But this means that $\langle a, b \rangle \notin \theta$. \square

Theorem 28 identifies the Suszko reduced models of \vdash^l that can be expressed in terms of Płonka sums of Suszko reduced models of \vdash . It is natural to wonder whether it is true that *all* Suszko reduced models of \vdash^l are of this kind. Example 42 shows that this does not hold in general. A full characterization of the class of Suszko reduced models can be given for the (left) variable inclusion companions of logics which have

stronger properties, such as equivalential and finitary, or having inconsistency terms. These descriptions are addressed in the following subsections.

5.1 Equivalential logics

It turns out that, in the setting of finitary equivalential logics \vdash , the class of matrices $\text{Mod}^{\text{Su}}(\vdash^l)$ has a very transparent description in terms of Płonka sums, as we proceed to prove (see Theorem 31).

Lemma 29 *Let \vdash be an equivalential finitary logic with a partition function. Then*

$$\text{Mod}^*(\vdash^l) \subseteq \mathcal{P}_l(\text{Mod}^*(\vdash)).$$

Proof Recall from Lemma 19 that also \vdash^l has a partition function. Then consider $\langle \mathbf{A}, F \rangle \in \text{Mod}^*(\vdash^l)$ and let

$$X = \langle \langle I, \leq \rangle, \{ \langle \mathbf{A}_i, F_i \rangle \}_{i \in I}, \{ f_{ij} : i \leq j \} \rangle$$

be the directed system of matrices given in Theorem 20. We know that $\mathcal{P}_l(X) = \langle \mathbf{A}, F \rangle$. Moreover, by Lemma 22, we know that each fiber of X is a model of \vdash . It only remains to prove that the fibers of X are Leibniz reduced.

We claim that $\bigcup_{i \in I} \Omega^{\mathbf{A}_i} F_i$ is a congruence of \mathbf{A} . To show this, let $\Delta(x, y)$ be a set of congruence formulas for \vdash . Then consider an n -ary basic operation λ and elements $a_1, \dots, a_n, b_1, \dots, b_n \in A$ such that $\langle a_j, b_j \rangle \in \bigcup_{i \in I} \Omega^{\mathbf{A}_i} F_i$, for all $1 \leq j \leq n$. This implies that are indexes $m_1, \dots, m_n \in I$ such that $a_j, b_j \in A_{m_j}$, for all $j \leq n$, and moreover that $\langle a_j, b_j \rangle \in \Omega^{\mathbf{A}_{m_j}} F_{m_j}$. The fact that Δ is a set of congruence formulas for \vdash implies that

$$\Delta^{\mathcal{P}_l}(a_j, b_j) = \Delta^{\mathbf{A}_{m_j}}(a_j, b_j) \subseteq F_{m_j}.$$

Set $k := m_1 \vee \dots \vee m_n$. We have that

$$\bigcup_{j \leq n} \Delta^{\mathbf{A}_k}(f_{m_j k}(a_j), f_{m_j k}(b_j)) \subseteq F_k \tag{16}$$

From the fact that Δ is a set of congruence formulas for \vdash it follows that (recall that λ is an n -ary arbitrary operation)

$$\bigcup_{j \leq n} \Delta(x_j, y_j) \vdash \Delta(\lambda(\vec{x}), \lambda(\vec{y})). \tag{17}$$

Together with (16) and (17), the fact that $\langle \mathbf{A}_k, F_k \rangle$ is a model of \vdash implies that

$$\begin{aligned} &\Delta^{\mathbf{A}_k}(\lambda^{\mathcal{P}_i}(a_1, \dots, a_n), \lambda^{\mathcal{P}_i}(b_1, \dots, b_n)) \\ &= \Delta^{\mathbf{A}_k}(\lambda(f_{m_{1k}}(a_1), \dots, f_{m_{nk}}(a_n)), \lambda(f_{m_{1k}}(b_1), \dots, f_{m_{nk}}(b_n))) \\ &\subseteq F_k. \end{aligned}$$

Together with the fact that Δ is a set of congruence formulas for \vdash , this implies that

$$\langle \lambda^{\mathcal{P}_i}(\vec{a}), \lambda^{\mathcal{P}_i}(\vec{b}) \rangle \in \Omega^{\mathbf{A}_k} F_k \subseteq \bigcup_{i \in I} \Omega^{\mathbf{A}_i} F_i.$$

This establishes the claim.

Since each $\Omega^{\mathbf{A}_i} F_i$ is compatible with F_i , we know that the congruence $\bigcup_{i \in I} \Omega^{\mathbf{A}_i} F_i$ is compatible with F . In particular, this implies that $\bigcup_{i \in I} \Omega^{\mathbf{A}_i} F_i \subseteq \Omega^{\mathbf{A}} F$. Since $\Omega^{\mathbf{A}} F$ is the identity relation, we conclude that so is each $\Omega^{\mathbf{A}_i} F_i$. Hence we obtain that $\langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^*(\vdash)$ for every $i \in I$ and, therefore, that

$$\langle \mathbf{A}, F \rangle = \mathcal{P}_i(X) \subseteq \mathcal{P}_i(\text{Mod}^*(\vdash)).$$

We conclude that $\text{Mod}^*(\vdash^l) \subseteq \mathcal{P}_i(\text{Mod}^*(\vdash))$, as desired. □

Corollary 30 *If \vdash is an equivalential finitary logic with a partition function, then*

$$\text{Mod}^{\text{Su}}(\vdash^l) \subseteq \mathcal{P}_i(\text{Mod}^*(\vdash)) = \mathcal{P}_i(\text{Mod}^{\text{Su}}(\vdash)).$$

Proof First recall that $\text{Mod}^{\text{Su}}(\vdash) = \text{Mod}^*(\vdash)$, since \vdash is equivalential. Thus it will be enough to prove that $\text{Mod}^{\text{Su}}(\vdash^l) \subseteq \mathcal{P}_i(\text{Mod}^*(\vdash))$. We have that

$$\text{Mod}^{\text{Su}}(\vdash^l) = \mathbb{P}_{\text{SD}} \text{Mod}^*(\vdash^l) \tag{18}$$

$$\subseteq \mathbb{SPP} \text{Mod}^*(\vdash^l) \tag{19}$$

$$\subseteq \mathbb{SPP} \mathcal{P}_i(\text{Mod}^*(\vdash)) \tag{20}$$

$$\subseteq \mathcal{P}_i(\mathbb{SPP} \text{Mod}^*(\vdash)) \tag{21}$$

$$= \mathcal{P}_i(\text{Mod}^*(\vdash)). \tag{22}$$

The non-trivial inclusions above are justified as follows: (20) is a consequence of Lemma 29, (21) follows from Lemma 9, and (22) from the fact that $\text{Mod}^*(\vdash)$ is closed under \mathbb{S} and \mathbb{P} , since \vdash is equivalential. Hence we conclude that $\text{Mod}^{\text{Su}}(\vdash^l) \subseteq \mathcal{P}_i(\text{Mod}^*(\vdash))$. □

We are now ready to provide a full characterization of the Suszko reduced models of the variable inclusion companion of a finitary equivalential logic (with partition function).

Theorem 31 *Let \vdash be an equivalential and finitary logic with a partition function, and $\langle \mathbf{A}, F \rangle$ be a matrix. The following conditions are equivalent:*

- (i) $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\text{Su}}(\vdash^I)$.
- (ii) *There exists a directed system of matrices $X \subseteq \text{Mod}^*(\vdash)$ indexed by a semilattice I such that $\langle \mathbf{A}, F \rangle = \mathcal{P}_I(X)$ and for every $n, i \in I$ such that $\langle \mathbf{A}_n, F_n \rangle$ is trivial and $n < i$, there exists $j \in I$ s.t. $n \leq j, i \not\leq j$ and \mathbf{A}_j is non-trivial.*

Proof This is a consequence of Theorem 28 and Corollary 30. □

Example 32 Observe that all substructural logics [31,49] are finitary, equivalential, and have a partition function. The same holds for all local and global consequences of normal modal logics [43]. As a consequence, the above result provides a description of the Suszko reduced models of the left variable inclusion companions of all substructural and modal logics (when the latter are understood as local and global consequences of normal modal logics [3,17,42]). □

5.2 Inconsistency terms

The following definition originates in [45], but see also [15,58]:

Definition 33 *A logic \vdash has a set of inconsistency terms if there is a set of formulas Σ such that $\sigma[\Sigma] \vdash \varphi$ for every substitution σ and formula φ .*

Example 34 For any formula φ , the set $\{\neg(\varphi \rightarrow \varphi)\}$ is a set of inconsistency terms for all superintuitionistic logics, all axiomatic extensions of MTL-logic [19,26] including Łukasiewicz logic [18], and all local and global consequences of normal modal logics. □

Remark 35 Observe that if \vdash has a set of inconsistency terms, then \vdash has a set of inconsistency terms only in variable x . If, moreover, \vdash is finitary, then it has a *finite* set of inconsistency terms only in variable x . □

The goal of this section is to show that if \vdash is a logic with a set of inconsistency terms, then the description of the Suszko reduced models of its variable inclusion companion can be substantially improved (see Theorems 28, 31), as we show in this section.

The next result discloses the semantic meaning of inconsistency terms. It should be observed that algebraic versions of it first appeared in [16,41] in the setting of varieties and quasi-varieties of algebras respectively.

Lemma 36 *Let \vdash be a logic. The following are equivalent:*

- (i) \vdash has a set of inconsistency terms Σ .
- (ii) *If $\langle \mathbf{A}, F \rangle \in \text{Mod}(\vdash)$ is non-trivial, then it has no trivial submatrix.*

Proof (i) \Rightarrow (ii): Suppose that \vdash has a set of inconsistency terms Σ . We can assume w.l.o.g. that Σ is in variable x only. Suppose, in view of a contradiction, that there is a non-trivial matrix $\langle \mathbf{A}, F \rangle \in \text{Mod}(\vdash)$ with a trivial submatrix $\langle \mathbf{B}, B \rangle$. Since $\langle \mathbf{A}, F \rangle$ is non trivial, there exists an element $a \in A \setminus F$. Consider any homomorphism $v: \mathbf{Fm} \rightarrow \mathbf{A}$ such that $v(x) = b$ and $v(y) = a$, where b is any element of B . Since

$\Sigma = \Sigma(x)$ and $\langle \mathbf{B}, B \rangle$ is a submatrix of $\langle \mathbf{A}, F \rangle$, we have that $v[\Sigma] \subseteq B \subseteq F$. Together with the fact that $\Sigma \vdash y$, this implies that $a = v(y) \in F$, which is a contradiction.

(ii) \Rightarrow (i): Let $Fm(x)$ be the set of formulas in variable x only. We show that $Fm(x)$ is a set of inconsistency terms for \vdash . To this end, consider a substitution σ and a formula ψ . It is enough to show that $\sigma[Fm(x)] \vdash \psi$. Let $\varphi := \sigma(x)$. Observe that $\sigma[Fm(x)]$ coincides with the universe of the subalgebra $\text{Sg}^{\mathbf{Fm}}(\varphi)$ of \mathbf{Fm} generated by φ . Consider the matrices

$$\begin{aligned} M_1 &:= \langle \mathbf{Fm}, \text{Cn}_{\vdash}(\text{Sg}^{\mathbf{Fm}}(\varphi)) \rangle \\ M_2 &:= \langle \text{Sg}^{\mathbf{Fm}}(\varphi), \text{Sg}^{\mathbf{Fm}}(\varphi) \rangle. \end{aligned}$$

Clearly, M_1 is a model of \vdash and M_2 a trivial submatrix of M_1 . By the assumption, we get that M_1 is a trivial matrix, i.e. $Fm = \text{Cn}_{\vdash}(\text{Sg}^{\mathbf{Fm}}(\varphi))$. Hence we conclude that

$$\psi \in Fm = \text{Cn}(\text{Sg}^{\mathbf{Fm}}(\varphi)) = \text{Cn}_{\vdash}(\sigma[Fm(x)]).$$

Clearly this implies that $\sigma[Fm(x)] \vdash \psi$, as desired. □

Remarkably, Theorem 28 can be substantially improved for logics possessing a set of inconsistency terms (whose presence is essential, as shown in Example 43):

Theorem 37 *Let \vdash be a logic with a partition function and a set of inconsistency terms. For every directed system X of matrices in $\text{Mod}^{\text{Su}}(\vdash)$, the following conditions are equivalent:*

- (i) $\mathcal{P}_I(X) \in \text{Mod}^{\text{Su}}(\vdash^I)$.
- (ii) X contains at most one trivial component.

Proof For the sake of simplicity, throughout the proof we set

$$X = \langle \langle I, \leq \rangle, \{ \langle \mathbf{A}_i, F_i \rangle \}_{i \in I}, \{ f_{ij} : i \leq j \} \rangle.$$

First we claim that if a component $\langle \mathbf{A}_n, F_n \rangle$ of X is trivial, then so is $\langle \mathbf{A}_k, F_k \rangle$, for every $k \geq n$. To prove this, consider a trivial component $\langle \mathbf{A}_n, F_n \rangle$ of X and $k \geq n$. Observe that

$$f_{nk}[A_n] = f_{nk}[F_n] \subseteq F_k.$$

Then $\langle f_{nk}[A_n], f_{nk}[F_n] \rangle$ is a trivial submatrix of $\langle \mathbf{A}_k, F_k \rangle$. Since \vdash has a set of inconsistency terms, we can apply Lemma 36 obtaining that $\langle \mathbf{A}_k, F_k \rangle$ is trivial. This establishes the claim.

(i) \Rightarrow (ii): Suppose, in view of a contradiction, that $\mathcal{P}_I(X) \in \text{Mod}^{\text{Su}}(\vdash^I)$ and that X contains two distinct trivial components $\langle \mathbf{1}_n, \{1_n\} \rangle$ and $\langle \mathbf{1}_k, \{1_k\} \rangle$ (their algebraic reducts are trivial, as the components of X belong to $\text{Mod}^{\text{Su}}(\vdash)$). Set $\langle \mathbf{A}, F \rangle := \mathcal{P}_I(X)$. Observe that, for every formula $\varphi(x, \vec{z})$ in which x really occurs, and every tuple $\vec{c} \in A$, we have that

$$\varphi^{\mathbf{A}}(1_n, \vec{c}), \varphi^{\mathbf{A}}(1_k, \vec{c}) \in F.$$

To prove this, observe that the element $\varphi^{\mathbf{A}}(1_n, \vec{c})$ belongs to a component $\langle \mathbf{A}_l, F_l \rangle$ of X with $n \leq l$. By the previous claim, we know that $\langle \mathbf{A}_l, F_l \rangle$ is trivial and, therefore, that $\varphi^{\mathbf{A}}(1_n, \vec{c}) \in F_l \subseteq F$, as desired. A similar argument shows that $\varphi^{\mathbf{A}}(1_k, \vec{c}) \in F$ as well. Hence for every unary polynomial function p of \mathbf{A} we have that

$$\text{Fg}_{\vdash}^{\mathbf{A}}(F \cup \{p(1_n)\}) = \text{Fg}_{\vdash}^{\mathbf{A}}(F \cup \{p(1_k)\}).$$

By Lemma 2 this implies that $\langle 1_n, 1_k \rangle \in \tilde{\mathfrak{M}}_{\vdash}^{\mathbf{A}} F$. Since $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\text{Su}}(\vdash^l)$, this implies that $1_n = 1_k$, which is a contradiction.

(ii) \Rightarrow (i): Suppose that X contains at most one trivial matrix. If X contains no trivial component, then, by Theorem 28, we obtain that $\mathcal{P}_l(X) \in \text{Mod}^{\text{Su}}(\vdash^l)$. Then consider the case where X contains exactly one trivial component. By the claim we obtain that this component is the maximum of $\langle I, \leq \rangle$. Again, with an application of Theorem 28, we conclude that $\mathcal{P}_l(X) \in \text{Mod}^{\text{Su}}(\vdash^l)$. \square

The assumption on the existence of a set of inconsistency terms for the logic \vdash in the above theorem is essential, as shown in Example 43.

Drawing consequences from Theorem 37, we obtain a very transparent description of the Suszko reduced models of the variable inclusion companion of a finitary equivalential logic with a partition function and inconsistency terms:

Theorem 38 *Let \vdash be an equivalential and finitary logic with a partition function and inconsistency terms, and $\langle \mathbf{A}, F \rangle$ be a matrix. The following conditions are equivalent:*

- (i) $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\text{Su}}(\vdash^l)$.
- (ii) *There exists a directed system of matrices $X \subseteq \text{Mod}^*(\vdash)$ with at most one trivial component such that $\langle \mathbf{A}, F \rangle = \mathcal{P}_l(X)$.*

Proof This is a combination of Theorems 37 and 31. \square

Example 39 It is worth to observe that the above result provides a full description of the Suszko reduced models of the left variable inclusion companions of most well-known logics, including all logics mentioned in Example 34. \square

6 Classification in the Leibniz hierarchy

We conclude this work by investigating the location of logics of variable inclusion in the Leibniz hierarchy. To this end, recall that a logic \vdash is inconsistent if $\Gamma \vdash \varphi$ for every $\Gamma \cup \{\varphi\} \subseteq Fm$. Equivalently, \vdash is inconsistent if $\emptyset \vdash x$ for some variable x . A logic is *consistent* when it is not inconsistent.

Theorem 40 *Let \vdash be a logic.*

- (i) *If \vdash is consistent, then \vdash^l is not protoalgebraic.*
- (ii) *If \vdash is finitary, algebraizable and has a partition function, then \vdash^l is truth-equational.*

Proof (i) We reason by contraposition. Suppose that \vdash^l is protoalgebraic. Then there is a set of formulas $\Delta(x, y)$ such that $\emptyset \vdash^l \Delta(x, x)$ and $x, \Delta(x, y) \vdash^l y$. Thus, the definition of \vdash^l implies that there is a subset $\Sigma(y) \subseteq \Delta(x, y)$ such that $\Sigma(y) \vdash y$. Since $\emptyset \vdash^l \Delta(x, x)$, we have that $\emptyset \vdash^l \Sigma(y)$. From $\Sigma(y) \vdash y$ and $\emptyset \vdash^l \Sigma(y)$ it follows that $\emptyset \vdash^l y$. By the definition of \vdash^l we conclude that $\emptyset \vdash y$ and, therefore, that \vdash is inconsistent.

(ii) Suppose that \vdash is finitary, algebraizable and has a partition function. In particular, \vdash is truth-equational with set of defining equations $\tau(x)$. We will show that $\tau(x)$ is a set of defining equations for \vdash^l as well. To this end, consider $\langle \mathbf{A}, F \rangle \in \text{Mod}^*(\vdash^l)$. Since \vdash is finitary, equivalential and with a partition function, we can apply Lemma 29 obtaining that there exists a directed system of matrices $X \subseteq \text{Mod}^*(\vdash)$ such that $\langle \mathbf{A}, F \rangle = \mathcal{P}_l(X)$. For the sake of simplicity, we set

$$X = \langle \langle I, \leq \rangle, \{ \langle \mathbf{A}_i, F_i \rangle \}_{i \in I}, \{ f_{ij} : i \leq j \} \rangle$$

and assume w.l.o.g. that $\langle \mathbf{A}, F \rangle = \mathcal{P}_l(X)$. Consider an element $a \in A$. There is $i \in I$ such that $a \in A_i$. We have that

$$\mathbf{A} \models \tau(a) \iff \mathbf{A}_i \models \tau(a) \iff a \in F_i \iff a \in F. \tag{23}$$

The above equivalences are justified as follows. The first one follows from the fact that $\mathbf{A} = \mathcal{P}_l(\mathbf{A}_i)_{i \in I}$. The second one follows from the fact that $\langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^*(\vdash)$ and that $\tau(x)$ is a set of defining equations for \vdash . The last one follows from the observation that $\langle \mathbf{A}, F \rangle = \mathcal{P}_l(X)$.

By (23) we obtain that for every $a \in A$,

$$\mathbf{A} \models \tau(a) \iff a \in F.$$

Hence we conclude that $\tau(x)$ is a set of defining equations for \vdash^l and, therefore, \vdash^l is truth-equational. □

In [10, Theorem 48] it is proved that the variety of involutive bisemilattices, i.e. the closure under Plonka sums of the variety of Boolean algebras [50], is not the equivalent algebraic semantics of any algebraizable logic. This result can be strengthened as follows:

Theorem 41 *Let K be a class of algebras containing two trivial algebras and closed under Plonka sums. There is no protoalgebraic logic \vdash such that $\text{Alg}(\vdash) = K$.*

Proof Suppose, in view of a contradiction, that there are a class of algebras K containing two trivial algebras and closed under Plonka sums, and a protoalgebraic logic \vdash such that $\text{Alg}(\vdash) = K$. Let $\mathbf{1}_a, \mathbf{1}_b \in K$ be distinct trivial algebras and consider the directed system obtained by the homomorphism $f_{ab} : \mathbf{1}_a \rightarrow \mathbf{1}_b$ ($a \leq b$ in the semilattice order of the indexes). Let $\mathbf{A} = \mathbf{1}_a \oplus \mathbf{1}_b$ be the Plonka sum of this directed system. Clearly $\mathbf{A} \in K$. Therefore there is $F \subseteq A$ such that $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\text{Su}}(\vdash)$. As $\text{Alg}(\vdash)$ contains a non-trivial algebra, it is not difficult to see that $x \not\vdash y$. Since \vdash is protoalgebraic, there is a set of formulas $\Delta(x, y)$ such that $\emptyset \vdash \Delta(x, x)$ and $x, \Delta(x, y) \vdash y$. Since $x \not\vdash y$

and $x, \Delta(x, y) \vdash y$, we conclude that $\Delta(x, y) \neq \emptyset$. Then consider $\varphi(x, y) \in \Delta(x, y)$. Since $\emptyset \vdash \Delta(x, x)$, we conclude that $\emptyset \vdash \varphi(x, x)$.

Now, observe that the variable x really occurs in $\varphi(x, x)$, since we do not allow the presence of constant symbols in this paper. Hence we obtain that

$$\varphi^{\mathbf{A}}(1_a, 1_a) = 1_a \text{ and } \varphi^{\mathbf{A}}(1_b, 1_b) = 1_b.$$

Together with the fact that $\emptyset \vdash \varphi(x, x)$, this implies that $A = \{1_a, 1_b\}$ is the smallest \vdash -filter on \mathbf{A} . In particular, this implies that A is the *unique* \vdash -filter on \mathbf{A} . Since F is a \vdash -filter on \mathbf{A} , we conclude that $A = F$. Hence $\langle \mathbf{A}, A \rangle$ is a Suszko reduced model of \vdash . This implies that \mathbf{A} is trivial, which is false. \square

Acknowledgements The first and the second author were both supported by the Grant GBP202/12/G061 of the Czech Science Foundation. The first author acknowledges also the ERC Grant: "Philosophy of Pharmacology: Safety, Statistical Standards, and Evidence Amalgamation", GA:639276. The second author was supported also by a Beatriz Galindo grant BEAGAL18/00040 of the Spanish Ministry of Science, Innovation and Universities. The third author gratefully acknowledges the support of the Regione Autonoma della Sardegna within the project: "Per un'estensione semantica della Logica Computazionale Quantistica-Impatto teorico e ricadute", CUP:F76C18001020002. We are grateful to an anonymous referee for his/her valuable comments and suggestions.


Appendix

Aim of this section is showing that some of the assumptions are indeed essential in order to prove certain results. In particular, the following example shows that, in general, there can be Suszko reduced models of the logic \vdash^I that are not Płonka sums of Suszko reduced models of \vdash .

Example 42 Consider the logic \vdash determined by the following class of matrices:

$$\mathbf{M} := \{ \langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a distributive lattice and } F \text{ is an upset} \}.$$

Let \mathbf{A}_1 be the three element lattice $a < b < c$ and let $F_1 = \{b, c\}$. Moreover, let \mathbf{A}_2 be the four-element Boolean lattice (with universe $\{0, d, e, 1\}$ with 0 as bottom element), and let $F_2 = A_2 \setminus \{0\}$. Clearly both $\langle \mathbf{A}_1, F_1 \rangle$ and $\langle \mathbf{A}_2, F_2 \rangle$ are models of \vdash (as they belong to \mathbf{M}). However, it is easy to see that $\langle \mathbf{A}_1, F_1 \rangle \notin \text{Mod}^{\text{Su}}(\vdash)$. Now, let $f: \mathbf{A}_1 \rightarrow \mathbf{A}_2$ be any of the two embeddings of \mathbf{A}_1 into \mathbf{A}_2 . Clearly these two matrices plus f give rise to a directed system X of matrices (of course one should pedantically add the identity endomorphisms) depicted in the following figure. We denote by $\langle \mathbf{B}, G \rangle$ the Płonka sum $\mathcal{P}_f(X)$.



Since $\langle \mathbf{A}_1, F_1 \rangle$ and $\langle \mathbf{A}_2, F_2 \rangle$ are models of \vdash , by Lemma 13 $\langle \mathbf{B}, G \rangle$ is a model of \vdash^I . We now show that it is indeed Suszko reduced. Elements belonging to the algebra \mathbf{A}_1 , as for example b and c (any other pair of elements in \mathbf{A}_1 is distinguished by the identity function), can be distinguished by means of the function $\wedge^{\mathbf{B}}$, the filter G and the element e , as follows:

$$\begin{aligned}
 b \wedge^{\mathbf{B}} e &= d \wedge^{\mathbf{A}_2} e = 0 \notin G \\
 c \wedge^{\mathbf{B}} e &= 1 \wedge^{\mathbf{A}_2} e = e \in G.
 \end{aligned}$$

One can reason similarly (using G as filter) for pairs of elements belonging to \mathbf{A}_2 (we illustrate the only interesting case):

$$\begin{aligned}
 d \wedge^{\mathbf{B}} b &= d \wedge^{\mathbf{A}_2} d = d \in G \\
 e \wedge^{\mathbf{B}} b &= e \wedge^{\mathbf{A}_2} d = 0 \notin G.
 \end{aligned}$$

On the other hand, pairs of elements belonging to different algebras are distinguished by considering the filter $H := F_1 \cup A_2$ on \mathbf{B} (the fact that it is a filter is guaranteed by Lemma 27), the function $\wedge^{\mathbf{B}}$ and the element a . Consider, for instance, the elements b and d :

$$\begin{aligned}
 b \wedge^{\mathbf{B}} a &= a \notin H; \\
 d \wedge^{\mathbf{B}} a &= d \wedge^{\mathbf{A}_2} 0 = 0 \in H.
 \end{aligned}$$


This is enough to show that $\langle \mathbf{B}, G \rangle$ is Suszko reduced.

To conclude the example we need to disprove that $\langle \mathbf{B}, G \rangle$ is a Płonka sum of any Suszko reduced models of \vdash . Suppose that $\langle \mathbf{B}, G \rangle$ is the Płonka sum of a directed system Y of Suszko reduced models $\langle \mathbf{B}_1, G_1 \rangle, \dots, \langle \mathbf{B}_n, G_n \rangle$ of \vdash . First observe that $n \leq 2$. Suppose the contrary towards a contradiction. Then $n \geq 3$. We choose three elements $b_1 \in B_1, b_2 \in B_2$ and $b_3 \in B_3$. Clearly b_1, b_2 and b_3 are different. Moreover, for every $1 \leq i < j \leq 3$ we have that either $b_i \cdot^{\mathbf{B}} b_j \neq b_i$ or $b_j \cdot^{\mathbf{B}} b_i \neq b_j$, where \cdot indicates the partition function, i.e. $x \cdot y := x \wedge (x \vee y)$. It is easy to see that no such three elements exist in \mathbf{B} , which is a contradiction. Hence $n \leq 2$. We have cases. If $n = 1$, then $\langle \mathbf{B}_1, G_1 \rangle = \langle \mathbf{B}, G \rangle$. In particular, this implies that $\langle \mathbf{B}, G \rangle \in \text{Mod}^{\text{Su}}(\vdash)$ and, therefore, $\mathbf{B} \in \text{Alg}(\vdash)$. By Lemma 3 this implies that \mathbf{B} is a lattice, which is false.

Thus, the only possible case is that $n = 2$. Now, by Lemma 3 we know that \mathbf{B}_1 and \mathbf{B}_2 are distributive lattices. Since the only way of partitioning \mathbf{B} into two subalgebras that are distributive lattices is $\{\mathbf{A}_1, \mathbf{A}_2\}$, we conclude that w.l.o.g. $\mathbf{B}_1 = \mathbf{A}_1$ and $\mathbf{B}_2 = \mathbf{A}_2$, i.e. $\langle \mathbf{B}, G \rangle$ can not be the Płonka sum of any Suszko reduced models of \vdash . \square

Example 43 The statement of Theorem 37 is in general false for logics without a set of inconsistency terms, as witnessed by the following example based on $\text{CL}^{\wedge, \vee}$, the conjunction and disjunction fragment of classical propositional logic (which does not possess a set of inconsistency terms). In particular, it happens to have a Suszko reduced model of \vdash^I , which is the Płonka sum of a directed system of Suszko reduced models of \vdash containing two trivial matrices.

Let \vdash be the \wedge, \vee -fragment of classical propositional logic. Moreover, let $\mathbf{1}$ be the trivial lattice and $\mathbf{L}_2 = \{\perp, \top\}$ the 2-element distributive lattice (with $\perp < \top$). Consider the directed system X of matrices formed by 6 copies of the matrix $\langle \mathbf{L}_2, \{\top\} \rangle$ and two trivial matrices $\langle \mathbf{1}, \{1\} \rangle$ sketched in the following figure (lines represent lattice order in the Płonka fibers, arrows, the homomorphisms, and circles, filters in any fiber).



Clearly each matrix in X , which contains two trivial matrices, is a Suszko reduced model of \vdash . Moreover, by applying Theorem 28, one immediately checks that $\mathcal{P}_I(X) \in \text{Mod}^{\text{Su}}(\vdash^I)$. \square

References

1. Bergman, C.: Universal Algebra: Fundamentals and Selected Topics. Chapman and Hall/CRC, Berlin (2011)
2. Bergman, C., Failing, D.: Commutative idempotent groupoids and the constraint satisfaction problem. Algebra Univers. **73**(3), 391–417 (2015)
3. Blackburn, P., de Rijke, M., Venema, Y.: Modal Logic. Cambridge University Press, Cambridge (2001)
4. Blok, W., Pigozzi, D.: Algebraizable Logics. American Mathematical Society, New York (1989)
5. Blok, W., Pigozzi, D.: Protoalgebraic logics. Stud. Log. **45**, 337–369 (1986)
6. Blok, W., Pigozzi, D.: Algebraic semantics for universal Horn logic without equality. In: Romanowska, A., Smith, J. (eds.) Universal Algebra and Quasigroup Theory, pp. 1–56. Heldermann, Berlin (1992)

7. Bochvar, D.: On a three-valued calculus and its application in the analysis of the paradoxes of the extended functional calculus. *Mathe. Sb.* **4**, 287–308 (1938)
8. Boem, F., Bonzio, S., Osimani, B.: A logic for scientific attitude? (2020) (**submitted**)
9. Bonzio, S.: Dualities for Płonka sums. *Log. Univers.* **12**(3), 327–339 (2018)
10. Bonzio, S., Gil-Férez, J., Paoli, F., Peruzzi, L.: On paraconsistent weak kleene logic: axiomatization and algebraic analysis. *Stud. Log.* **105**(2), 253–297 (2017)
11. Bonzio, S., Loi, A., Peruzzi, L.: A duality for involutive bisemilattices. *Stud. Log.* **107**(2), 423–444 (2019)
12. Bonzio, S., Pra Baldi, M.: Containment logics: algebraic completeness and axiomatization (2020) (**submitted**)
13. Bonzio, S., Pra Baldi, M., Valota, D.: Counting finite linearly ordered involutive bisemilattices. In: Desharnais, J., Guttman, W., Joosten, S. (eds.) *Relational and Algebraic Methods in Computer Science*, pp. 166–183. Springer, Berlin (2018)
14. Burris, S., Sankappanavar, H.P.: *A Course in Universal Algebra. The Millennium Edition* (2012)
15. Campercholi, M.A., Rafferty, J.G.: Relative congruence formulas and decompositions in quasivarieties. *Algebra Univers.* **78**(3), 407–425 (2017)
16. Campercholi, M.A., Vaggione, D.J.: Implicit definition of the quaternary discriminator. *Algebra Univers.* **68**(1), 1–16 (2012)
17. Chagrov, A., Zakharyashev, M.: *Modal Logic*. Oxford University Press, Oxford (1997)
18. Cignoli, R., D'Ottaviano, I.M.L., Mundici, D.: *Algebraic Foundations of Many-Valued Reasoning*. Kluwer Academic Publishers, Berlin (2000)
19. Cintula, P., Hájek, P., Noguera, C. (eds.): *Handbook of Mathematical Fuzzy Logic. Volumes 1 and 2. Studies in Logic. Mathematical Logic and Foundations*. College Publications, Berlin (2011)
20. Ciuni, R., Carrara, M.: Characterizing logical consequence in paraconsistent weak Kleene. In: Felline, L., Ledda, A., Paoli, F., Rossanese, E. (eds.) *New Directions in Logic and the Philosophy of Science*, pp. 165–176. College Publications, Berlin (2016)
21. Ciuni, R., Ferguson, T.M., Szmuc, D.: Logics based on linear orders of contaminating values. *J. Log. Comput.* **29**(5), 631–663 (2019)
22. Czelakowski, J.: *Protoalgebraic Logics*. Kluwer Academic Publishers, Berlin (2001)
23. Da Ré, B., Pailos, F., Szmuc, D.: Theories of truth based on four-valued infectious logics. *Log. J. IGPL* **53**, 251–252 (2018)
24. Denecke, K., Wismath, S.L.: *Universal Algebra and Applications in Theoretical Computer Science*. Chapman and Hall, London (2002)
25. Diego, A.: *Sobre álgebras de Hilbert. Notas de Lógica Matemática, vol. 12. Universidad Nacional del Sur, Bahía Blanca (Argentina)* (1965)
26. Esteva, F., Godo, L.: Monoidal t-norm based logic: towards a logic for left-continuous t-norms. *Fuzzy Sets Syst.* **124**, 271–288 (2001)
27. Ferguson, T.: A computational interpretation of conceptivism. *J. Appl. Nonclass. Log.* **24**(4), 333–367 (2014)
28. Font, J.M.: *Abstract Algebraic Logic: An Introductory Textbook*. College Publications, Berlin (2016)
29. Font, J.M., Jansana, R.: *A General Algebraic Semantics for Sentential Logics. A.S.L., New York* (2009)
30. Font, J.M., Jansana, R., Pigozzi, D.: A survey on abstract algebraic logic. *Stud. Log. Spec. Issue Abstr. Algebraic Log. Part II* **74**(1–2), 13–97 (2003)
31. Galatos, N., Jipsen, P., Kowalski, T., Ono, H.: *Residuated Lattices: An Algebraic Glimpse at Substructural Logics*. Elsevier, Amsterdam (2007)
32. Gierz, G., Romanowska, A.: Duality for distributive bisemilattices. *J. Aust. Math. Soc. A* **51**, 247–275 (1991)
33. Halldén, S.: *The Logic of Nonsense*. Uppsala Universitets Årsskrift (1949)
34. Harding, J., Romanowska, A.: Varieties of Birkhoff systems: part I. *Order* **34**(1), 45–68 (2017)
35. Harding, J., Romanowska, A.: Varieties of Birkhoff systems: part ii. *Order* **34**(1), 69–89 (2017)
36. Jansana, R., Moraschini, T.: The poset of all logics I: interpretations and lattice structure (2019) (**submitted**)
37. Jansana, R., Moraschini, T.: The poset of all logics II: Leibniz classes and hierarchy (2019) (**submitted**)
38. Jansana, R., Moraschini, T.: The poset of all logics III: finitely presentable logics. *Stud. Log.* (2020)
39. Kalman, J.: Subdirect decomposition of distributive quasilattices. *Fundam. Math.* **2**(71), 161–163 (1971)
40. Kleene, S.: *Introduction to Metamathematics*. North Holland, Amsterdam (1952)

41. Kollár, J.: Congruences and one-element subalgebras. *Algebra Univers.* **9**, 266–267 (1979)
42. Kracht, M.: *Tools and Techniques in Modal Logic*. North-Holland Publishing Co., Amsterdam (1999)
43. Kracht, M.: Modal Consequence Relations, Chapter 8 of the *Handbook of Modal Logic*. Elsevier, Berlin (2006)
44. Lakser, H., Padmanabhan, R., Platt, C.R.: Subdirect decomposition of Płonka sums. *Duke Math. J.* **39**, 485–488 (1972)
45. Lávička, T.: An abstract study of completeness in infinitary logics. Ph.D. Thesis, Charles University (2018)
46. Ledda, A.: Stone-type representations and dualities for varieties of bisemilattices. *Stud. Log.* **106**(2), 417–448 (2018)
47. Libkin, L.: Aspects of partial information in databases. Ph.D. Thesis, University of Pennsylvania (1994)
48. McKenzie, R. N., McNulty, G. F., Taylor, W. F.: *Algebras, Lattices, Varieties*, vol. I. Wadsworth & Brooks/Cole Advanced Books & Software, Berlin (1987)
49. Paoli, F.: *Substructural Logics: A Primer*. Trends in Logic-Studia Logica Library, vol. 13. Kluwer Academic Publishers, Dordrecht (2002)
50. Peruzzi, L.: Algebraic approach to paraconsistent weak Kleene logic. Ph.D. Thesis, University of Cagliari (2018)
51. Płonka, J.: On a method of construction of abstract algebras. *Fundam. Math.* **61**(2), 183–189 (1967)
52. Płonka, J.: On distributive quasilattices. *Fundam. Math.* **60**, 191–200 (1967)
53. Płonka, J., Romanowska, A.: Semilattice sums. In: Romanowska, A., Smith, J. (eds.) *Universal Algebra and Quasigroup Theory*, pp. 123–158. Heldermann, New York (1992)
54. Priest, G.: The logic of paradox. *J. Philos. Log.* **8**, 219–241 (1979)
55. Prior, A.: *Time and Modality*. Oxford University Press, Oxford (1957)
56. Puhlmann, H.: The snack powerdomain for database semantics. In: Borzyszkowski, A.M., Sokołowski, S. (eds.) *Mathematical Foundations of Computer Science 1993*, pp. 650–659. Springer, Berlin (1993)
57. Raftery, J.G.: A perspective on the algebra of logic. *Quaest. Math.* **34**, 275–325 (2011)
58. Raftery, J.G.: Inconsistency lemmas in algebraic logic. *Math. Log. Quart.* **59**(6), 393–406 (2013)
59. Romanowska, A., Smith, J.: *Modes*. World Scientific, Singapore (2002)
60. Romanowska, A., Smith, J.D.: Duality for semilattice representations. *J. Pure Appl. Algebra* **115**(3), 289–308 (1997)
61. Szmuc, D.: Defining LFIs and LFUs in extensions of infectious logics. *J. Appl. Nonclass. Log.* **26**(4), 286–314 (2016)
62. Urquhart, A.: Basic many-valued logic. In: Gabbay, D.M., Guenther, F. (eds.) *Handbook of Philosophical Logic*, vol. 2, pp. 249–295. Springer, Berlin (2001)
63. Wójcicki, R.: *Theory of Logical Calculi*. Basic Theory of Consequence Operations. Reidel, Dordrecht (1988)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.