

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

International Journal of Approximate Reasoning

www.elsevier.com/locate/ijar

Probability over Płonka sums of Boolean algebras: States, metrics and topology

Stefano Bonzio ^{a,*}, Andrea Loi ^b^a Artificial Intelligence Research Institute, Spanish National Research Council, Spain^b Department of Mathematics and Computer Science, University of Cagliari, Italy

ARTICLE INFO

Article history:

Received 24 July 2020

Received in revised form 8 May 2021

Accepted 11 May 2021

Available online 26 May 2021

Keywords:

Involutive bisemilattice

Płonka sum

Boolean algebra

Probability measure

Pseudometric

ABSTRACT

The paper introduces the notion of state for involutive bisemilattices, a variety which plays the role of algebraic counterpart of weak Kleene logics and whose elements are represented as Płonka sums of Boolean algebras. We investigate the relations between states over an involutive bisemilattice and probability measures over the (Boolean) algebras in the Płonka sum representation and, the direct limit of these algebras. Moreover, we study the metric completion of involutive bisemilattices, as pseudometric spaces, and the topology induced by the pseudometric.

© 2021 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Probability theory is grounded on the notion of *event*. Events are traditionally interpreted as elements of a (σ -complete) Boolean algebra. Intrinsically, this means that classical propositional logic is the most suitable formal language to speak about events. The direct consequence of this standard assumption is that any event can either happen (to be the case) or not happen, and, thus, its negation is taking place. One could claim that this criterion does not encompass all situations: certain events might not be either true or false (simply happen or not happen). Think about the coin toss to decide which one, among two tennis players, is choosing whether to serve or respond at the beginning of a match. Although being statistically extremely rare, the coin may fall on the edge, instead of on one face. Pragmatically, the issue is solved re-tossing the coin (hoping to have it landing on one face). Theoretically, one should admit that there are circumstances in which the event “head” (and so also its logical negation “tail”) could be indeterminate (or, undefined). Interestingly enough, not all kinds of events are well modelled by classical logic. This is the case, for instance, of assertions about the properties of quantum systems (which motivated Von Neumann to introduce quantum logic [7], see also [18]), or assertions that can be neither true nor false (like paradoxes), or also propositions regarding vague, or fuzzy properties (like “being tall” or “being smart”). Yet, we are convinced that adopting (some) non-classical logical formalism to describe certain situations is not a good objection to renounce to measure their probability. On the contrary, we endorse the idea of those who think that it constitutes a good reason to look beyond classical probability, namely to render probability when events under consideration do not belong to classical propositional logic.

The idea of studying probability maps over algebraic structures connected to non-classical logic formalisms is at the heart of the theory of *states*. A theory that is well developed for different structures involved in the study of fuzzy logics,

* Corresponding author.

E-mail addresses: stefano@iia.csic.es (S. Bonzio), loi@unica.it (A. Loi).

including MV-algebras [39,24]), Gödel-Dummett [4], Gödel_Δ [1], the logic of nilpotent minimum [3] and product logic [23]. Within the same strand of research, probability maps have been defined and studied also for other algebraic structures (connected to logic), such as Heyting algebras [51], De Morgan algebras [42], orthomodular lattices [5] and effect algebras [25].

The idea motivating the present work is to further extend the theory of states to non-classical events; in particular, to the three-valued logics in the weak Kleene family, whose algebraic semantics is played by the variety of involutive bisemilattices (see [9] and [14]). The peculiarity of such variety is that each of its members has a representation in terms of Płonka sums of Boolean algebras. This abstract construction, originally introduced in universal algebra by J. Płonka [47,48], is performed over direct systems of algebras whose index set is a semilattice. The axiomatisation of states we propose, which is mainly motivated by the logic PWK (Paraconsistent Weak Kleene) but can be easily adapted to Bochvar logic, allows to “break” a state into a family of (finitely additive) probability measures over the Boolean algebras in the Płonka sum representation of an involutive bisemilattice. In other words, our notion of state accounts for (and is strictly connected to) all the Boolean algebras in the Płonka sum. Moreover, we show that states over an involutive bisemilattice are in bijective correspondence with finitely additive probability measures over the Boolean algebra constructed as the direct limit of the algebras in the (semilattice) direct system of the representation. This allows to prove that each state corresponds to an integral over the dual space of the direct limit (the inverse limit of the dual spaces).

The results obtained explore, on the one hand, the possibility of defining probability measures over the algebraic counterpart of certain Kleene logics. On the other, it shows how (finitely additive) probability measures can be lifted from Boolean algebras to the Płonka sum of Boolean algebras.

The paper is organised as follows. Section 2 recaps all the necessary preliminary notions helpful for the reader to go through the entire paper. In Section 3, states over involutive bisemilattices are introduced. We show that each (non-trivial) involutive bisemilattice, having no trivial algebra in its Płonka sum representation, carries at least a state. This class has been introduced (see [44]) for its logical relevance, as algebraic counterpart of an extension of PWK. Moreover, we show the correspondence with probability measures of the direct limit of the Boolean algebras in the semilattice system of **B**. We dedicate Section 4 to the analysis of strictly positive states, which we refer to as *faithful states* (in accordance with the nomenclature for MV-algebras). In particular, the presence of a faithful state motivates the introduction of the subclass of *injective involutive bisemilattices*, characterised by injective homomorphisms in the Płonka sum representation. In Section 5 and 6, respectively, we approach involutive bisemilattices carrying a state as pseudometric spaces and topological spaces (with the topology induced by the pseudometric), respectively. In the former we study metric completions, while in the latter we insist on the relation between involutive bisemilattices and the correspondent direct limits, as topological spaces. We close the paper with Section 7, introducing possible further works and two Appendixes, discussing, respectively, an alternative notion of state (and motivating why we discard it) and some details of an unsolved problem stated in Section 4.

2. Preliminaries

2.1. Płonka sums

A *semilattice* is an algebra $\mathbf{I} = \langle I, \vee \rangle$ of type $\langle 2 \rangle$, where \vee is a binary commutative, associative and idempotent operation. Given a semilattice \mathbf{I} , it is possible to define a partial order relation between the elements of its universe, as follows

$$i \leq j \iff i \vee j = j,$$

for each $i, j \in I$. We say that a semilattice has a *least element*, if there exists an element $i_0 \in I$ such that $i_0 \leq i$ (equivalently, $i_0 \vee i = i$), for all $i \in I$.

Definition 1. A *semilattice direct system of algebras* is a triple $\mathbb{A} = \langle \{\mathbf{A}_i\}_{i \in I}, I, p_{ij} \rangle$ consisting of

- (1) a semilattice¹ $I = \langle I, \vee \rangle$ with least element i_0 ;
- (2) a family of algebras $\{\mathbf{A}_i\}_{i \in I}$ of the same type with disjoint universes;
- (3) a homomorphism $p_{ij} : \mathbf{A}_i \rightarrow \mathbf{A}_j$, for every $i, j \in I$ such that $i \leq j$,

where \leq is the order induced by the binary operation \vee .

Moreover, p_{ii} is the identity map for every $i \in I$, and if $i \leq j \leq k$, then $p_{ik} = p_{jk} \circ p_{ij}$.

Organising a family of algebras $\{\mathbf{A}_i\}_{i \in I}$ into a semilattice direct system means, substantially, requiring that the index set I forms a semilattice and that algebras whose indexes are comparable with respect to the order are “connected” by homomorphisms, whose “direction” is bottom up, namely from algebras whose index is lower to algebras with a greater index. The nomenclature in Definition 1 is deliberately chosen to emphasise the presence of an index set equipped with the

¹ With a slight abuse of notation, we identify semilattice \mathbf{I} with its universe I .

structure of *semilattice*.² We will often refer to a semilattice direct system simply as $\{\mathbf{A}_i\}_{i \in I}$ (instead of $\mathbb{A} = \langle \{\mathbf{A}_i\}_{i \in I}, I, p_{ij} \rangle$) and, in order to indicate homomorphisms, we sometimes write $p_{i,j}$ instead of p_{ij} (in situations where confusion may arise).

The Płonka sum is a new algebra that is defined given a semilattice direct systems of algebras.

Definition 2. Let $\mathbb{A} = \langle \{\mathbf{A}_i\}_{i \in I}, I, p_{ij} \rangle$ be a semilattice direct system of algebras of type τ . The Płonka sum over \mathbb{A} , in symbols $\mathcal{P}_I(\mathbb{A})$ or $\mathcal{P}_I(\mathbf{A}_i)_{i \in I}$, is the algebra such that

- (1) the universe of $\mathcal{P}_I(\mathbb{A})$ is the disjoint union $\bigsqcup_{i \in I} A_i$;
- (2) for every n -ary basic operation f (with $n \geq 1$) in τ , and $a_1, \dots, a_n \in \bigcup_{i \in I} A_i$, we set

$$f^{\mathcal{P}_I(\mathbf{A}_i)_{i \in I}}(a_1, \dots, a_n) := f^{\mathbf{A}_j}(p_{i_1 j}(a_1), \dots, p_{i_n j}(a_n))$$

where $a_1 \in A_{i_1}, \dots, a_n \in A_{i_n}$ and $j = i_1 \vee \dots \vee i_n$;

for every 0-ary operation c (constant) in τ ,

$$c^{\mathcal{P}_I(\mathbf{A}_i)_{i \in I}} := c^{\mathbf{A}_{i_0}},$$

where i_0 is the least element in I .

In words, non-nullary operations $f^{\mathcal{P}_I(\mathbb{A})}$ on the elements a_1, \dots, a_n of the Płonka sum are defined by computing the operation f in the algebra \mathbf{A}_j , whose index is the join of the indexes (a notion that is well defined since the index set I is a semilattice) corresponding to the algebras where the elements a_1, \dots, a_n live, respectively (this idea is clarified through Example 5). The constants of the Płonka sum coincide with the constants of the algebra \mathbf{A}_{i_0} (whose index set is the least element in I).

The theory of Płonka sums is intrinsically connected with the notion of partition function which we recall in the following.

Definition 3. Let \mathbf{A} be an algebra of type τ . A function $\cdot : A^2 \rightarrow A$ is a *partition function* in \mathbf{A} if the following conditions are satisfied for all $a, b, c \in A$, $a_1, \dots, a_n \in A^n$ and for any operation $g \in \tau$ of arity $n \geq 1$, and $c \in \tau$ of arity $n = 0$.

- (PF1) $a \cdot a = a$,
- (PF2) $a \cdot (b \cdot c) = (a \cdot b) \cdot c$,
- (PF3) $a \cdot (b \cdot c) = a \cdot (c \cdot b)$,
- (PF4) $g(a_1, \dots, a_n) \cdot b = g(a_1 \cdot b, \dots, a_n \cdot b)$,
- (PF5) $b \cdot g(a_1, \dots, a_n) = b \cdot a_1 \dots \cdot a_n$,
- (PF6) $a \cdot c = a$.

A comment on the above definition is in order. The upshot of (PF1)-(PF3) is that (A, \cdot) is a *left normal band*. Left normal bands (see e.g. [45, Ch. 2] or [30, Sec. 4.4-4.6]) are important classes of algebras in semigroup theory. (PF4)-(PF6) demand some compatibility between \cdot and the operations in the type τ (which may include constants). It is useful to recall that the above definition uses the minimal number of conditions, in case the type τ admits also constants³; it is not difficult to check that the above introduced definition is equivalent to the one in [49] (see also [50, Section 6]).

The connection between partition functions and Płonka sums is provided by the following result, which states that algebras possessing a (term-definable) partition function can be decomposed as Płonka sums.

Theorem 4. [49, Thm. II] Let \mathbf{A} be an algebra of type τ with a partition function \cdot . The following conditions hold:

- (1) A can be partitioned into $\{A_i\}_{i \in I}$ where any two elements $a, b \in A$ belong to the same component A_i exactly when

$$a = a \cdot b \text{ and } b = b \cdot a.$$

- (2) The relation \leq on I given by the rule

$$i \leq j \iff \text{there exist } a \in A_i, b \in A_j \text{ s.t. } b \cdot a = b$$

is a partial order and $\langle I, \leq \rangle$ is a semilattice with least element.⁴

² Systems of this kind are special cases of *direct systems* (of algebras), which differentiate with respect to the index set which is assumed to be a directed preorder.


³ The construction of the Płonka sum and the notion of partition function were originally introduced by Płonka in [48,47] considering only types without constants and, subsequently, extended to types with constants [49].

⁴ With a slight abuse of notation we indicate here the semilattice of indexes I by the order \leq and not by the binary operation \vee .


- (3) For all $i, j \in I$ such that $i \leq j$ and $b \in A_j$, the map $p_{ij}: \mathbf{A}_i \rightarrow \mathbf{A}_j$, defined by the rule $p_{ij}(x) = x \cdot b$ is a homomorphism.
- (4) $\mathbb{A} = \langle \{\mathbf{A}_i\}_{i \in I}, \langle I, \leq \rangle, \{p_{ij} : i \leq j\} \rangle$ is a direct system of algebras such that $\mathcal{P}_I(\mathbb{A}) = \mathbf{A}$.

Theorem 4 is enunciated in the general form which allows the presence of constants in the type τ . The following example may be useful for the reader to understand how the construction of Płonka sum is performed over a semilattice direct system of Boolean algebras (and the relation with the partition function).

Example 5. Consider the four-elements semilattice $I = \{i_0, i, j, k\}$ whose order is given as follows:



Consider the family $\{\mathbf{A}_{i_0}, \mathbf{A}_i, \mathbf{A}_j, \mathbf{A}_k\}$ of Boolean algebras, organised into a semilattice direct system, whose index is I and homomorphisms are defined as extensions of the following maps: $p_{ik}(a) = c, p_{jk}(b) = e$ (thus, $p_{ik}(a') = c', p_{jk}(b') = e'$), p_{i_0m} is defined in the unique obvious way, for any $m \in \{i, j, k\}$.



According to Definition 2, the Płonka sum over the above introduced semilattice direct system is the new algebra \mathbf{B} , whose universe is $B = A_{i_0} \sqcup A_i \sqcup A_j \sqcup A_k$. The constants in \mathbf{B} are the constants of the Boolean algebra \mathbf{A}_{i_0} . We just give an example of how binary operations are computed in \mathbf{B} (as should be clear that $'$ coincides with negation in each respective algebra).

$$a \wedge^{\mathbf{B}} a' = p_{ii}(a) \wedge^{\mathbf{A}_i} p_{ii}(a') = a \wedge^{\mathbf{A}_i} a' = 0_i.$$

In words, a binary operation between elements belonging to the same algebra (e.g. \mathbf{A}_i) is computed as in that algebra. As for binary operations between elements “living” in (the universes of) two different algebras (e.g. $\mathbf{A}_i, \mathbf{A}_j$) it is necessary to recur to the algebra \mathbf{A}_k (since $k = i \vee j$) and homomorphisms in the system:

$$a \vee^{\mathbf{B}} b = p_{ik}(a) \vee^{\mathbf{A}_k} p_{jk}(b) = c \vee^{\mathbf{A}_k} e = d'.$$

The algebra \mathbf{B} just introduced is an example of involutive bisemilattice (see Definition 6) and may be helpful to clarify also the notion of partition function and the content of Theorem 4.⁵ Observe that \mathbf{B} does not satisfy absorption. Indeed, $a \wedge^{\mathbf{B}} (a \vee^{\mathbf{B}} b) = p_{ik}(a) \wedge^{\mathbf{A}_k} (p_{ik}(a) \vee^{\mathbf{A}_k} p_{jk}(b)) = c \neq a$. However, all the algebras $\{\mathbf{A}_{i_0}, \mathbf{A}_i, \mathbf{A}_j, \mathbf{A}_k\}$ do satisfy absorption (they

⁵ We thank an anonymous reviewer for urging the need for a clarification.

are Boolean algebras). It is indeed easy to verify that the term function $x \cdot y := x \wedge (x \vee y)$ is a partition function on \mathbf{B} (it does satisfy all the properties in Definition 3). In order to clarify the content of Theorem 4, consider $\mathbf{B} = \langle B, \vee, \wedge, ', 0, 1 \rangle$ as an algebra of type $(2, 2, 1, 0, 0)$, and the term operation $x \cdot y := x \wedge (x \vee y)$ playing the role of partition function. Then \mathbf{B} can be partitioned into (four) components each of which satisfies $x \cdot y \approx x$, namely absorption: these components clearly are $\mathbf{A}_{i_0}, \mathbf{A}_i, \mathbf{A}_j, \mathbf{A}_k$. Moreover, the set of indexes $\{i_0, i, j, k\}$ can be equipped with the structure of semilattice, according to condition (2) in Theorem 4. For instance, $i_0 \leq i$ as $1_i \cdot 1 = 1_i \wedge^{\mathbf{B}} (1_i \vee^{\mathbf{B}} 1) = 1_i$; $i \leq k$ since $1_k \cdot 1_i = 1_k \wedge^{\mathbf{B}} (1_k \vee^{\mathbf{B}} 1_i) = 1_k$. Condition (3) guarantees the presence of homomorphisms between the (Boolean) algebras $\{\mathbf{A}_{i_0}, \mathbf{A}_i, \mathbf{A}_j, \mathbf{A}_k\}$. For instance, $p_{ik} : \mathbf{A}_i \rightarrow \mathbf{A}_k$ is defined by the rule $p_{ik}(x) = x \cdot y$ (for some $y \in \mathbf{A}_k$). For instance, $p_{ik}(a) = a \cdot 1_k = c \wedge (c \vee 1_k) = c$. It is not difficult to check that the definition of p_{ik} does not depend on the choice of a specific element in the algebra \mathbf{A}_k .

2.2. Involutive bisemilattices


Definition 6. An involutive bisemilattice is an algebra $\mathbf{B} = \langle B, \wedge, \vee, ', 0, 1 \rangle$ of type $(2, 2, 1, 0, 0)$ satisfying:

- 11. $x \vee x \approx x$;
- 12. $x \vee y \approx y \vee x$;
- 13. $x \vee (y \vee z) \approx (x \vee y) \vee z$;
- 14. $(x')' \approx x$;
- 15. $x \wedge y \approx (x' \vee y')'$;
- 16. $x \wedge (x' \vee y) \approx x \wedge y$;
- 17. $0 \vee x \approx x$;
- 18. $1 \approx 0'$.

It follows from the definition, that the \vee -reduct of an involutive bisemilattice is a (join) semilattice and, consequently, the binary operation \vee induces a partial order \leq_{\vee} . Similarly, in virtue of 15, also the \wedge -reduct is a (meet) semilattice, inducing a partial order \leq_{\wedge} . In general, the two orders are different⁶ and are not lattice-orders: they are (and actually coincide) in case an involutive bisemilattice is a Boolean algebra. The class of involutive bisemilattices forms a variety which we denote by \mathcal{IBSL} , which satisfies all (and only) the regular identities⁷ holding for Boolean algebras. Thus, for instance, it fails to satisfy absorption (see [9] for details). The importance of involutive bisemilattice is connected to logic, as (one of its subquasivarieties) plays the role of algebraic semantics of Paraconsistent Weak Kleene logic (see [9]).

Involutive bisemilattices are connected to Płonka sums. In particular, every involutive bisemilattice is the Płonka sum over a semilattice direct system of Boolean algebras and, conversely, the Płonka sum over any semilattice direct system of Boolean algebras is an involutive bisemilattice (see [9,50]). It follows that the Płonka sum introduced in Example 5 is an example of involutive bisemilattice. Some details of the representation theorem are exemplified in the following (for further details, we refer the reader to [9]).

Example 7. Let $\mathbf{B} \in \mathcal{IBSL}$, with $B = \{0, 1, a, a', b, b'\}$, whose order \leq_{\vee} is represented in the following Hasse diagram.



Define $x \cdot y := x \wedge (x \vee y)$. Observe that \mathbf{B} does not satisfy absorption, indeed: $a \cdot b = a \wedge (a \vee b) = a \wedge b = (a' \vee b')' = (b')' = b$. It is not difficult to check that \cdot is a partition function on \mathbf{B} . By Theorem 4-(1), B can be partitioned in two components: $\mathbf{A}_{i_0} = \{0, a, a', 1\}$, $\mathbf{A}_j = \{b, b'\}$. Indeed (just to show one case), $a \cdot a' = a \wedge (a \vee a') = a \wedge 1 = (a' \vee 0)' = (a')' = a$ (and, similarly, $a' \cdot a = a'$). Condition (2) in Theorem 4 implies that the semilattice of indexes is indeed the two-elements chain $\{i_0, j\}$ with $i_0 < j$. Indeed, for instance, $b \cdot a = b \wedge (b \vee a) = b \wedge b = b$. Condition (3) fixes the (unique) homomorphism $p_{i_0j} : \mathbf{A}_{i_0} \rightarrow \mathbf{A}_j$. To exemplify just one case: $p_{i_0j}(a) = a \cdot b = b$ (it is immediate to check that $p_{i_0j}(1) = b$, $p_{i_0j}(0) = p_{i_0j}(a') = b'$). The elements b and b' are the top and bottom element, respectively, of the Boolean algebra \mathbf{A}_j . This illustrates the Płonka sum representation of \mathbf{B} .

⁶ It is easy to check that $x \leq_{\vee} y$ if and only if $y' \leq_{\wedge} x'$ (see [9]).
⁷ An identity $\varphi \approx \psi$ is regular provided that $Var(\varphi) = Var(\psi)$.

Remark 8. Throughout the whole paper, when considering $\mathbf{B} \in \mathcal{IBSL}$ we will always identify it with its Płonka sum representation $\mathcal{P}_I(\mathbf{A}_i)$, without explicitly mentioning it.

Notation. We denote by $1_i, 0_i$ the top and bottom element, respectively, of the Boolean algebras \mathbf{A}_i (in the Płonka sum representation of $\mathbf{B} \in \mathcal{IBSL}$).

2.3. Finitely additive probability measures over Boolean algebras

Let \mathbf{A} be a Boolean algebra. A finitely additive probability measure over \mathbf{A} is a real-valued map $m: \mathbf{A} \rightarrow [0, 1]$ such that:

- (1) $m(1) = 1$;
- (2) $m(a \vee b) = m(a) + m(b)$, if $a \wedge b = 0$.

With a slight abuse of notation here, and elsewhere, we use the same symbol (“1”) to denote both the top element of a Boolean algebra and the unit of \mathbb{R} . A probability measure m over a Boolean algebra \mathbf{A} is called *regular* (or, strictly positive) provided that $m(a) > 0$ for any $a \neq 0$.

For our purposes, it is useful to recall a relevant result which connects probability maps over Boolean algebras to probability measures over an appropriate topological space. This result (proved, independently, by Kroupa [35] and Panti [43]), holds, more in general, for MV-algebras.⁸ For the convenience of our reader, we opt to formulate it for Boolean algebras, recalling that every Boolean algebra is a semisimple MV-algebra (where the operations \oplus and \odot coincide with \vee and \wedge , respectively). The fact that a Boolean algebra \mathbf{A} is semisimple, as MV-algebra, allows to represent it as an algebra of $[0, 1]$ -valued continuous functions defined over its (dual) Stone space \mathbf{A}^* (see [24, Theorem 2.1.7]). It follows that one can associate to each element $a \in A$, an unique continuous function $a^*: \mathbf{A}^* \rightarrow [0, 1]$. Moreover, we refer to the space of all the (finitely additive) probability measures over a Boolean algebra \mathbf{A} as $\mathcal{S}(\mathbf{A})$.

Theorem 9. Let \mathbf{A} be a Boolean algebra and $m: \mathbf{A} \rightarrow [0, 1]$ a (finitely additive) probability measure. Then

- (1) There is a homeomorphism $\Psi: \mathcal{S}(\mathbf{A}) \rightarrow \mathcal{M}(\mathbf{A}^*)$, where $\mathcal{M}(\mathbf{A}^*)$ is the space of all regular⁹ Borel probability measures on its dual space \mathbf{A}^* .
- (2) For every $a \in A$,

$$m(a) = \int_{\mathbf{A}^*} a^*(M) d\mu_s(M),$$

where a^* is the unique function associated to a , $M \in \mathbf{A}^*$ and $d\mu_s = \Psi(s)$.

A standard reference for the above theorem is [24, Theorem 4.0.1] (where it is stated and proved for MV-algebras).

2.4. Booleanisation of an involutive bisemilattice

Given a semilattice direct system of algebras, the Płonka sum is only one way to construct a new algebra. Another is the *direct limit*. We recall this construction in the special case of Boolean algebras (we refer the reader interested into the details to [28]).

Consider an involutive bisemilattice $\mathbf{B} \cong \mathcal{P}_I(\mathbf{A}_i)$. The *direct limit* over a direct system $\{\mathbf{A}_i\}_{i \in I}$ of Boolean algebras is the Boolean algebra defined as the quotient:

$$\lim_{\rightarrow i \in I} \mathbf{A}_i := (\bigsqcup_{i \in I} \mathbf{A}_i) / \sim,$$

where, for $a \in A_i$ and $b \in A_j$, $a \sim b$ if and only if there exist $c \in A_k$, for some $k \in I$ with $i, j \leq k$, such that $p_{ik}(a) = c = p_{jk}(b)$. It is immediate to check that \sim is a congruence on the (disjoint) union, thus, operations on the direct limit are defined as usual for quotient algebras.

It is always possible to associate to $\mathbf{B} \in \mathcal{IBSL}$, the Boolean algebra $\lim_{\rightarrow} \mathbf{A}_i$, the direct limit of the algebras (in the system) $\{\mathbf{A}_i\}_{i \in I}$, which we will call the *Booleanisation*¹⁰ of \mathbf{B} , and we will indicate it by \mathbf{A}_∞ . Given an involutive bisemilattice \mathbf{B} , we can define the map $\pi: \mathbf{B} \rightarrow \mathbf{A}_\infty$, as $\pi(a) := [a]_\sim$, for any $a \in B$.

⁸ We opted not to recall basic facts about MV-algebras as they are unnecessary to go through the paper. Standard references are [17] or [21].

⁹ Notice that this use of the term “regular” is different from above and refers to Borel measures over topological spaces (see, for instance, [24]).

¹⁰ It is useful to recall that the present use of the term “Booleanisation” differs from other usages in literature, in lattice theory (see [16]) and in the theory of Boolean (inverse) semigroups (see [36]).

Remark 10. The map π is a surjective homomorphism. We provide the details of one binary operation only (the meet \wedge). Suppose $a, b \in B$ with $a \in A_i, b \in A_j$ and $k = i \vee j$, then:

$$\pi(a \wedge b) = [a \wedge b]_{\sim} = [p_{ik}(a) \wedge p_{jk}(b)]_{\sim} = [p_{ik}(a)]_{\sim} \wedge [p_{jk}(b)]_{\sim} = \pi(a) \wedge \pi(b),$$

where the second last equality is justified by observing that $c \in [p_{ik}(a) \wedge p_{jk}(b)]_{\sim}$ (for an arbitrary $c \in A_l$ and $m = k \vee l$) if and only if $p_{lm}(c) = p_{km}(p_{ik}(a) \wedge p_{jk}(b)) = p_{km}(p_{ik}(a)) \wedge p_{km}(p_{jk}(b)) = p_{im}(a) \wedge p_{jm}(b)$ and this is equivalent to say $c \in [p_{ik}(a)]_{\sim} \wedge [p_{jk}(b)]_{\sim}$.

Notation. To indicate elements of the Booleanisation we sometimes drop the subscript \sim when no danger of confusion may arise.

3. States over involutive bisemilattices

Definition 11. Let \mathbf{B} be an involutive bisemilattice. A state over \mathbf{B} is a map $s: \mathbf{B} \rightarrow [0, 1]$ such that:

- (1) $s(1) = 1$;
- (2) $s(a \vee b) = s(a) + s(b)$, provided that $a \wedge b \in \bigcup_{i \in I} \{0_i\}$.

Moreover, a state $s: \mathbf{B} \rightarrow [0, 1]$ is *faithful* if $s(a) > 0$, for every $a \neq 0$.

The following resumes the basic properties of a state over an involutive bisemilattice.

Proposition 12. Let s be a state over an involutive bisemilattice \mathbf{B} . Then

- (1) $s(0) = 0$;
- (2) $s(1_i) = 1$ and $s(0_i) = 0$, for every $i \in I$.
- (3) $s(a') = 1 - s(a)$, for every $a \in B$.

Proof. (1) Since $0 \wedge 1 = 0 \in \bigcup_{i \in I} \{0_i\}$, then $s(1) = s(0 \vee 1) = s(0) + s(1)$, hence $s(0) = 0$.

(2) Observe that $0_i \wedge 1 = 0_i \wedge p_{i0_i}(1) = 0_i \wedge 1_i = 0_i$. Then $s(0_i) + s(1) = s(0_i \vee 1) = s(0_i \vee 1_i) = s(0_i) + s(1_i)$. Therefore $s(1_i) = s(1) = 1$. $s(0_i) = 0$ follows observing that $0_i \vee 1_i = 1_i$.

(3) Let $a \in A_i$ for some $i \in I$. Then $a' \in A_i$ and $a \wedge a' = 0_i$. Therefore $1 = s(1_i) = s(a \vee a') = s(a) + s(a')$. ■

The motivating idea for introducing states over involutive bisemilattices is that of having maps expressing the probability of events that are different from classical ones (corresponding to elements of a Boolean algebra). Our definition of states relies on the algebraic characterisation of the logic PWK given in [9]. We try to motivate the logical reason behind our choice, looking at the involutive bisemilattice introduced in Example 5, which will refer to as \mathbf{B} (in this paragraph). According to that analysis carried out in [9], the only logical filter turning matrix $\langle \mathbf{B}, F \rangle$ into a reduced model of PWK is $F = \{1_{i_0}, 1_i, 1_j, 1_k\}$, i.e. the union of the top elements of the Boolean algebras in the Płonka sum representation of \mathbf{B} (this can be deduced also from the results in [12]). In other words, $F = \{1_{i_0}, 1_i, 1_j, 1_k\}$ expresses the notion of truth with respect to the logic PWK (on the algebra \mathbf{B}). Requiring a state s to map the constant 1 (of an involutive bisemilattice) into 1 (Condition (1) in Definition 11) is equivalent to say that true events (according to the logic PWK) have probability 1 (this is a consequence also of the choice of Condition (2), since it follows from Proposition 12). Condition (2) is introduced in analogy with the classical condition according to which the probability of logically incompatible events is additive. In classical logic, two events x and y are incompatible when their conjunction is the impossible event ($x \wedge y = 0$, in a Boolean algebra). Shifting from classical logic to PWK implies adopting a different notion of incompatibility between events. Two different ways of rendering incompatibility appear then “natural” for an involutive bisemilattice: 1) to say that events x and y are incompatible if $x \wedge y = 0$ or; 2) that events x and y are incompatible if $x \wedge y = 0_i$, where 0_i is the bottom element of the Boolean algebra (in the Płonka sum) where $x \wedge y$ is computed. We opted for 2) persuaded by the idea that two events in the logic PWK are *not compatible* when their conjunction (\wedge in an involutive bisemilattice) is not simply untrue (not belonging to a logical filter), but, more specifically, the *opposite* of truth. And this is obviously obtained choosing the union of zeros in the Boolean algebra in the Płonka sum representation. The choice of the other mentioned notion of incompatibility leads to a different scenario, briefly discussed in Appendix A.

Definition 13. Let $\mathbf{B} \in \mathcal{IBSL}$. For every $i, j \in I$ and a family of finitely additive probability measures $\{m_i\}_{i \in I}$ over $\{\mathbf{A}_i\}_{i \in I}$ (each \mathbf{A}_i carries the measure m_i), the homomorphism p_{ij} preserves the measures if $m_j(p_{ij}(a)) = m_i(a)$, for any $a \in A_i$.

Notation. We indicate the restriction of a state s on the Boolean components $\mathbf{A}_i, \mathbf{A}_j, \mathbf{A}_k, \dots$ of the Płonka sum representation of an involutive bisemilattice \mathbf{B} , as s_i, s_j, s_k, \dots instead of $s_{|_{A_i}}, s_{|_{A_j}}, s_{|_{A_k}}, \dots$ to make notation less cumbersome.

Proposition 14. Let s be a map from \mathbf{B} to $[0, 1]$. The following are equivalent:

- (1) s is a state over \mathbf{B} ;
- (2) $s_i: \mathbf{A}_i \rightarrow [0, 1]$ is a (finitely additive) probability measure over \mathbf{A}_i , for every $i \in I$, and p_{ij} preserves the measures, for each $i \leq j$.

Proof. (1) \Rightarrow (2). Assume that s is a state over \mathbf{B} . Then, $s_i(1_i) = s(1_i) = 1$ for each $i \in I$, by Proposition 12-(2). Moreover, let $a, b \in A_i$ be two elements such that $a \wedge^{A_i} b = 0_i$. Observe that $a \wedge^{\mathbf{B}} b = a \wedge^{A_i} b$ and $a \vee^{\mathbf{B}} b = a \vee^{A_i} b$. Then $s_i(a \vee b) = s(a \vee b) = s(a) + s(b)$, using (2) in Definition 11. This shows that s_i is a (finitely additive) probability measure over \mathbf{A}_i , for every $i \in I$. Now, let $a \in A_i$, for some $i \in I$, and $j \in I$ with $i \leq j$. Observe that $a \wedge^{\mathbf{B}} 0_j = p_{ij}(a) \wedge^{A_j} 0_j = 0_j$, thus

$$s_j(p_{ij}(a)) = s(p_{ij}(a)) = s(p_{ij}(a) \vee^{A_j} 0_j) = s(a \vee^{\mathbf{B}} 0_j) = s(a) + s(0_j) = s(a) = s_i(a),$$

where we have used the additivity of a state and Proposition 12-(1). This shows that any homomorphism p_{ij} preserves the measures s_i, s_j , for every $i \leq j$.

(2) \Rightarrow (1) Assume that every Boolean algebra \mathbf{A}_i in the direct system carries a finitely additive probability measure s_i and that each homomorphism p_{ij} ($i \leq j$) preserves the measures. Let $s: \mathbf{B} \rightarrow [0, 1]$ be the map defined as

$$s(x) := s_i(x),$$

for $x \in A_i$, with $i \in I$.

(1) $s(1) = s_{i_0}(1) = 1$.

(2) Let $a, b \in B$ such that $a \wedge b \in \bigcup_{i \in I} \{0_i\}$. W.l.o.g. let $a \in A_i$ and $b \in A_j$, then $a \wedge^{\mathbf{B}} b = p_{ik}(a) \wedge^{A_k} p_{jk}(b) = 0_k$, with $k = i \vee j$. Consequently, $s(a \vee^{\mathbf{B}} b) = s(p_{ik}(a) \vee^{A_k} p_{jk}(b)) = s_k(p_{ik}(a) \vee^{A_k} p_{jk}(b)) = s_k(p_{ik}(a)) + s_k(p_{jk}(b)) = s_i(a) + s_j(b) = s(a) + s(b)$, where we have essentially used (finite) additivity of probability measures. ■

Example 15. Consider the involutive bisemilattice \mathbf{B} introduced in Example 5 and define the map $s: \mathbf{B} \rightarrow [0, 1]$ as follows:

$$\begin{aligned} s(1) &= s(1_l) = 1, \text{ for any } l \in \{i, j, k\}, \\ s(0) &= s(0_l) = 0, \text{ for any } l \in \{i, j, k\}, \\ s(a) &= s(a') = \frac{1}{2}, \\ s(b) &= \frac{1}{3}, \quad s(b') = \frac{2}{3}, \\ s(c) &= s(c') = \frac{1}{2}, \quad s(d) = \frac{1}{6}, \quad s(e) = \frac{1}{3}, \quad s(d') = \frac{5}{6}, \quad s(e') = \frac{2}{3}. \end{aligned}$$

It is not difficult to check that s is a (faithful) state over \mathbf{B} . Indeed, it is immediate to see that the restrictions of s to the Boolean components of the Płonka sum are finitely additive probability measures which are, moreover, preserved by homomorphisms of the Płonka sum. ■

Observe that, in Proposition 14, it is not necessary to assume the existence of a state over the involutive bisemilattice \mathbf{B} (or, over all the Boolean algebras in the Płonka sum). However, it already gives a gist on which involutive bisemilattices actually carry a state: those whose Płonka sum representation contains no trivial Boolean algebra, as the trivial Boolean algebra carries no (finitely additive) probability measure (this is shown in Corollary 17).

Theorem 16. Let $\mathbf{B} \in \mathcal{IBSL}$. There exists a bijection $\Phi: \mathcal{S}(\mathbf{B}) \rightarrow \mathcal{S}(\mathbf{A}_\infty)$ such that Φ is state preserving, i.e. $s(b) = \Phi(s)(\pi(b))$.

Proof. Consider $\Phi: \mathcal{S}(\mathbf{B}) \rightarrow \mathcal{S}(\mathbf{A}_\infty)$ defined as follows:

$$s(b) \mapsto \Phi(s) = m_\infty([b]_\sim) := s(b), \tag{1}$$

for every $s \in \mathcal{S}(\mathbf{B})$ and $b \in B$, a representative of the equivalence class $[b]_\sim$. Observe that the definition of Φ does not depend on the choice of the representative of $[a]_\sim$. Indeed, let $a, b \in B$ with $a \neq b$, such that $a, b \in [a]_\sim$. W.l.o.g. assume that $a \in A_i$ and $b \in A_j$, for some $i, j \in I$. Since $a \sim b$, then there exists $k \in I$ with $i, j \leq k$ such that $p_{ik}(a) = p_{jk}(b)$. Then, by Proposition 14, $s(a) = s_i(a) = s_k(p_{ik}(a)) = s_k(p_{jk}(b)) = s_j(b) = s(b)$.

Let us prove that $\Phi(s) = m_\infty$ is indeed a (finitely additive) probability measure over the Boolean algebra \mathbf{A}_∞ . To this end, observe that $\bigcup_{i \in I} 1_i \subseteq [1]_\sim$ and choose 1_j (for some $j \in I$) as representative for $[1]_\sim$. Then $m_\infty([1]_\sim) = s(1_j) = 1$, by Proposition 12. Let $[a]_\sim, [b]_\sim \in \mathbf{A}_\infty$ two elements such that $[a]_\sim \wedge [b]_\sim = [0]_\sim$. W.l.o.g. we can assume that $a \in A_i, b \in A_j$, for some $i, j \in I$, with $i \neq j$, and set $k = i \vee j$. The assumption that $[a]_\sim \wedge [b]_\sim = [0]_\sim$ implies that there exists $l \geq i, j$ such that $p_{il}(a) \wedge p_{jl}(b) = 0_l$ (and, obviously, $k \leq l$). Then:

$$\begin{aligned}
m_\infty([a]_\sim \vee [b]_\sim) &= m_\infty([a \vee b]_\sim) \\
&= s(a \vee b) \\
&= s_k(p_{ik}(a) \vee^{A_k} p_{jk}(b)) \\
&= s_l(p_{kl}(p_{ik}(a) \vee^{A_k} p_{jk}(b))) \\
&= s_l(p_{il}(a) \vee^{A_k} p_{jl}(b)) \\
&= s_l(p_{il}(a)) + s_l(p_{jl}(b)) && \text{(Proposition 14)} \\
&= s_i(a) + s_j(b) && \text{(Proposition 14)} \\
&= s(a) + s(b) \\
&= m_\infty([a]_\sim) + m_\infty([b]_\sim).
\end{aligned}$$

To complete the proof, let us check that Φ is invertible. Let $m: \mathbf{A}_\infty \rightarrow [0, 1]$ be a (finitely additive) probability measure over \mathbf{A}_∞ . Define $\Phi^{-1}(m[a]_\sim) = s_i(a) := m([a]_\sim)$, for any $[a] \in \mathbf{A}_\infty$, with $a \in A_i$ (for $i \in I$). Let us check that $s_i: \mathbf{A}_i \rightarrow [0, 1]$ is a (finitely additive) probability measure (over \mathbf{A}_i). $m_i(1_i) = m([1_i]_\sim) = m([1]_\sim) = 1$. Moreover, let $a, b \in A_i$ two elements such that $a \wedge b = 0_i$. Then, $[a]_\sim \wedge [b]_\sim = [a \wedge b]_\sim = [0_i]_\sim = [0]_\sim$. Therefore, $s_i(a \vee b) = m([a \vee b]_\sim) = m([a]_\sim \vee [b]_\sim) = m([a]_\sim) + m([b]_\sim) = m_i(a) + m_i(b)$. Observe that, the homomorphism p_{ij} preserves the measures s_i and s_j , for any $i \leq j$. Indeed (for $[a]_\sim \in \mathbf{A}_\infty$) $a \sim p_{ij}(a)$, thus $s_i(a) = m([a]_\sim) = m([p_{ij}(a)]_\sim) = s_j(p_{ij}(a))$. Therefore, since m_i is a finitely additive probability measure over \mathbf{A}_i , for each $i \in I$, and homomorphisms preserve the measures, by Proposition 14, we get that $\Phi^{-1}(m)$ is a state over \mathbf{B} (it is immediate to check that Φ^{-1} is the inverse of Φ). Finally, it follows from the definition that Φ is state preserving. ■

Corollary 17. Let $\mathbf{B} \in \mathcal{IBSL}$ with $\mathcal{P}_I(\mathbf{A}_i)$ its Płonka sum representation. The following are equivalent:

- (1) \mathbf{B} carries a state;
- (2) $\{\mathbf{A}_i\}_{i \in I}$ contains no trivial algebra.

Proof. In virtue of the correspondence established in Theorem 16, \mathbf{B} carries a state if and only if the Booleanisation \mathbf{A}_∞ does carry a (finitely additive) probability measure, i.e. \mathbf{A}_∞ is non trivial (since the trivial Boolean algebra carries no such map). Moreover, observe that \mathbf{A}_∞ is trivial if and only if there is an algebra \mathbf{A}_k (for some $k \in I$) in the system that is trivial. To show the latter claim, suppose that \mathbf{A}_∞ is trivial, i.e. $[0]_\sim = [1]_\sim$, then there is some $k \in I$ such that $p_{ik}(0_i) = p_{jk}(1_j)$ (for some $i, j \in I$), i.e. $0_k = 1_k$, showing that \mathbf{A}_k is the trivial algebra. For the converse direction, suppose that \mathbf{A}_∞ is non-trivial, i.e. $[0]_\sim \neq [1]_\sim$, and, by contradiction, that $\{\mathbf{A}_i\}_{i \in I}$ contains a trivial algebra \mathbf{A}_k (for some $k \in I$), i.e. $0_k = 1_k$, but $0_k \in [0]_\sim$ and $1_k \in [1]_\sim$, a contradiction with $[0]_\sim \neq [1]_\sim$. ■

Interestingly, the class of involutive bisemilattices admitting no trivial algebra in the Płonka sum representation forms a subquasivariety of \mathcal{IBSL} , known as \mathcal{NGIB} , which has been introduced in [44] and plays the role of the algebraic counterpart of the non-paraconsistent extension of the logic PWK by adding the ex-falso quodlibet (this consists of the unique extension of PWK, different from classical logic). \mathcal{NGIB} is axiomatised¹¹ (with respect to \mathcal{IBSL}) by the quasi-identity $x \approx x' \Rightarrow y \approx z$.

From now on, we will consider only (non-trivial) involutive bisemilattices whose Płonka sum representation contains no trivial Boolean algebra, thus the non-trivial elements belonging to the class \mathcal{NGIB} . Hence, in view of Corollary 17, any involutive bisemilattice considered carries at least a state. We indicate by $\mathcal{S}(\mathbf{B})$ and $\mathcal{S}(\mathbf{A}_\infty)$ the spaces of states and of (finitely additive) probability measures of an involutive bisemilattice \mathbf{B} and of its Booleanisation \mathbf{A}_∞ , respectively.

The correspondence established in Theorem 16 allows to provide an integral representation of states over an involutive bisemilattice. Let \mathbf{A}_∞ be the Booleanisation of an involutive bisemilattice \mathbf{B} and \mathbf{A}_∞^* the Stone space dually equivalent to \mathbf{A}_∞ . It is useful to recall that \mathbf{A}_∞^* corresponds to the inverse limit over the direct system whose elements $\{\mathbf{A}_i^*\}_{i \in I}$ are the dual spaces of the Boolean algebras $\{\mathbf{A}_i\}_{i \in I}$ in the representation of \mathbf{B} (see [28]).

Theorem 18 (Integral representation of states). Let $\mathbf{B} \in \mathcal{NGIB}$ and $s: \mathbf{B} \rightarrow [0, 1]$ be a state. Then

- (1) There is a bijection $\chi: \mathcal{S}(\mathbf{B}) \rightarrow \mathcal{M}(\mathbf{A}_\infty^*)$, where $\mathcal{M}(\mathbf{A}_\infty^*)$ is the space of all regular Borel probability measures on \mathbf{A}_∞^* .

¹¹ It shall be mentioned that, in [44], the authors work with generalised involutive bisemilattices, namely defined in the type containing no constant. However, this difference is not significant for the purpose of our discussion.

(2) For every $b \in B$,

$$s(b) = \int_{\mathbf{A}_\infty^*} [\widehat{b}]_{\sim}(M) d\mu_s(M),$$

where $[\widehat{b}]_{\sim}$ is the unique function associated to $[b]_{\sim} \in A_\infty$, $M \in \mathbf{A}_\infty^*$ and $d\mu_s = \chi(s)$.

Proof. It follows from the bijective correspondence between states of an involutive bisemilattices and of its Booleanisation (Theorem 16) and integral representation for probability measures over Boolean algebras (Theorem 9). ■

One may wonder whether it is possible to provide a different integral representation of states which makes use of the dual space of an involutive bisemilattice (for instance, the Płonka product described in [10]) instead of the inverse limits of the dual spaces of the Boolean algebras involved in the Płonka sum representation. This is a question that we do not address in the present paper.

4. Faithful states

Recall that a state s over an involutive bisemilattice \mathbf{B} is faithful (cfr. Definition 11) when $s(a) > 0$, for any $a \neq 0$. By Proposition 14, this is equivalently expressed by saying that $s(a) > 0$, for every $a \notin \{0_i\}_{i \in I}$. A (finitely additive) probability measure m over a Boolean algebra \mathbf{C} is *regular* provided that $m(a) > 0$, when $a \neq 0$.

The presence of a faithful state s over an involutive bisemilattice \mathbf{B} has a non-trivial consequence on the structure of its Płonka sum representation, as expressed in the following.

Proposition 19. *Let s be a state over $\mathbf{B} \in \mathcal{IBSL}$. The following are equivalent:*

- (1) s is faithful;
- (2) $s_i : \mathbf{A}_i \rightarrow [0, 1]$ is a regular (finitely additive) probability measure over \mathbf{A}_i , for every $i \in I$, and p_{ij} is an injective homomorphism preserving the measures, for each $i \leq j$.

Proof. We just show the non-trivial direction (1) \Rightarrow (2). By Proposition 14, we only have to prove that s_i is regular and that p_{ij} is an injective homomorphism, for every $i \leq j$. The former is immediate, indeed for $a \in A_i$, (with $i \in I$) such that $a \neq 0_i$, then $s_i(a) = s(a) > 0$. As for the latter, let $i \leq j$ for some $i, j \in I$. Let $a \in \ker(p_{ij})$, then $p_{ij}(a) = 0_j$ (we are using here the fact that congruences are determined by their 0-class, see for instance [26]). Then, $s_j(p_{ij}(a)) = s(p_{ij}(a)) = s(0_j) = 0$, by Proposition 12. By Proposition 14, p_{ij} preserves the measures (and s_i, s_j are probability measures), hence $s_i(a) = 0$ and, since s_i is regular (as shown above), then $a = 0_i$. This shows that $\ker(p_{ij}) = \{0_i\}$, i.e. p_{ij} is injective. ■

Definition 20. Let $\mathbf{B} \in \mathcal{IBSL}$. We say that \mathbf{B} is *injective* if, for every $i, j \in I$ such that $i \leq j$, the homomorphism $p_{ij} : \mathbf{A}_i \rightarrow \mathbf{A}_j$ is injective.

We refer to the class of injective involutive bisemilattices, that has also been introduced in [44], as \mathcal{IGIB} . It is not difficult to see that \mathcal{IGIB} is closed under subalgebras and products but not under homomorphic images.

Recall that the variety of involutive bisemilattices admits a partition function \cdot , that can be defined as $x \cdot y := x \wedge (x \vee y)$ (see Examples 5 and 7).

Proposition 21. *Let $\mathbf{B} \in \mathcal{IBSL}$ with partition function \cdot . The following are equivalent:*

- (1) $\mathbf{B} \in \mathcal{IGIB}$;
- (2) $\mathbf{B} \models x \cdot y \approx x \ \& \ y \cdot x \approx y \ \& \ x \cdot z \approx y \cdot z \Rightarrow x \approx y$.

Proof. (1) \Rightarrow (2) Suppose $\mathbf{B} \in \mathcal{IGIB}$, i.e. p_{ij} is an embedding for each $i \leq j$. Suppose, in view of a contradiction, that \mathbf{B} does not satisfy condition (2). Then, there exist elements $a, b, c \in B$ such that $a \cdot b = a$ and $b \cdot a = b$ and $a \cdot c = b \cdot c$ but $a \neq b$. By Theorem 4-(1), $a \cdot b = a$ and $b \cdot a = b$ imply that $a, b \in A_i$, for some $i \in I$. W.l.o.g. assume that $c \in A_j$, for some $j \in I$ and set $k = i \vee j$. Then, applying Theorem 4 and the assumption that $a \cdot c = b \cdot c$, we have $p_{ik}(a) = p_{ik}(a) \cdot p_{jk}(c) = a \cdot c = b \cdot c = p_{ik}(b) \cdot p_{jk}(c) = p_{ik}(b)$. Since $\mathbf{B} \in \mathcal{IGIB}$, p_{ik} is injective, namely $a = b$, in contradiction with our hypothesis.

(2) \Rightarrow (1) Suppose \mathbf{B} satisfies condition (2) and, by contradiction, that $\mathbf{B} \notin \mathcal{IGIB}$, namely there exists a homomorphism p_{ij} (for some $i \leq j$) which is not an embedding. Hence, there exists element $a \neq b$ such that $p_{ij}(a) = p_{ij}(b)$. Clearly $a, b \in A_i$ (otherwise p_{ij} is not well defined), so, by Theorem 4-(1), $a \cdot b = a$ and $b \cdot a = b$. Let $c = p_{ij}(a) = p_{ij}(b)$, then $a \cdot c = p_{ij}(a) \cdot p_{ij}(a) = p_{ij}(a) = p_{ij}(b) = p_{ij}(b) \cdot p_{ij}(b) = b \cdot c$. Then, by condition (2), we have that $a = b$, a contradiction. ■

It follows from the above Proposition that \mathcal{IGIB} is a quasi-variety, which can be axiomatised relatively to \mathcal{IBSL} by the quasi-identity $x \cdot y \approx x \& y \cdot x \approx y \& x \cdot z \approx y \cdot z \Rightarrow x \approx y$ (a different quasi-equational axiomatisation can be found in [44]).

Theorem 22. *Let $\mathbf{B} \in \mathcal{IGIB}$. Then there is a bijective correspondence between faithful states over \mathbf{B} and regular measures over \mathbf{A}_∞ .*

Proof. The correspondence is given by the map Φ , defined in (1). Indeed, let $\mathbf{B} \in \mathcal{IGIB}$ and $s: \mathbf{B} \rightarrow [0, 1]$ a faithful state. Assume that $[a]_\sim \in \mathbf{A}_\infty$ with $[a]_\sim \neq [0]_\sim$. Then $a \neq 0$, hence $s(a) > 0$ and $m_\infty([a]_\sim) = s(a) > 0$, which shows that $\Phi(s)$ is regular. For the other direction, it is immediate to check that, given a regular measure $m: \mathbf{A}_\infty \rightarrow [0, 1]$ then $s_i(a) = \Phi^{-1}(m([a]_\sim))$ (for $a \in A_i$) is also a regular measure and since \mathbf{B} is injective, Proposition 19 guarantees that $\Phi^{-1}(m)$ is a faithful state over \mathbf{B} . ■

Combining [24, Proposition 3.1.7] and Theorem 16 we directly get the following.

Corollary 23. *The space $\mathcal{S}(\mathbf{B})$ of states of an involutive bisemilattice \mathbf{B} can be identified (via Φ) with a non-empty compact subspace of $[0, 1]^{\mathbf{A}_\infty}$.*


It is natural to wonder under which conditions an involutive bisemilattice \mathbf{B} (whose Płonka sum representation contains no trivial algebra) carries a faithful state. Theorem 22 suggests that this might be the case provided that \mathbf{B} is injective and its Booleanisation \mathbf{A}_∞ actually carries a regular probability measure. In general, as observed in [33], not every (non-trivial) Boolean algebra carries a regular probability measure (necessary and sufficient conditions for a Boolean algebra to carry a regular measure are stated in [33, Theorem 4]).

It is possible to define the categories of injective involutive bisemilattices and Boolean algebras “with faithful state”, “with regular probability measure”, respectively. Objects are pairs (\mathbf{B}, s) and (\mathbf{A}, m) , where $\mathbf{B} \in \mathcal{IBSL}$, \mathbf{A} is a Boolean algebra, s is a state over \mathbf{B} and m is a (finitely additive) probability measure over \mathbf{A} . A morphism between two objects (\mathbf{B}_1, s_1) and (\mathbf{B}_2, s_2) is a homomorphism (between the corresponding algebras in the first component) which preserves the measures, namely $s_1(b) = s_2(h(b))$, for every $b \in B_1$ and every homomorphism $h: \mathbf{B}_1 \rightarrow \mathbf{B}_2$. It is immediate to check that involutive bisemilattices carrying a faithful state (Boolean algebras carrying a probability measure, resp.) form a category, which we indicate by the pair $(\mathcal{IBSL}, \mathcal{S}(\mathbf{B}))$ ($\mathcal{BA}, \mathcal{S}(\mathbf{A})$), resp.

Let us define a functor $\mathcal{F}: (\mathcal{IBSL}, \mathcal{S}(\mathbf{B})) \rightarrow (\mathcal{BA}, \mathcal{S}(\mathbf{A}))$, which associates, to each object (\mathbf{B}, s) , the object $\overline{\mathcal{F}}(\mathbf{B}, s) = (\mathbf{A}_\infty, \Phi(s))$, with Φ defined as in (1) and, to each morphism $h: (\mathbf{B}_1, s_1) \rightarrow (\mathbf{B}_2, s_2)$, the morphism $\overline{\mathcal{F}}(h) = \overline{h}$ with $\overline{h}: \mathbf{A}_{1_\infty} \rightarrow \mathbf{A}_{2_\infty}$, defined as $\overline{h}[a_1] = [h(a_1)]$.

Theorem 24. *\mathcal{F} is a covariant functor between the categories of injective involutive bisemilattices with faithful states and Boolean algebras with regular probability measures.*

Proof. Let (\mathbf{B}, s) an object in $(\mathcal{IBSL}, \mathcal{S}(\mathbf{B}))$. Then, by Theorem 16, $\overline{\mathcal{F}}(\mathbf{B}, s) = (\mathbf{A}_\infty, \Phi(s))$ is an object in $(\mathcal{BA}, \mathcal{S}(\mathbf{A}))$. We have to prove that, for every homomorphism $h: (\mathbf{B}_1, s_1) \rightarrow (\mathbf{B}_2, s_2)$ preserving states, the map $\overline{h}: (\mathbf{A}_{1_\infty}, \Phi(s_1)) \rightarrow (\mathbf{A}_{2_\infty}, \Phi(s_2))$ is a Boolean homomorphism, which preserves the (finitely additive) probability measures. To see that \overline{h} is indeed a Boolean homomorphism, we show just one case (all the others are proved analogously). Let $[a], [b] \in \mathbf{A}_{1_\infty}$, then $\overline{h}([a] \wedge [b]) = \overline{h}([a \wedge b]) = [h(a \wedge b)] = [h(a) \wedge h(b)] = [h(a)] \wedge [h(b)] = \overline{h}(a) \wedge \overline{h}(b)$. Moreover, \overline{h} preserves the states. Indeed let $[a_1] \in \mathbf{A}_{1_\infty}$ and $\Phi(s_1) \in \mathcal{S}(\mathbf{A}_{1_\infty})$. Then $\Phi(s_1)([a_1]) = s_1(a_1) = s_2(h(a_1)) = \Phi(s_2)([h(a_1)]) = \Phi(s_2)(\overline{h}[a_1])$, where we have used the fact that h preserves states. Finally, it follows from the definition of \overline{h} that the following diagram is commutative and this concludes the proof of our claim.



The functor \mathcal{F} admits many adjoints, depending on the number of injective involutive bisemilattices having the same Booleanisation.

Problem. Characterise all the injective involutive bisemilattices having the same Booleanisation.

The problem is not of easy solution. Taking up the suggestion of an anonymous reviewer, we have drawn some considerations about it in Appendix B.

5. States, metrics and completion

Definition 25. Let X be a set. A *pseudometric* on X is a map $d: X \times X \rightarrow \mathbb{R}$ such that:

- (1) $d(x, y) \geq 0$,
- (2) $d(x, x) = 0$,
- (3) $d(x, y) = d(y, x)$,
- (4) $d(x, z) \leq d(x, y) + d(y, z)$ (triangle inequality),

for all $x, y, z \in X$.

A pair (X, d) given by a set with a pseudometric d is called *pseudometric space*. A pseudometric d , on X , is a *metric*, if $d(x, y) = 0$ implies that $x = y$. Notice that, as for metric spaces, the triangle inequality is enough to show that the pseudo-metric d is continuous (a fact that we will use in several proofs), when X is topologised with the topology induced by the pseudo-metric (with no explicit mention).

Let us recall that in any Boolean algebra \mathbf{A} it is possible to define the *symmetric difference* $\Delta: A \times A \rightarrow A$, $a \Delta b := (a \wedge b') \vee (a' \wedge b)$. The presence of a (finitely additive) probability measure m over a Boolean algebra \mathbf{A} allows to define a pseudo-metric $d := m \circ \Delta$ on \mathbf{A} , which becomes a metric, in case m is regular (see [34] for details).

The symmetric difference can be (analogously) defined also for an involutive bisemilattice \mathbf{B} . Let $a, b \in B$, with $\mathbf{B} \cong \mathcal{P}_I(\mathbf{A}_i)$, $a \in A_i$ and $b \in A_j$, for some $i, j \in I$. Then $a \Delta b = (a \wedge^{\mathbf{B}} b') \vee^{\mathbf{B}} (a' \wedge^{\mathbf{B}} b) = p_{ik}(a) \Delta^{\mathbf{A}_k} p_{jk}(b)$, where $k = i \vee j$. Clearly, in case \mathbf{B} carries a state s , then one can define the map $d_s: \mathbf{B} \rightarrow [0, 1]$ as:

$$d_s := s \circ \Delta. \tag{2}$$

We will sometimes refer to the (pseudo)metrics induced by states and finitely additive probability measures as distances.

Proposition 26. Let \mathbf{B} be an involutive bisemilattice carrying a state s . Then d_s is a pseudo-metric on B .

Proof. (1) obviously holds, since $s: \mathbf{B} \rightarrow [0, 1]$.

(2) Let $a \in A_i$, for some $i \in I$. Then $d_s(a, a) = s((a \wedge a') \vee (a' \wedge a)) = s(0_i \vee 0_i) = s(0_i) = 0$, by Proposition 12.

(3) holds since \vee and \wedge are commutative operations.

(4) Let $a \in A_i$, $b \in A_j$ and $c \in A_k$ with $i \neq j \neq k$. Preliminarily, observe that

$$\begin{aligned} d_s(a, c) &= s((a \wedge^{\mathbf{B}} c') \vee^{\mathbf{B}} (a' \wedge^{\mathbf{B}} c)) \\ &= s((p_{i,i \vee k}(a) \wedge^{\mathbf{A}_{i \vee k}} p_{k,i \vee k}(c')) \vee^{\mathbf{A}_{i \vee k}} (p_{i,i \vee k}(a)' \wedge^{\mathbf{A}_{i \vee k}} p_{k,i \vee k}(c))) \\ &= s_{i \vee k}((p_{i,i \vee k}(a) \wedge^{\mathbf{A}_{i \vee k}} p_{k,i \vee k}(c')) \vee^{\mathbf{A}_{i \vee k}} (p_{i,i \vee k}(a)' \wedge^{\mathbf{A}_{i \vee k}} p_{k,i \vee k}(c))) \\ &= s_{i \vee k \vee j}(p_{i \vee k, i \vee k \vee j}((p_{i,i \vee k}(a) \wedge^{\mathbf{A}_{i \vee k}} p_{k,i \vee k}(c')) \vee^{\mathbf{A}_{i \vee k}} (p_{i,i \vee k}(a)' \wedge^{\mathbf{A}_{i \vee k}} p_{k,i \vee k}(c)))) \quad (\text{Proposition 14}) \\ &= s_{i \vee k \vee j}(a \Delta^{\mathbf{A}_{i \vee k \vee j}} c). \end{aligned}$$

Similarly, it is possible to show that $d_s(a, b) = s_{i \vee k \vee j}(a \Delta^{\mathbf{A}_{i \vee k \vee j}} b)$ and $d_s(b, c) = s_{i \vee k \vee j}(b \Delta^{\mathbf{A}_{i \vee k \vee j}} c)$. Hence $d_s(a, c) = s_{i \vee k \vee j}(a \Delta^{\mathbf{A}_{i \vee k \vee j}} c) \leq s_{i \vee k \vee j}(a \Delta^{\mathbf{A}_{i \vee k \vee j}} b) + s_{i \vee k \vee j}(b \Delta^{\mathbf{A}_{i \vee k \vee j}} c) = s \circ d(a, b) + s \circ d(b, c) = d_s(a, b) + d_s(b, c)$. ■

Remark 27. Recall that, for Boolean algebras (and, more in general, MV-algebras [37]), the choice of a regular probability measure m (or, a faithful state in the case of MV-algebras) implies that the induced pseudometric d_m is indeed a metric. This fact does not hold for involutive bisemilattices, where the choice of a faithful state s is not enough to guarantee that d_s is a metric. Indeed, consider any $\mathbf{B} \in \mathcal{NGLB}$ and two distinct elements $a, b \in B$, $a \in A_i$, $b \in B_j$, such that $p_{i,i \vee j}(a) = p_{j,i \vee j}(b)$. Then $d_s(a, b) = s(a \Delta^{\mathbf{B}} b) = s(p_{i,i \vee j}(a) \Delta^{\mathbf{A}_{i \vee j}} p_{j,i \vee j}(b)) = s(0_{i \vee j}) = 0$; however, $a \neq b$. Notice, moreover, that $a \sim b$, i.e. they belong to the same equivalence class in the Booleanisation \mathbf{A}_∞ . Indeed, it is easy to check that d_s is a metric over \mathbf{B} if and only if $\mathbf{B} = \mathbf{A}_\infty$.

We now recall a well known fact about Boolean algebras.

Lemma 28. Let \mathbf{A} be a Boolean algebra carrying a finitely additive probability measure m and let d_m be the pseudometric ($d_m = m \circ \Delta$) defined via m . Then:

- (1) $d_m(a, b) = d_m(a', b')$;
- (2) $d_m(a \vee b, c \vee d) \leq d_m(a, c) + d_m(b, d)$,

for any $a, b, c, d \in A$.

Proof. The proof can be adapted from [37, Lemma 3.1], observing that Boolean algebras are MV-algebras where the operations \oplus and \vee coincide. ■

We decide to give explicit proofs of all the following results in the particular case of pseudometric (injective) involutive bisemilattices (and relative Booleanisations), although some of them could be derived from the general theory of pseudo-metric spaces (see for instance [32]).

In analogy to what is done in [37] for MV-algebras, in the remaining part of this section we study the metric completion for involutive bisemilattices. The completion is a standard construction for metric spaces (see [32]), which can be analogously applied to pseudo-metric spaces.

Recall that a (pseudo)metric space is *complete* if every Cauchy sequence is convergent. Given a pseudometric space (X, d) , a completion $(\widehat{X}, \widehat{d})$ of (X, d) is such that:

- (1) $(\widehat{X}, \widehat{d})$ is complete;
- (2) there exists an isometric injective map $j: X \rightarrow \widehat{X}$ such that $j(X)$ is dense in \widehat{X} .

Observe that the second condition means that, for every $\widehat{x} \in \widehat{X}$, there exists a sequence $x_n \in X$ such that $j(x_n) \rightarrow \widehat{x}$.

In the sequel, we can assume, up to isometry, that the embedding j is the natural inclusion $X \hookrightarrow \widehat{X}$. Notice that the completion $(\widehat{X}, \widehat{d})$ is uniquely determined, by the above conditions (1)-(2), up to isometries.

We proceed in the same way for involutive bisemilattices. In detail, given (\mathbf{B}, d_s) a pair where \mathbf{B} is an involutive bisemilattice (carrying a state s) and d_s the pseudometric induced by the state s , we can associate to it, on the one hand, the completion $(\widehat{\mathbf{B}}, \widehat{d}_s)$; on the other hand, we can consider its Płonka sum representation $\mathcal{P}_I(\mathbf{A}_i)$. By Proposition 14, for every $i \in I$, (\mathbf{A}_i, d_{s_i}) is a pseudo-metric space (since s_i is a probability measure over \mathbf{A}_i), which is metric in case s is faithful (see Proposition 19). Therefore, it makes sense to consider the pseudo-metric space $(\widetilde{\mathbf{B}}, \widetilde{d})$, where $\widetilde{\mathbf{B}} = \mathcal{P}_I(\widehat{\mathbf{A}}_i)_{i \in I}$ (the Płonka sum of the completions of the Boolean algebras¹² in the system representing \mathbf{B}), $\widetilde{d}(\widetilde{a}, \widetilde{b}) = \lim_{n \rightarrow \infty} d_s(a_n, b_n)$, where $\widetilde{a} \in \widehat{\mathbf{A}}_i, \widetilde{b} \in \widehat{\mathbf{A}}_j$ (for some $i, j \in I$), and a_n, b_n are sequences (of elements) in $\mathbf{A}_i, \mathbf{A}_j$, respectively, such that $a_n \rightarrow \widetilde{a}, b_n \rightarrow \widetilde{b}$. We are going to show (see Theorem 31 below) that $\widetilde{\mathbf{B}} \in \mathcal{IBSL}$ and, moreover, that $(\widetilde{\mathbf{B}}, \widetilde{d})$ is the completion of (\mathbf{B}, d_s) .

Lemma 29. *Let $(\mathbf{A}_1, d_1), (\mathbf{A}_2, d_2)$ be two Boolean algebras with distance (induced by a probability measure) and $h: \mathbf{A}_1 \rightarrow \mathbf{A}_2$ be a distance preserving homomorphism. Then there exists a distance preserving homomorphism $\widehat{h}: \widehat{\mathbf{A}}_1 \rightarrow \widehat{\mathbf{A}}_2$ such that $\widehat{h}|_{\mathbf{A}_1} = h$.*

Proof. Let $\widehat{a} \in \widehat{\mathbf{A}}_1$ and define $\widehat{h}: \widehat{\mathbf{A}}_1 \rightarrow \widehat{\mathbf{A}}_2$ as

$$\widehat{h}(\widehat{a}) := \lim_{n \rightarrow \infty} h(a_n),$$

where $a_n \rightarrow \widehat{a}$ is a sequence of (elements of) \mathbf{A}_1 convergent to \widehat{a} . Observe that \widehat{h} is well defined. Indeed, suppose that $a'_n \rightarrow \widehat{a}$, i.e. a'_n is a different sequence convergent to the element \widehat{a} . Then $\lim_{n \rightarrow \infty} \widehat{d}_2(h(a_n), h(a'_n)) = \lim_{n \rightarrow \infty} d_2(h(a_n), h(a'_n)) = \lim_{n \rightarrow \infty} d_1(a_n, a'_n) = \widehat{d}_1(\widehat{a}, \widehat{a}) = 0$, where the second equality is justified by the fact that h is an isometry (distance preserving map). This shows that $\lim_{n \rightarrow \infty} h(a_n) = \lim_{n \rightarrow \infty} h(a'_n)$. Moreover, it follows by construction that $\widehat{h}|_{\mathbf{A}_1} = h$. It only remains to show that \widehat{h} is distance preserving and a Boolean homomorphism. Let $\widehat{a}, \widehat{b} \in \widehat{\mathbf{A}}_1$. Then

$$\begin{aligned} \widehat{d}_2(\widehat{h}(\widehat{a}), \widehat{h}(\widehat{b})) &= \widehat{d}_2(\lim_{n \rightarrow \infty} h(a_n), \lim_{n \rightarrow \infty} h(b_n)) \\ &= \lim_{n \rightarrow \infty} \widehat{d}_2(h(a_n), h(b_n)) && (\widehat{d}_2 \text{ is continuous}) \\ &= \lim_{n \rightarrow \infty} d_2(h(a_n), h(b_n)) \\ &= \lim_{n \rightarrow \infty} d_1(a_n, b_n) && (h \text{ is isometry}) \\ &= \widehat{d}_1(\widehat{a}, \widehat{b}). \end{aligned}$$

To see that \widehat{h} is a Boolean homomorphism, we only show the cases of negation and one of the two binary operations.

$$\begin{aligned} \widehat{h}(\widehat{a}') &= \widehat{h}(\lim_{n \rightarrow \infty} a'_n) \\ &= \lim_{n \rightarrow \infty} \widehat{h}(a'_n) && (\widehat{h} \text{ is continuous}) \end{aligned}$$

¹² The fact that the completion of a Boolean algebra is still a Boolean algebra is a routine exercise: one may follow the strategy applied to MV-algebras in [37] (using the fact that Boolean algebras are MV-algebras where \oplus is idempotent).

$$\begin{aligned}
 &= \lim_{n \rightarrow \infty} h(a'_n) && (\widehat{h}|_{\mathbf{A}_1} = h) \\
 &= \lim_{n \rightarrow \infty} h(a_n)' && (h \text{ homomorphism}) \\
 &= \widehat{h}(\widehat{a})',
 \end{aligned}$$

where the last equality holds, since, by Lemma 28-(1), $\widehat{d}_2(h(a_n)', \widehat{h}(\widehat{a})') = \widehat{d}_2(h(a_n), \widehat{h}(\widehat{a})) = 0$. We reason similarly for \vee :

$$\begin{aligned}
 \widehat{h}(\widehat{a} \vee \widehat{b}) &= \widehat{h}(\lim_{n \rightarrow \infty} a_n \vee b_n) \\
 &= \lim_{n \rightarrow \infty} \widehat{h}(a_n \vee b_n) && (\widehat{h} \text{ is continuous}) \\
 &= \lim_{n \rightarrow \infty} h(a_n \vee b_n) && (\widehat{h}|_{\mathbf{A}_1} = h) \\
 &= \lim_{n \rightarrow \infty} h(a_n) \vee h(b_n) && (h \text{ homomorphism}) \\
 &= \widehat{h}(\widehat{a}) \vee \widehat{h}(\widehat{b}),
 \end{aligned}$$

where the last equality holds, since, by Lemma 28-(2), $\widehat{d}_2(h(a_n) \vee h(b_n), \widehat{h}(\widehat{a}) \vee \widehat{h}(\widehat{b})) \leq \widehat{d}_2(h(a_n), \widehat{h}(\widehat{a})) + \widehat{d}_2(h(b_n), \widehat{h}(\widehat{b})) = 0$. ■

Notation. Given the pseudo-metric space (\mathbf{B}, d_s) , where $\mathbf{B} \in \mathcal{NGLB}$ and d_s is the pseudo-metric induced by a state s , we indicate by d_∞ (instead of $d_{\Phi(s)}$) the pseudo-metric on its Booleanisation, obtained via the bijection in Theorem 16.

Lemma 30. Let $\mathbf{B} \in \mathcal{IBSL}$ carrying a state s . Then (\mathbf{B}, d_s) is complete if and only if $(\mathbf{A}_\infty, d_\infty)$ is complete.

Proof. Observe that a sequence $\{x_n\}_{n \in \mathbb{N}}$ of elements in \mathbf{B} is a Cauchy sequence if and only if the sequence $\{[x_n]_{\sim}\}_{n \in \mathbb{N}}$ is a Cauchy sequence in \mathbf{A}_∞ . Indeed, if, for each $\varepsilon > 0$, there is a $n_0 \in \mathbb{N}$ such that $d_s(x_n, x_m) < \varepsilon$, for every $n, m > n_0$, then also $d_\infty([x_n]_{\sim}, [x_m]_{\sim}) < \varepsilon$, since $d_s(x_n, x_m) = d_\infty([x_n]_{\sim}, [x_m]_{\sim})$ by Theorem 16. The result follows by observing that $\lim_{n \rightarrow \infty} x_n = x$ if and only if $\lim_{n \rightarrow \infty} [x_n]_{\sim} = [x]_{\sim}$. ■

Theorem 31. Let (\mathbf{B}, d_s) be an involutive bisemilattice with the pseudo-metric d_s induced by a state s , then:

- (1) $\widetilde{\mathbf{B}} = \mathcal{P}_I(\widehat{\mathbf{A}}_I)$ is an involutive bisemilattice;
- (2) $(\widetilde{\mathbf{B}}, \widetilde{d}_s)$ is isometric to $(\widetilde{\mathbf{B}}, \widetilde{d})$.

Proof. (1) $\widetilde{\mathbf{B}} = \bigsqcup_{i \in I} \widehat{\mathbf{A}}_i$, where $\widehat{\mathbf{A}}_i$, for each $i \in I$, is a Boolean algebra (since Boolean algebras are closed under completions). Moreover, by Lemma 29, the system $\langle \{\widehat{\mathbf{A}}_i\}_{i \in I}, I, \{\widehat{h}_{ij}\}_{i \leq j} \rangle$ is a semilattice direct systems of Boolean algebras (given that $\langle \{\mathbf{A}_i\}_{i \in I}, I, \{h_{ij}\}_{i \leq j} \rangle$ is such) and this is enough to conclude that $\widetilde{\mathbf{B}}$ is an involutive bisemilattice.

(2) Preliminarily observe that $\widetilde{\mathbf{A}}_\infty = \widehat{\mathbf{A}}_\infty$ (the Booleanisations of $\widetilde{\mathbf{B}}$ and $\widehat{\mathbf{B}}$ coincide). Then, by Lemma 30, it follows that $\widetilde{\mathbf{B}}$ is complete, as its Booleanisation is complete. We claim that (\mathbf{B}, d_s) is a dense subset of $(\widetilde{\mathbf{B}}, \widetilde{d})$. Indeed, let $\widetilde{b} \in \widetilde{\mathbf{B}}$, then $\widetilde{b} \in \widehat{\mathbf{A}}_i$, for some $i \in I$ (since $\widetilde{\mathbf{B}} = \bigsqcup_{i \in I} \widehat{\mathbf{A}}_i$). Thus, there exists a sequence $\{a_n\}_{n \in \mathbb{N}} \in A_i$ such that $\lim_{n \rightarrow \infty} d_s(a_n, \widetilde{b}) = 0$. Then, by definition of \widetilde{d} , $\widetilde{d}(a_n, \widetilde{b}) \rightarrow 0$, which implies that a_n (as element of \mathbf{B}) converges to \widetilde{b} , i.e. \mathbf{B} is dense in $\widetilde{\mathbf{B}}$. It follows (from the previous claim) that there is an isometric bijection $f : (\widetilde{\mathbf{B}}, \widetilde{d}_s) \rightarrow (\widetilde{\mathbf{B}}, \widetilde{d})$. ■

6. The topology of involutive bisemilattices

An involutive bisemilattice \mathbf{B} carrying a state s can be topologised with the topology induced by the pseudo-metric d_s , defined in (2). In virtue of Theorem 16 (and Theorem 22 for faithful states), also the Booleanisation \mathbf{A}_∞ (of \mathbf{B}) can be topologised via the corresponding probability measure which we indicate as $d_\infty = \Phi(s) \circ \Delta$ (where Φ is the map defined in (1)). In this section, we confine ourselves only to faithful states over (injective) involutive bisemilattices and when referring to \mathbf{B} and \mathbf{A}_∞ as topological spaces, we think them as equipped with the topologies \mathcal{T}_{d_s} and \mathcal{T}_{d_∞} induced by the respective (pseudo) metric. To achieve this we should assume that the involutive bisemilattices under consideration carry faithful states. Recall that, in this topology, a subset $U \subset B$ is open if and only if for every $x \in U$, there exists $r > 0$ such that $D_r(x) \subset U$, where $D_r(x) = \{y \in B \mid d_s(x, y) < r\}$ is the open disk centred in x with radius r . Moreover, one base of both topologies is given by the family of all open disks with respect to d_s and d_∞ , respectively.

Remark 32. Notice that, in an involutive bisemilattice \mathbf{B} , carrying a state s , $s(b) = d_s(b, 0)$, for every $b \in B$. Indeed let $b \in A_i$, for some A_i in the Płonka sum representation of \mathbf{B} , then $d_s(b, 0) = s \circ \Delta^{\mathbf{B}}(b, 0) = s((b \wedge 1_i) \vee (b' \wedge 0_i)) = s(b \vee 0_i) = s(b)$. This implies that $s : \mathbf{B} \rightarrow [0, 1]$ is continuous (since the pseudo-metric is continuous) when \mathbf{B} is topologised via \mathcal{T}_{d_s} .

Definition 33. Let X be a topological space and let $\equiv \subseteq X \times X$ the equivalence relation defined as $x \equiv y$ if and only if x and y have the same open neighbourhoods. Then, the space $X_{/\equiv}$ is the *Kolmogorov quotient* of X .

In words, two points x, y belonging to the same equivalence class with respect to \equiv are *topologically indistinguishable*.

Remark 34. Notice that the topology on $X_{/\equiv}$ is the quotient topology, namely a set U is open in $X_{/\equiv}$ if and only if $\pi^{-1}(U)$ is open in X , where $\pi : X \rightarrow X_{/\equiv}$ is the (natural) projection onto $X_{/\equiv}$.

Proposition 35. Let $\mathbf{B} \in \mathcal{IGTB}$ which carries a faithful state s . Then the Kolmogorov quotient of \mathbf{B} is its Booleanisation \mathbf{A}_∞ .

Proof. We have to show that two elements $a, b \in B$ are topologically indistinguishable (namely, $d_s(a, b) = 0$) if and only if $a \sim b$. Assume w.l.o.g. that $a \in A_i$ and $b \in A_j$ with $i \neq j$ and set $k = i \vee j$.

(\Rightarrow) Let $d_s(a, b) = 0$ (i.e. a, b are indistinguishable). Then $s(a \Delta^{\mathbf{B}} b) = 0$ and, since s is faithful, $a \Delta^{\mathbf{B}} b = p_{ik}(a) \Delta^{\mathbf{A}_k} p_{jk}(b) = 0_k$. Then $p_{ik}(a) = p_{jk}(b)$, i.e. $a \sim b$.

(\Leftarrow) Let $a \sim b$. It follows that there exists $l \in I$ such that $i, j \leq l$ and $p_{il}(a) = p_{jl}(b)$. Then

$$\begin{aligned} s(a \Delta b) &= s(p_{ik}(a) \Delta^{\mathbf{A}_k} p_{jk}(b)) \\ &= s_k(p_{ik}(a) \Delta^{\mathbf{A}_k} p_{jk}(b)) \\ &= s_l(p_{kl}(p_{ik}(a) \Delta^{\mathbf{A}_k} p_{jk}(b))) && \text{(Proposition 14)} \\ &= s_l(p_{kl} \circ p_{ik}(a) \Delta^{\mathbf{A}_l} p_{kl} \circ p_{jk}(b)) \\ &= s_l(p_{il}(a) \Delta^{\mathbf{A}_l} p_{jl}(b)) \\ &= s(0_l) \\ &= 0. \end{aligned}$$

This shows that a, b are indeed topologically indistinguishable points. ■

The proof of Proposition 35 shows another interesting fact worth being highlighted: while \mathbf{B} is a pseudo-metric space, its Booleanisation \mathbf{A}_∞ becomes a *metric* space.

By combining Proposition 35 and Theorem 37 we immediately get the following.

Corollary 36. Let $\mathbf{B} \in \mathcal{IGTB}$ with a faithful state s . Then $\widehat{\mathbf{B}}_{/\equiv} = \widehat{\mathbf{A}}_\infty$, i.e. the Kolmogorov quotient of the completion (\mathbf{B}, d_s) is the completion of the metric space $(\mathbf{A}_\infty, d_\infty)$.

Recall that, given a surjective map $f : X \rightarrow Y$ between topological spaces X and Y , a *section* of f is a continuous map $g : Y \rightarrow X$ such that $f \circ g = id_Y$.

Theorem 37. The following facts hold for the topological spaces \mathbf{B} and \mathbf{A}_∞ :

- (1) There exists a section $\sigma : \mathbf{A}_\infty \rightarrow \mathbf{B}$ of π such that $\sigma(\mathbf{A}_\infty)$ is dense in \mathbf{B} ;
- (2) σ preserves states, namely $s \circ \sigma = \Phi(s)$.

Proof. (1) Consider $\sigma : \mathbf{A}_\infty \rightarrow \mathbf{B}$ defined as follows:

$$\sigma([a]_\sim) = a,$$

where a is picked by a choice function among the elements of the equivalence class $[a]_\sim$ (this can be done recurring to the Axiom of choice). We first show that σ is continuous. Let $D_r(b)$ an open disk of radius r centred in b , for some $b \in B$. $\sigma^{-1}(D_r(b)) = \{[y]_\sim \in \mathbf{A}_\infty \mid \sigma([y]_\sim) \in D_r(b)\} = \{[y]_\sim \in \mathbf{A}_\infty \mid y \in D_r(b)\} = \{[y]_\sim \in \mathbf{A}_\infty \mid d_s(y, b) < r\} = \{[y]_\sim \in \mathbf{A}_\infty \mid d_\infty([b]_\sim, [y]_\sim) < r\} = D_r([b]_\sim)$, the open disk of \mathbf{A}_∞ centred in $[b]_\sim$. This shows the continuity of σ .

The fact that $\pi \circ \sigma = id$ immediately follows from the definition of σ .

To show that $\sigma(\mathbf{A}_\infty)$ is dense in \mathbf{B} , we have to check that $\sigma(\mathbf{A}_\infty) \cap U \neq \emptyset$, for every non-void open set $U \subset B$. Since U is non empty, there exists $b \in U$. Moreover, since U is open, then, for some $r > 0$, $D_r(b) \subset U$ (where $D_r(b) = \{y \in B \mid d_s(b, y) < r\}$). W.l.o.g. we can assume that $b \in A_i$, for some $i \in I$. Observe that, for each $j \in I$ with $i \leq j$, we have that $p_{ij}(b) \in U$. Indeed $d_s(b, p_{ij}(b)) = s(b \Delta^{\mathbf{B}} p_{ij}(b)) = s_j(p_{ij}(b) \Delta^{\mathbf{A}_j} p_{ij}(b)) = 0$, which implies that $p_{ij}(b) \in D_r(b) \subset U$. If $\sigma([b]) = b$, then we have finished. So, assume that $\sigma([b]) = a$, with $a \neq b$. W.l.o.g. let $a \in A_j$ (for some $j \in I$). By definition of σ , $a \in [b]_\sim$, i.e. there exists some $k \in I$, such that $i, j \leq k$ and $p_{ik}(b) = p_{jk}(a)$. Since $p_{ik}(b) \in U$ (for the above observation), then also $p_{jk}(a) \in U$. Reasoning as above, one checks that $d_s(a, p_{jk}(a)) = 0$, which implies that $a \in U$. This shows that $\sigma(\mathbf{A}_\infty) \cap U \neq \emptyset$, for every

non-void open set $U \subset B$, i.e. $\sigma(\mathbf{A}_\infty)$ is dense in \mathbf{B} .
 (2) follows from the definition of σ and Theorem 16. ■

Remark 38. Observe that the projection π admits many sections (depending on the cardinality of the fiber $\pi^{-1}(\{b\})$, for $b \in B$) and all of them are topological embeddings by construction.

Recall that, if $f: X \rightarrow Y$ be a continuous map between two topological spaces (X and Y), $V \subseteq X$ is f -saturated (or saturated with respect to f) if $V = f^{-1}(f(V))$.

Lemma 39. Every open and closed set of an involutive bisemilattice \mathbf{B} is saturated with respect to the projection $\pi: \mathbf{B} \rightarrow \mathbf{A}_\infty$. In particular, π is (continuous) open and closed.

Proof. Since the basis of the topology (over \mathbf{B}) is the family of open disks, then, with respect to open sets, it is enough to check that $\pi^{-1}(\pi(D_r)) = D_r$, for some open disk D_r . Let $D_r(b)$ an open disk (of radius r) centred in b , for some $b \in B$. Then $\pi^{-1}(\pi(D_r(b))) = \{x \in B \mid \pi(x) \in \pi(D_r(b))\} = \{x \in B \mid d_\infty([x], [b]) < r\} = \{x \in B \mid d_s(x, b) < r\} = D_r(b)$, where the second last equality holds by Theorem 16.
 Let $C \subseteq B$ a closed set. Then $C = B \setminus U$, for some open set U . Observe that $C \subseteq \pi^{-1}(\pi(C))$ holds in general, so we have to show only the converse inclusion. To this end $\pi^{-1}(\pi(C)) = \pi^{-1}(\pi(B \setminus U)) \subseteq \pi^{-1}(\pi(B) \setminus \pi(U)) = \pi^{-1}(\pi(B)) \setminus \pi^{-1}(\pi(U)) = B \setminus U = C$, where the second last equality holds since open sets are saturated with respect to π . Finally, the fact that π is open and closed follows from what we have just proved, observing that \mathbf{A}_∞ is the Kolmogorov quotient of \mathbf{B} (Proposition 35), topologised with the quotient topology (see Remark 34). ■

Remark 40. In the proof of the following results we will use some well-known facts in general topology that we briefly recap (see, for instance, [41]). Let $f: X \rightarrow Y$, be an open and closed continuous function between topological spaces. Then

- (1) $f^{-1}(\text{Int}(B)) = \text{Int}(f^{-1}(B))$, for every $B \subseteq Y$;
- (2) $f(\bar{A}) = \overline{f(A)}$, for every $A \subseteq X$.

Lemma 41. Let $C \subseteq B$ a closed set of an involutive bisemilattice \mathbf{B} . Then $\pi(\text{Int}(C)) = \text{Int}(\pi(C))$.

Proof. Let C be a closed set in B . Observe that, from Lemma 39, we have that π is an open and closed continuous map. Hence

$$\pi(\text{Int}(C)) = \pi(\text{Int}(\pi^{-1}(\pi(C)))) = \pi(\pi^{-1}(\text{Int}(\pi(C)))) = \text{Int}(\pi(C)),$$

where we have applied Lemma 39 and the properties of open and closed continuous maps (see Remark 40). ■

Observe that the statement of Lemma 41 does not hold for all topological spaces (see [38] for details). Recall that a map $f: X \rightarrow Y$ (between two topological spaces) preserves the interiors if $\text{Int}(f(A)) = f(\text{Int}(A))$, for all $A \subseteq X$. Interior preserving maps are studied in [38]. One can wonder whether the statement of Lemma 41 could be extended to any subset of involutive bisemilattice (not only for closed subsets). Interestingly enough, the next result shows that the projection π is interior preserving if and only if \mathbf{B} is a Boolean algebra.

Theorem 42. Let \mathbf{B} an involutive bisemilattice carrying a faithful state. The following facts are equivalent:

- (1) $\mathbf{B} = \mathbf{A}_\infty$;
- (2) $\pi: \mathbf{B} \rightarrow \mathbf{A}_\infty$ is an interior preserving map;
- (3) $\sigma(\mathbf{A}_\infty)$ is open (closed, saturated) in \mathbf{B} , for every section $\sigma: \mathbf{A}_\infty \rightarrow \mathbf{B}$.

Proof. (1) \Rightarrow (2) is trivial (as $\pi = id$).
 (2) \Rightarrow (1). We reason by contraposition, and suppose that $\mathbf{B} \neq \mathbf{A}_\infty$. This implies that there exists an element $[a] \in A_\infty$ such that $|\pi^{-1}([a])| \geq 2$. Let $b \in \pi^{-1}([a])$. Observe that $\mathbf{A}_\infty \setminus \{[a]\}$ is open (as $\{[a]\}$ is closed), thus, since π is continuous, $\mathbf{B} \setminus \{\pi^{-1}([a])\}$ is open. This implies that $\text{Int}(\mathbf{B} \setminus \{b\}) = \mathbf{B} \setminus \{\pi^{-1}([a])\}$. Then, $\pi(\text{Int}(\mathbf{B} \setminus \{b\})) = \pi(\mathbf{B} \setminus \{\pi^{-1}([a])\}) = \mathbf{A}_\infty \setminus \{[a]\}$. On the other hand, $\text{Int}(\pi(\mathbf{B} \setminus \{b\})) = \mathbf{A}_\infty$, since $|\pi^{-1}([a])| \geq 2$, which shows that π does not preserve interiors.
 (1) \Rightarrow (3) is obvious.
 (3) \Rightarrow (1). Let $\sigma(\mathbf{A}_\infty)$ be open (closed, saturated) in \mathbf{B} . Then, by Lemma 39, $\sigma(\mathbf{A}_\infty)$ is π -saturated, i.e. $\sigma(\mathbf{A}_\infty) = \pi^{-1}(\pi(\sigma(\mathbf{A}_\infty))) = \pi^{-1}(\mathbf{A}_\infty) = \mathbf{B}$, so π is a bijection being σ its inverse. ■

Recall that, for a topological space X , an open set $U \subseteq X$ is an open regular set if $U = \text{Int}(\bar{U})$ (where \bar{U} indicates the closure of U). To keep in mind the difference between an open, and an open regular set, consider \mathbb{R} topologised (as usual)

with the Euclidian topology. Then $(0, 1)$ is an example of an open regular set, while $U = (0, 1) \cup (1, 2)$ is an open set which is not regular, as $\text{Int}(\overline{U}) = (0, 2)$. The set of open regular sets $\text{Reg}(X)$ of a topological space X can be turned into a (complete) Boolean algebra (see, for instance, [26]) $\mathbf{Reg}(X) = \langle \text{Reg}(X), \cap, \vee, \setminus, \emptyset, X \rangle$, where $U \vee V := \text{Int}(\overline{U \cup V})$. Moreover, the Boolean algebra of **Clopen**(X) (of the clopen sets of X) is a subalgebra of $\mathbf{Reg}(X)$. Despite the fact that an involutive bisemilattice \mathbf{B} and its Booleanisation \mathbf{A}_∞ are not homeomorphic (except in the trivial case $\mathbf{B} = \mathbf{A}_\infty$), surprisingly enough, the Boolean algebras of regular sets arising from \mathbf{B} and \mathbf{A}_∞ are isomorphic, as shown in the following.

Theorem 43. *Let $\mathbf{B} \in \mathcal{IGIB}$ carrying a faithful state. The projection $\pi : \mathbf{B} \rightarrow \mathbf{A}_\infty$ induces a bijection between $\text{Open}(\mathbf{B})$ and $\text{Open}(\mathbf{A}_\infty)$, the open sets of \mathbf{B} and \mathbf{A}_∞ , respectively. Moreover, the Boolean algebras $\mathbf{Reg}(\mathbf{B})$ and $\mathbf{Reg}(\mathbf{A}_\infty)$ are isomorphic.*

Proof. The fact that π is a bijection between $\text{Open}(\mathbf{B})$ and $\text{Open}(\mathbf{A}_\infty)$ follows from Lemma 39 (and the surjectivity of π). The isomorphism between $\mathbf{Reg}(\mathbf{B})$ and $\mathbf{Reg}(\mathbf{A}_\infty)$ is given by the projection π , restricted to $\mathbf{Reg}(\mathbf{B})$. We first show that the map is well defined, i.e. that given an open regular set $U \in \mathbf{Reg}(\mathbf{B})$, then $\pi(U) \in \text{Reg}(\mathbf{A}_\infty)$. To show regularity, observe that

$$\begin{aligned} \pi(U) &= \pi(\text{Int}(\overline{U})) && (U \text{ is regular}) \\ &= \text{Int}(\pi(\overline{U})) && (\text{Lemma 41}) \\ &= \text{Int}(\overline{\pi(U)}) && (\pi \text{ is continuous, open and closed}) \end{aligned}$$

To conclude the proof, we only need to check that π is a homomorphism (with respect to the Boolean operations of $\mathbf{Reg}(\mathbf{B})$ and $\mathbf{Reg}(\mathbf{A}_\infty)$). With respect to the constants, observe that $\pi(\emptyset) = \emptyset$ and, since π is surjective, $\pi(\mathbf{B}) = \mathbf{A}_\infty$. Now, let $U, V \in \text{Reg}(\mathbf{B})$, then $\pi(U) \cap \pi(V) = \pi \circ \pi^{-1}(\pi(U) \cap \pi(V)) = \pi(\pi^{-1}(\pi(U)) \cap \pi^{-1}(\pi(V))) = \pi(U \cap V)$, where the last equality follows from Lemma 39. Moreover,

$$\begin{aligned} \pi(U \vee V) &= \pi(\text{Int}(\overline{U \cup V})) \\ &= \text{Int}(\pi(\overline{U \cup V})) && (\text{Lemma 41}) \\ &= \text{Int}(\overline{\pi(U \cup V)}) && (\pi \text{ is continuous, open and closed}) \\ &= \text{Int}(\overline{\pi(U) \cup \pi(V)}) \\ &= \pi(U) \vee \pi(V). \end{aligned}$$

Since we have shown that π preserves the constants and the binary operations, it follows that it preserves also the unary operation \setminus , hence we are done. ■

Theorem 44 (Topological characterization of states). *Let s be a faithful state over \mathbf{B} and $t : \mathbf{B} \rightarrow [0, 1]$ a continuous map such that $t \circ \sigma = \Phi(s)$, for any section $\sigma : \mathbf{A}_\infty \rightarrow \mathbf{B}$. Then $t = s$.*

Proof. By assumption, $t \circ \sigma = \Phi(s)$, for any section $\sigma : \mathbf{A}_\infty \rightarrow \mathbf{B}$. This implies that the two maps s and t coincide over a dense subset $\sigma(\mathbf{A}_\infty)$ of \mathbf{B} (in virtue of Theorem 37-(2)). Therefore, since both s and t are continuous (s by Remark 32, t by assumption) and $[0, 1]$ is Hausdorff, we have $t = s$. ■

Remark 45. As we have seen, in general, the spaces \mathbf{B} and \mathbf{A}_∞ are not homeomorphic. However, it follows from the general theory of Kolmogorov quotients (see [46, Theorem 8.6]) that, under the assumption that they are both Alexandrov discrete¹³ they are homotopically equivalent. Obviously, the equivalence holds in the particular case whether \mathbf{B} is finite.

7. Conclusion and further work

In this work, we have shown how to define a notion of state on Płonka sums of Boolean algebras, with the aim of expressing the probability for elements of an involutive bisemilattice, a variety associated to the logic PWK. In particular, we have shown that the (non-trivial) elements of the class \mathcal{NGIB} (of involutive bisemilattices with no trivial algebra in the Płonka sum representation) always carry a state. The class \mathcal{NGIB} plays a relevant role in logic, as algebraic counterpart of the extension of PWK by adding the ex-falso quodlibet. Moreover, we have exploited the connections between such notion, the probability measures carried by Boolean algebras in a Płonka sum and the Booleanisation of an involutive bisemilattice. These connections are crucial in the study of the completion and the topology induced by a state over an involutive bisemilattice.

¹³ A topological space X is *Alexandrov discrete* when the arbitrary intersection of open sets is an open set.

This work sheds a further light on the possibility of developing the theory of probability beyond the boundaries of classical logic, i.e. when events are elements of a Boolean algebra. To the best of our knowledge, this consists of the first attempt to lift (finitely additive) probability measures from Boolean algebras to Płonka sums of Boolean algebras. For this reason, many theoretical problems, as well as potential applications are not examined in the present work. At first, it shall be noticed that there is nothing special behind the choice of Boolean algebras, a part the fact that Płonka sums of Boolean algebras play the important role of characterising the algebraic counterparts of weak Kleene logics and its extensions. The ideas developed here could be used, in principle, to define states for varieties that are represented as Płonka sums of classes of algebras admitting states, such as MV-algebras, Goedel algebras, Heyting algebras, just to mention some for which a theory of states has been developed. On the other hand, a deeper investigation about the connection between Płonka sums and certain logics has been conducted in [12], [13] and [14].

A relevant question that we leave for further investigations is the possibility of characterising states over involutive bisemilattices as *coherent books* over a (finite) set of events of the extension of the logic PWK. Coherent books have been introduced, in the classical case, by de Finetti [19,20], via a specific (reversible) betting game and are shown to be in one-to-one correspondence with (finitely additive) probability measures over the Boolean algebra generated by the events considered. This kind of abstract betting scenario has been used also to characterise states for non-classical structures [40].

We have shown (see Theorem 18) that states over involutive bisemilattices correspond to integrals on the dual space of the Booleanisation. It makes sense to ask whether this correspondence can be extended to faithful states, relying on the integral representation proved for faithful states over free MV-algebras in [22].

The theory of states we developed could, perhaps, find potential applications also in the field of *knowledge representation*. This is mainly due to the fact that states break into probability measures over the Boolean algebras in the Płonka sum representation. One may interpret the semilattice of indexes, involved in the representation, to model, for instance, situations of branching time¹⁴ (as the index set is, in general, not a chain). A state, then, encapsulates information related to the probabilities of classical events (Boolean algebras) located in every point (indexes) of the structure. This might be used, in principle, also to analyse conditional bettings or counterfactual situations, under the assumption, for instance, that events are related when there is a homomorphism connecting the algebras they belong to.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

S.B. gratefully acknowledges the financial support of the PRIN project “From models to decisions” (Italian Ministry of Scientific Research grant n. 201743F9YE, Turin unit) and the support of the postdoctoral fellowship Beatriu de Pinos, funded by the Secretary of Universities and Research (Government of Catalonia) and by the Horizon 2020 program of research and innovation of the European Union under the Marie Skłodowska-Curie grant agreement No. 801370. A.L. acknowledges the support of the Kasba, Regione Autonoma della Sardegna Fondo di Sviluppo e Coesione 2014 - 2020 - Interventi di sostegno alla ricerca. Both authors also gratefully acknowledges the INDAM GNSAGA, Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni. We finally thank Roberto Giuntini, Francesco Paoli, Tommaso Flaminio and two anonymous reviewers for their fruitful comments on previous versions of the paper.

Appendix A

Our definition of state relies (see Definition 11) on the assumption that two elements $a, b \in B$ of an involutive bisemilattice \mathbf{B} are logically incompatible provided that $a \wedge^{\mathbf{B}} b = 0_i$, where 0_i is the bottom element of the Boolean algebra (in the Płonka sum representation of \mathbf{B}) where the operation \wedge is computed. One could question this principle and understand two elements $a, b \in B$ as incompatible, in case $a \wedge^{\mathbf{B}} b = 0$. This leads to a different definition of state obtained, by replacing condition (2) in Definition 11 with the following:

$$s(a \vee b) = s(a) + s(b) \text{ provided that } a \wedge b = 0. \quad (3)$$

However, since the element 0 of an involutive bisemilattice always belongs to the Boolean algebra (in the Płonka sum) whose index is the least element in the semilattice (I, \leq) of indexes, this latter choice leads to the following consequence.

Proposition 46. *Let \mathbf{B} and involutive bisemilattice. Then the following are equivalent:*

(1) $s: \mathbf{B} \rightarrow [0, 1]$ satisfies $s(1) = 1$ and condition (3);

¹⁴ A similar idea is developed from the construction of horizontal sums in [2].

(2) s_{i_0} is a (finitely additive) probability measure over the Boolean algebra \mathbf{A}_{i_0} where i_0 is the minimum element in I .

Proof. (1) \Rightarrow (2). Immediate by observing that $1 \in A_{i_0}$ and that, for any arbitrary pair of elements $a, b \in B$, $a \wedge b = 0$ implies that $a, b \in A_{i_0}$.

(2) \Rightarrow (1). Let $s_{i_0} : \mathbf{A}_{i_0} \rightarrow [0, 1]$ be any finitely additive probability measure over \mathbf{A}_{i_0} . Then, the map $s : \mathbf{B} \rightarrow [0, 1]$

$$s(x) := \begin{cases} s_{i_0}(x) & \text{if } x \in A_{i_0}, \\ \alpha & \text{otherwise,} \end{cases}$$

for $\alpha \in (0, 1)$ a fixed number, satisfies that $s(1) = 1$ and that $s(a \vee b) = s(a) + s(b)$, when $a \wedge b = 0$. Moreover, s_{i_0} is the restriction of s over \mathbf{A}_{i_0} . ■

In words, the above result suggests that, this different notion of state, obtained by replacing (2) in Definition 11 with (3), implies that only the elements belonging to the Boolean algebra \mathbf{A}_{i_0} are actually measured following the standard rules of probability.

Appendix B

We have shown that (faithful) states on involutive bisemilattices are in correspondence with (regular) probability measures over the corresponding Booleanisations (see Theorem 16 and Theorem 22). However, there could be many non-isomorphic (injective) involutive bisemilattices having the same Booleanisation.¹⁵ The problem of characterising all injective involutive bisemilattices having the same Booleanisation (outlined at the end of Section 4) is not an easy task and the aim of present appendix is (to try) to provide some reasons.

At first, we observe that a knowledge of the lattice of subalgebras of Boolean algebras (see [27,6]) is not sufficient to give an answer to the problem, even in case the index set I is finite and, consequently, the Booleanisation coincides with the Boolean algebra (in the Płonka sum) whose index is the top element of the lattice of indexes. Indeed the structure of an injective involutive bisemilattices strongly relies on all possible embeddings one can have between algebras in the sum. To give a more precise intuition, consider the involutive bisemilattice introduced in Example 5 and a different one, constructed recurring to the same index set and the same algebras, but with different embeddings, namely $p_{ik}(a) = d = p_{jk}(b)$. The two involutive bisemilattices have the same Booleanisation \mathbf{A}_k , although they are not isomorphic; this can be checked directly, or also reasoning the categorical equivalence between involutive bisemilattices and semilattice direct systems of Boolean algebras shown in [11] (see also [8]).

Given a Boolean algebra \mathbf{A}_∞ , an obvious example of (finite) injective involutive bisemilattice is one constructed over a direct system formed by subalgebras of \mathbf{A}_∞ (with \mathbf{A}_∞ as top element of the index set) and whose homomorphisms are inclusions. We will call these kinds of involutive bisemilattices *inclusive* (see Definition below). We wonder whether it is possible to count the number of non-isomorphic *inclusive* involutive bisemilattices having the same Booleanisation. We will show some peculiar cases, which, in our view, give a gist of the hardness of the problem announced in Section 4 (which we cannot solve here).¹⁶

In order to define inclusive involutive bisemilattices, let \mathbf{A}_∞ a finite Boolean algebra such that $|A_\infty| = 2^n$ and consider a collection $\{\mathbf{A}_1, \dots, \mathbf{A}_n\}$ of distinct subalgebras of \mathbf{A}_∞ such that $\mathbf{A}_1 = \mathbf{A}_{i_0}$, $\mathbf{A}_n = \mathbf{A}_\infty$ and, for each $i, j \in \{1, \dots, n\}$, either $A_i \subset A_j$ or $A_j \subset A_i$. In other words, the collection of subalgebras $\{\mathbf{A}_1, \dots, \mathbf{A}_n\}$ consists of a maximal subchain in the lattice of all subalgebras of \mathbf{A}_∞ .¹⁷

Definition 47. A finite involutive bisemilattice \mathbf{B} with Booleanisation \mathbf{A}_∞ is *inclusive* if it is the Płonka sum over a direct system whose elements are subalgebras in the collection $\{\mathbf{A}_1, \dots, \mathbf{A}_n\}$ and homomorphisms are inclusions from A_l to A_m in case $l \leq m$, for some $l, m \in \{1, \dots, n\}$.

We will call *weight* k of an involutive bisemilattice \mathbf{B} , the number of algebras in its Płonka sum representation. Observe that all inclusive involutive bisemilattice are injective and that, since, we are considering finite algebras then the index set forms a lattice.

Problem. Compute the number $N(\mathbf{A}_\infty, k)$ of all non-isomorphic inclusive involutive bisemilattices of weight k and Booleanisation \mathbf{A}_∞ .


¹⁵ It is clear that two isomorphic involutive bisemilattices have isomorphic Booleanisations.

¹⁶ A similar problem, connecting with counting the number of a specific subclass of involutive bisemilattices has been addressed in [15].

¹⁷ By *maximal subchain* we mean a sublattice which is a chain and contains a copy of any Boolean subalgebra (of \mathbf{A}_∞) with cardinality less or equal to 2^n .

Expecting to find a formula counting $N(\mathbf{A}_\infty, k)$ is a hopeless effort. Indeed, counting the number of non-isomorphic inclusive involutive bisemilattice constructed over a direct system containing k copies of the same algebra (\mathbf{A}_∞), is equivalent to counting the number of all non-isomorphic finite lattices with k elements. This is a problem for which no formula is known to work and which is indeed solved by a specific algorithm [29] (implemented and improved in the case of modular lattices in [31]). However, there are fortunate cases, where it is possible to find a formula counting $N(\mathbf{A}_\infty, k)$. We address the case where \mathbf{B} is an inclusive involutive bisemilattice such that $B \setminus \{A_1, A_\infty\}$ consists of $k - 2$ distinct subalgebras of \mathbf{A}_∞ (thus, $k - 2 \leq n$). We refer to the number of all non-isomorphic inclusive involutive bisemilattices, in this first case, as $N_d(\mathbf{A}_\infty, k)$.

To make an example of the case under consideration here, let \mathbf{A}_∞ be the eight-elements Boolean algebra ($n = 3$). Then the collection $\{\mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3\}$ consists of a copy of the two-elements Boolean algebra (\mathbf{A}_1), a copy of the four-elements Boolean algebra (\mathbf{A}_2) and the eight-elements (\mathbf{A}_3) one. It can be immediately checked that, for instance, for $k = 4$, we have $N_d(\mathbf{A}_\infty, k) = 8$. The Plonka sum representation of the eight non-isomorphic involutive bisemilattices are depicted in the following drawing (where arrows stands for inclusions).



Fortunately, in this case, it is not necessary to recur to the algorithm counting all (non-isomorphic) finite lattices. To see why, we begin by providing an useful lemma, which allows us to count the number of (non-isomorphic) inclusive involutive bisemilattice which differs only for the Boolean algebra whose index is the least element in the index lattice. To this end, let $X = \{1, \dots, n\}$ consisting of the first n natural numbers. Let $1 \leq h \leq n$ be fixed, and, for each $s \in X$, let $\mathcal{P}_s(h)$ be the set consisting of all subsets of X of cardinality h , having s as minimum element (with respect to the natural linear order over X).

Lemma 48. $\sum_{s=1}^n s |\mathcal{P}_s(h)| = \binom{n+1}{h+1}$.

Proof. Observe that there are $n - s$ possible choices of subsets of X of h elements containing s , thus $|\mathcal{P}_s(h)| = \binom{n-s}{h-1}$.

Therefore $\sum_{s=1}^n s |\mathcal{P}_s(h)| = \sum_{s=1}^n s \binom{n-s}{h-1} = \binom{n+1}{h+1}$, where the last equality easily follows by induction over n (for fixed h). ■

A simple example may help to grasp the content of the previous Lemma and its utility for our purposes. Let $n = 3$, so $X = \{1, 2, 3\}$, and $h = 2$. Then, $\mathcal{P}_1(2) = \{\{1, 2\}, \{1, 3\}\}$ and $\mathcal{P}_2(2) = \{\{2, 3\}\}$. Thus, $\sum_{s=1}^n s |\mathcal{P}_s(h)| = 1 \cdot 2 + 2 \cdot 1 = 4 = \binom{3+1}{2+1} = \binom{4}{3}$.

Theorem 49. Let \mathbf{B} be an inclusive involutive bisemilattice with Booleanisation \mathbf{A}_∞ , whose Plonka sum representation consists of $k - 2$ distinct subalgebras of \mathbf{A}_∞ . Then

$N_d(\mathbf{A}_\infty, k) = \binom{n+1}{k-1} a(k-2)$, where $a(k-2)$ is the number of acyclic graph with $k - 2$ vertices.

Proof. Observe that all the possible different (non isomorphic) inclusive involutive bisemilattices obtained by setting a different algebra in the place of the least element in the index set are counted, via Lemma 48 (setting $h = k - 2$), by $\binom{n+1}{k-1}$. Finally, we have to count the possible posets with $k - 2$ elements and this number is equivalent to the number $a(k - 2)$ is the number of acyclic graph with $k - 2$ vertices. This is counted by the following inductive formula:

$$a(k - 2) = \sum_{q=1}^{k-2} \binom{k-3}{q-1} q^{(q-2)} a(k - 2 - q),$$

where $q^{(q-2)}$ is the Cayley's formula, counting the number of trees over q vertices (of the acyclic graph), where, by convention, we assume $a(0) = 1$. ■

Solving the general case, namely determining a way to count $N(\mathbf{A}_\infty, k)$ could be delivered by “combining” two relevant cases: the one where the involutive bisemilattice is constructed via $k - 2$ distinct subalgebras of the Booleanisation \mathbf{A}_∞ and the relevant one where it contains an arbitrary number of copies of the same subalgebra of \mathbf{A}_∞ . As mentioned, the combination of the two cases gives itself raise to a hard problem which requires the application of an algorithm for counting the number of non-isomorphic (finite) lattices. Yet, an understanding of “how many” those algebras are (the number of injective involutive bisemilattices is greater than the inclusive ones!) does not provide a criterion to characterise when two injective involutive bisemilattices have the same Booleanisation (our original problem), but gives a hint of the difficulty of the enterprise, which we were not able to solve in the present work.

References

- [1] S. Aguzzoli, M. Bianchi, B. Gerla, D. Valota, Probability measures in Gödel $_{\Delta}$ logic, in: A. Antonucci, L. Cholvy, O. Papini (Eds.), *Symbolic and Quantitative Approaches to Reasoning with Uncertainty*, Springer, 2017, pp. 353–363.
- [2] S. Aguzzoli, M. Bianchi, V. Marra, A temporal semantics for basic logic, *Stud. Log.* 92 (2) (2009).
- [3] S. Aguzzoli, B. Gerla, Probability measures in the logic of nilpotent minimum, *Stud. Log.* 94 (2010) 151–176.
- [4] S. Aguzzoli, B. Gerla, V. Marra, De Finetti's no-Dutch-book criterion for Goedel logic, *Stud. Log.* 90 (2008) 25–41.
- [5] M. Bennett, States on orthomodular lattices, *J. Nat. Sci. Math.* 8 (1968) 47–52.
- [6] K. Bhaskara Rao, M. Bhaskara Rao, On the lattice of subalgebras of a Boolean algebra, *Czechoslov. Math. J.* 29 (4) (1979) 530–545.
- [7] G. Birkhoff, J. Von Neumann, The logic of quantum mechanics, *Ann. Math.* 37 (4) (1936) 823.
- [8] S. Bonzio, Dualities for Plonka sums, *Log. Univers.* 12 (2018) 327–339.
- [9] S. Bonzio, J. Gil-Férez, F. Paoli, L. Peruzzi, On paraconsistent weak Kleene logic: axiomatization and algebraic analysis, *Stud. Log.* 105 (2) (2017) 253–297.
- [10] S. Bonzio, A. Loi, The Plonka product of topological spaces, *Algebra Univers.* 29 (80) (2019).
- [11] S. Bonzio, A. Loi, L. Peruzzi, A duality for involutive bisemilattices, *Stud. Log.* 107 (2019) 423–444.
- [12] S. Bonzio, T. Moraschini, M. Pra Baldi, Logics of left variable inclusion and Plonka sums of matrices, *Arch. Math. Log.* 60 (2021) 49–76.
- [13] S. Bonzio, M. Pra Baldi, Containment logics: algebraic completeness and axiomatization, *Stud. Log.* (2021), <https://doi.org/10.1007/s11225-020-09930-1>, in press.
- [14] S. Bonzio, M. Pra Baldi, Containment logics: algebraic counterpart and reduced models, Submitted Manuscript (2021).
- [15] S. Bonzio, M. Pra Baldi, D. Valota, Counting finite linearly ordered involutive bisemilattices, in: J. Desharnais, W. Guttmann, S. Joosten (Eds.), *Relational and Algebraic Methods in Computer Science*, Springer International Publishing, Cham, 2018, pp. 166–183.
- [16] C. Borlido, M. Gehrke, A. Krebs, H. Straubing, Difference hierarchies and duality with an application to formal languages, *Topol. Appl.* 273 (2020) 106975.
- [17] R. Cignoli, I. d'Ottaviano, D. Mundici, *Algebraic Foundations of Many-Valued Reasoning*. Trends in Logic, Springer, Netherlands, 2000.
- [18] M. Dalla Chiara, R. Giuntini, R. Greechie, *Reasoning in Quantum Theory: Sharp and Unsharp Quantum Logics*. Trends in Logic, Springer, Netherlands, 2004.
- [19] B. de Finetti, Sul significato soggettivo della probabilità, *Fundam. Math.* 17 (1931) 289–329.
- [20] B. de Finetti, *Theory of Probability*, vol. 1, John Wiley and Sons, New York, 1974.
- [21] A. Di Nola, I. Leustean, Mv-algebras and Łukasiewicz logic, in: P. Cintula, P. Hájek, C. Noguera (Eds.), *Handbook of Mathematical Fuzzy Logic*, vol. 2, College Publications, 2011, pp. 469–583.
- [22] T. Flaminio, Three characterizations of strict coherence on infinite-valued events, *Rev. Symb. Log.* 13 (3) (2020) 593–610.
- [23] T. Flaminio, L. Godo, S. Ugolini, Towards a probability theory for product logic: states, integral representation and reasoning, *Int. J. Approx. Reason.* 93 (2018) 199–218.
- [24] T. Flaminio, T. Kroupa, States of MV Algebras, *Handbook of Mathematical Fuzzy Logic*, vol. 3, College Publications, London, 2015, pp. 1191–1245.
- [25] D. Foulis, R. Greechie, Probability weights and measures on finite effect algebras, *Int. J. Theor. Phys.* 38 (1999) 3189–3208.
- [26] S. Givant, P. Halmos, *Introduction to Boolean Algebras*, Undergraduate Texts in Mathematics, Springer, New York, 2008.
- [27] K. Grätzer, M. Koh, M. Makkai, On the lattice of subalgebras of a Boolean algebra, *Proc. Am. Math. Soc.* 36 (1) (1972) 87–92.
- [28] F. Haimo, Some limits of Boolean algebras, *Proc. Am. Math. Soc.* 2 (4) (1951) 566–576.
- [29] J. Heitzig, J. Reinhold, Counting finite lattices, *Algebra Univers.* 48 (1) (2002) 43–53.
- [30] J. Howie, *Fundamentals of Semigroup Theory*, Oxford University Press, 1995.
- [31] P. Jipsen, N. Lawless, Generating all finite modular lattices of a given size, *Algebra Univers.* 74 (2015) 253–264.
- [32] J. Kelley, *General Topology*, Graduate Texts in Mathematics, Springer, New York, 1975.
- [33] J.L. Kelley, Measures on Boolean algebras, *Pac. J. Math.* 9 (4) (1959) 1165–1177.
- [34] A.N. Kolmogorov, Complete metric Boolean algebras, *Philos. Stud.* 77 (1) (1995).
- [35] T. Kroupa, Every state on semisimple MV-algebra is integral, *Fuzzy Sets Syst.* 157 (20) (2006) 2771–2782.
- [36] M.V. Lawson, The Booleanization of an inverse semigroup, *Semigroup Forum* 100 (1) (2020) 283–314.
- [37] I. Leustean, Metric completions of MV-algebras with states: an approach to stochastic independence, *J. Log. Comput.* 21 (3) (2009) 493–508.
- [38] T. Memp, Interior preserving maps, *Bull. Aust. Math. Soc.* 1 (2) (1969) 209–212.
- [39] D. Mundici, Averaging the truth-value in Łukasiewicz logic, *Stud. Log.* 55 (1) (1995) 113–127.

- [40] D. Mundici, Bookmaking over infinite-valued events, *Int. J. Approx. Reason.* 43 (3) (2006) 223–240.
- [41] J.R. Munkres, *Topology*, 2nd edition, Prentice Hall, 2000.
- [42] M. Negri, A probability measure for partial events, *Stud. Log.* 94 (2010) 271–290.
- [43] G. Panti, Invariant measures in free MV-algebras, *Commun. Algebra* 36 (8) (2008) 2849–2861.
- [44] F. Paoli, M. Pra Baldi, Extensions of paraconsistent weak Kleene logic, *Log. J. IGPL* (2021), <https://doi.org/10.1093/jigpal/jzaa024>, in press.
- [45] M. Petrich, *Lectures in Semigroups*, Wiley, London, 1977.
- [46] T. Pirttmäki, A survey of Kolmogorov quotients, *ArXiv pre-print*, 2019.
- [47] J. Plonka, On a method of construction of abstract algebras, *Fundam. Math.* 61 (2) (1967) 183–189.
- [48] J. Plonka, On distributive quasilattices, *Fundam. Math.* 60 (1967) 191–200.
- [49] J. Plonka, On the sum of a direct system of universal algebras with nullary polynomials, *Algebra Univers.* 19 (2) (1984) 197–207.
- [50] J. Plonka, A. Romanowska, Semilattice sums, in: A. Romanowska, J. Smith (Eds.), *Universal Algebra and Quasigroup Theory*, Heldermann, 1992, pp. 123–158.
- [51] B. Weatherston, From classical to intuitionistic probability, *Notre Dame J. Form. Log.* 44 (2) (2003) 111–123.