

1 **Structure–Activity Relationship of Pyrrolyl Diketo Acid Derivatives as Dual**
2 **Inhibitors of HIV-1 Integrase and Reverse Transcriptase Ribonuclease H Domain**

3 Giuliana Cuzzucoli Crucitti,† Mathieu Metfiot,‡ Luca Pescatori,† Antonella Messori,†
4 Valentina Noemi Madia,† Giovanni Pupo,† Francesco Saccoliti,† Luigi Scipione,† Silvano
5 Tortorella,† Francesca Esposito,§ Angela Corona,§ Marta Cadeddu,§ Christophe
6 Marchand,‡ Yves Pommier,‡ Enzo Tramontano,§ Roberta Costi,* ,†,|| and Roberto Di
7 Santo†,||

8

9 †Dipartimento di Chimica e Tecnologie del Farmaco, Istituto Pasteur-Fondazione Cenci
10 Bolognetti, “Sapienza” Università di Roma, Rome, I-00185, Italy

11 ‡Developmental Therapeutics Branch and Laboratory of Molecular Pharmacology, Center
12 for Cancer Research, National Cancer Institute, National Institutes of Health, Building 37,
13 Room 5068, Bethesda, Maryland 20892-4255, United States

14 §Department of Life and Environmental Sciences, University of Cagliari, Cagliari, 09124,
15 Italy

16 *S Supporting Information

17 **AUTHOR INFORMATION**

18

19 Corresponding Author:

20

21 *Phone: +39-06-49693247. Fax: +39-06-49913133. E-mail:

22

23 roberta.costi@uniroma1.it.

24

25

26 **Author Contributions**

27

28 ||R.C. and R.D.S. contributed equally.

29 The manuscript was written through contribution of all authors. All authors have given
30 approval to the final version of the manuscript.

31

32 **Notes**

33

34

35 The authors declare no competing financial interest.

36

37 **ABBREVIATIONS USED**

38 IN, integrase; RT, reverse transcriptase; RNase H, ribonuclease H; 3'-P, 3'-processing;

39 ST, strand transfer; DKA, diketo acid; BTDBA, 4-[5-(benzoylamino)thien-2-yl]-2,4-

40 dioxobutanoic acid; SI, selectivity index; TosMIC, toluene-4-sulfonylmethyl isocyanide;

41 NIS, N-iodosuccinimide; GP, general procedure; DMEM, Dulbecco's modified Eagle

42 medium

43

44

45

46

47

48

49

50

51


52

53 **ABSTRACT:**

54

55 The development of HIV-1 dual inhibitors is a highly innovative approach aimed at reducing
56 drug toxic side effects as well as therapeutic costs. HIV-1 integrase (IN) and reverse
57 transcriptase-associated ribonuclease H (RNase H) are both selective targets for HIV-1
58 chemotherapy, and the identification of dual IN/RNase H inhibitors is an attractive strategy
59 for new drug development. We newly synthesized pyrrolyl derivatives that exhibited good
60 potency against IN and a moderate inhibition of the RNase H function of RT, confirming the
61 possibility of developing dual HIV-1 IN/RNase H inhibitors and obtaining new information for
62 the further development of more effective dual HIV-1 inhibitors.

63



73 INTRODUCTION

74

75 The development of dual-action drugs is a promising approach to ameliorate drug–drug
76 interactions, reduce toxic side effects, and suppress viral resistance selection.^{1–4} Among
77 dual-action drugs, dual inhibitors are single compounds that are able to inhibit two enzyme
78 activities. Several reports have shown that dual inhibitors may have a role in the treatment
79 of different diseases such as Alzheimer,⁵ Parkinson,⁶ inflammation,⁷ and cancer.^{1,8,9} This
80 approach had been attempted also in the virological arena, aiming to inhibit rhinovirus
81 replication.¹⁰ Recently, tropolones,^{11–13} madurahydroxylactone,¹⁴ and 2-hydroxyisoquinolin-
82 1,3(2H,4H)-diones^{15,16} have been reported to act as dual inhibitors against HIV-1, targeting
83 viral integrase (IN) and reverse transcriptase (RT) ribonuclease H (RNase H) activities.

84

85 HIV-1 IN is the viral enzyme responsible for the integration of the proviral dsDNA into the
86 cell host chromosome through two coordinated enzyme functions, both accomplished by the
87 same active site.¹⁷ In the first reaction, termed 3'-end processing (3'-P), IN removes the two
88 terminal nucleotides (GT) from each 3'-end of the dsDNA.¹⁷ In the second reaction, termed
89 strand-transfer (ST), IN catalyzes a nucleophilic attack by the free 3'-OH of the viral
90 processed DNA to the target chromosomal DNA, resulting in covalent joining of the two
91 molecules. Several classes of integrase inhibitors have been identified;¹⁸ among these the
92 diketoacids (DKAs) showed greatest promise, and the first DKA bioisoster, raltegravir (1),
93 has been approved in 2007 for HIV-1 therapy.¹⁹ HIV-1 IN belongs to the functionally diverse
94 superfamily of DDE(D) nucleotidyltransferases, whose other notable mem-bers include
95 RNaseH and MuA, Tn5, and Mos1 transposases.¹⁷ The active sites of these enzymes
96 typically contain three essential carboxylates that coordinate a pair of divalent metal cations
97 (usually Mg²⁺). Thus, chelating inhibitors can be active across several classes of viral metal-

98 dependent enzymes and chelation has been successfully used in drug design, also of dual
99 inhibitors.²⁰ In particular, DKAs have been reported to chelate the divalent cations in the IN
100 active site,^{17,18} and notably, DKAs originally developed against HIV-1 IN have been also
101 reported to inhibit the HIV-1 RNase H.^{21,22} The HIV-1 RT-associated RNase H function
102 hydrolyzes the RNA strand of the replicative intermediate RNA:DNA hybrid and, hence, is
103 essential for viral replication.²³ Even though several compounds have been recently
104 identified to inhibit this RT activity,²⁴⁻³⁰ up to today no RNase H inhibitor has been approved
105 for HIV therapy.

106 Therefore, this viral function is a very attractive target for drug development.

107 The first DKA IN inhibitor later described also as RNase H inhibitor was the 4-[5-
108 (benzoylamino)thien-2-yl]-2,4-dioxobutanoic acid (BTDBA, 2) discovered by Merck.²¹
109 Recently we reported that 6-[1-(4-fluorophenyl)methyl-1H-pyrrol-2-yl]-2,4-dioxo-5-hexenoic
110 acid ethyl ester (RDS 1643, 3) inhibited the HIV-1 RT-associated RNase H function in
111 biochemical assays with an IC₅₀ value of 8 μM, the HIV-1 IN ST (the IC₅₀ value was 98
112 μM) and blocked the HIV-1 replication in cell based assays with an EC₅₀ value of <0.2 μM
113 (Chart 1).²² Starting from these observations, more recently we designed a small library of
114 3 analogues with the aim of obtaining dual IN/ RNase H HIV-1 inhibitors and found
115 compounds active at micromolar concentration against RNase H and low nanomolar IC₅₀
116 values against IN in recombinant assays.³¹ The best dual inhibition was found for
117 compounds showing a diketo ester group and fluorine atoms as substituents on the benzyl
118 portion, as exemplified by compound 4 (Chart 1, IC_{50,IN} = 0.6 μM, IC_{50,RH} = 3.0 μM, EC₅₀
119 = 2 μM). Furthermore, we extended these results to a quinolonyl diketo acid series, in which
120 less marked dual activity was found, having in the best case IC_{50,IN} = 26.2 μM, IC_{50,RH} =
121 2.4 μM, and EC₅₀ = 3.6 μM (compound 5).³²

122

123 Herein we present the design, synthesis, and biological evaluation of new compounds
124 related to 3 to better define the structure–activity relationship (SAR) within the most interest-
125 ing pyrrolyl diketo acid series. All the newly designed pyrrolyl DKAs, both ester 6 and acid 7
126 derivatives, are shown in Chart 2.

127 Basically, starting from 3, we fixed the pyrrole ring and the DKA chain while wider
128 transformations included one or more of the following modifications: (i) introduction of
129 aromatic substituent in position 4 of the pyrrole ring; (ii) shift of the diketo hexenoic chain
130 from 2 to 3 position of the pyrrole moiety; (iii) shortening of the diketo hexenoic branch into
131 a diketo butanoic group; (iv) introduction of alkyl or aryl group replacing the fluorobenzyl
132 moiety; (v) replacement of carboxylic function with a triazole ring; (vi) introduction of alkyl
133 group within the DKA branch; (vii) replacement of keto group of DKA moiety with NH₂
134 function. Notably, among the compounds described in this paper, 7k has been the first DKA
135 derivative reported by Merck as selective ST IN inhibitor.³³ We decided to include this
136 compound in this study to define its properties as dual inhibitor of IN and RNase H.

137

138

139

140

141

142

143

144

145

146 RESULTS AND DISCUSSION

147

148 Chemistry

149 The synthesis of derivatives 6a–l and 7a–m is outlined in Schemes 1–4. Derivatives 6a–d
150 and 7a–d were synthesized according to the pathway described in Scheme 1. The acetyl or
151 propionyl pyrrole intermediates 9a–c were obtained by two different procedures: (1) the
152 alkylation with 4-fluorobenzyl bromide in alkaline medium (K₂CO₃) of 1-(4-phenyl-1H-
153 pyrrol-3-yl)ethanone³⁴ or derivatives 8, achieved via toluene-4-sulfonylmethyl isocyanide
154 (TosMIC) reaction from (E)-1-phenylpent-1-en-3-one;³⁵ (2) thermal transposition of the
155 acetyl chain of 2-acetyl-1-[(4-fluorophenyl)methyl]pyrrole³⁶ from 2- to 3- position of the
156 pyrrole ring in the presence of CF₃COOH. Derivatives 9a–c were condensed in turn with
157 diethyl oxalate in the presence of sodium ethoxide to provide the diketobutanoic ethyl esters
158 6a–c. Compound 6a was used as substrate to provide (i) 6d, obtained by reacting the enol
159 6° with ammonium acetate in acid medium (CH₃COOH) following a known procedure
160 reported for DKA derivatives,³⁷ and (ii) 7a, achieved by hydrolysis of ester 6a in the
161 presence of 6 N NaOH. The last conditions have also been used to obtain 7b–c starting from
162 6b–c, respectively. A slightly different condition has been used to obtain 7d, as previously
163 described.³⁷ Derivatives 6e–h and 7e–h were obtained according to Scheme 2. The
164 pyrroles 10 and 11 were used as starting materials to obtain the key intermediates 12e–h.
165 Compound 10 was obtained by alkylation of commercially available 4-iodo-2-formylpyrrole
166 in alkaline medium (K₂CO₃) with 4-F-benzyl bromide. Derivative 11 was obtained starting
167 from pyrrole that underwent a two-step one-pot reaction comprising (i) formylation in the
168 presence of oxalyl chloride and DMF and (ii) Friedel–Crafts reaction with 4-F-benzoyl
169 chloride. Thus, compound 11 was arylated to 12f or alkylated on nitrogen atom of pyrrole
170 ring with the appropriate alkyl halide furnishing 12e and 12g, while intermediate 10

171 underwent a Suzuki coupling in position 4 to obtain 12h. The formylpyrroles 12e–h were
172 condensed with acetone in alkaline medium (4 N NaOH) to obtain α,β -unsaturated ketones
173 13e–h. The last compounds were in turn condensed with diethyl oxalate in the presence of
174 sodium ethoxide to provide the diketobutanoic ethyl esters 6e–h that were finally hydrolyzed
175 by reaction with 1 N NaOH to give the corresponding carboxylic acids 7e–h. The synthetic
176 pathway to obtain derivatives 6i,j and 7i,j is outlined in Scheme 3. The enones 17i,j were
177 obtained starting from 15 and 16, respectively. Pyrrole 15 has been achieved starting from
178 pyrrole-2-carboxaldehyde that was alkylated in alkaline medium (K_2CO_3) to obtain 14. The
179 thermal transposition of the formyl chain from position 2 to 3 of the pyrrole ring in the presence
180 of trifluoroacetic acid led to derivative 15. In a parallel pathway (3E,5E)-6-phenylhexa-3,5-
181 dien-2-one³⁸ underwent ring closure by reacting with TosMIC, giving the pyrrole derivatives
182 16. Interestingly, the attack of the TosMIC reagent was specific on the 5,6 double bond of
183 the starting dienone. Afterward, derivative 15 was condensed with acetone, affording 17i;
184 conversely, compound 16 was converted into 17j by reaction with 4-F-benzyl bromide in
185 alkaline medium (K_2CO_3). Finally, intermediates 17i,j were converted into esters 6i,j by
186 condensation with diethyl oxalate and then converted to acids 7i,j by alkaline hydrolysis.

187 Derivatives 6k,l and 7k,l were obtained according to the pathway described in Scheme 4.
188 The iodination of 2-acetyl-1-[(4-fluorophenyl)methyl]pyrrole in the presence of
189 Nodosuccinimide (NIS) afforded derivative 18, which underwent a Suzuki coupling reaction
190 to give 19. Intermediates 19 and 2-acetyl-1-[(4-fluorophenyl)methyl]pyrrole were converted
191 into the diketobutanoic acid derivatives 7k,l through the ethyl esters 6k,l, according to the
192 condensation/hydrolysis procedure described above. Finally, 2-acetyl-1-[(4-
193 fluorophenyl)methyl]pyrrole was subjected to a condensation reaction in the presence of 1-
194 trityl-1H-[1,2,4]triazole-3-carboxylic acid ethyl ester³⁹ to afford the triazole-protected

195 derivative 20, which was deprotected by hydrolysis in the presence of 3 M HCl to obtain the
196 triazole derivative 7m.

197 **Evaluation of Biological Activities.**

198 All newly synthesized compounds 6a–l and 7a–m were tested in vitro in enzyme assays
199 against both recombinant RNase H and IN. The IC₅₀ values obtained for each compound
200 in the inhibition of both the IN ST reaction and HIV-1 RT-associated RNase H function were
201 plotted against each other in correlation plots (Figure 1). In these plots, single dots
202 correspond to single compounds. The compounds are distributed around two perpendicular
203 axes crossing the IN IC₅₀ axis (X axis) at 1 μM and the RNase H IC₅₀ axis (Y axis) at 10
204 μM (bolded crosshair in the center of each graph, Figure 1). These two axes splice the graph
205 into four quarters corresponding to RNase H/IN dual inhibitors (lower left quarter), RNase
206 H-selective inhibitors (lower right quarter), IN selective inhibitors (upper left quarter), and
207 inhibitors of lower potency (upper right quarter). As seen in Figure 1, these graphs do not
208 show any particular correlation between RNase H and IN inhibition.

209 The newly synthesized pyrrolyl derivatives 6a–l and 7a–m exhibited good potency of
210 inhibition when tested on the HIV-1 IN ST (Table 1). In general, as reported previously,^{31,32}
211 the acid derivatives 7a–m were endowed with the highest potency showing IC₅₀ values in
212 the range of 26–0.019 μM. In our assay Merck compound 7k was confirmed as potent IN
213 ST inhibitor showing IC₅₀ = 0.057 μM (literature data IC₅₀ = 0.08 μM).³³

214 As seen in Figure 1A, 80% of the acid compounds (red dots) are distributed in the left half
215 of the graph, suggesting the acid function is critical for IN inhibition but not critical for RNase
216 H inhibition. Among them, seven compounds (7a,c,d,e,h,k,l) were proven to have
217 submicromolar activity (IC₅₀ value were in the range of 66–19 nM), while two derivatives
218 (7b,i) can be considered less active to almost inactive (IC₅₀ values were 111 and 26 μM,

219 respectively). The most active compound was the 4-phenylpyrrole derivative 7l with an IC₅₀
220 value of 19 nM, 3 times less active than 1 in inhibiting of the ST reaction of IN.

221 Interestingly, even though not all the synthesized analogues were tested on the HIV-1 3'-P
222 activity, results showed that the newly synthesized acids were selective inhibitors of the ST
223 step, with the IC₅₀ values on the 3'-P step 2–3 orders of magnitude higher with respect to
224 the ones obtained on ST, thus confirming that the DKAs are selective ST inhibitors (data not
225 shown). The newly synthesized pyrrolyl DKA derivatives 6a–l and 7a–m can be divided into
226 two classes: the diketobutanoic (6a– d,k,l and 7a–d,k,l) and the diketohexenoic (6e–j and
227 7e–j) derivatives. In the diketobutanoic structures of derivatives 7a– d, 7k,l, two main
228 differences involving the substitution of the pyrrole ring emerged: the diketobutanoic chain
229 can be linked on the pyrrole ring into two different positions (2- or 3- position), along with a
230 phenyl ring, which can be substituted in position 4 of the pyrrole ring. The two variables did
231 not influence the IN inhibitory activity. Only the phenyl substitution at position 4 of the pyrrole
232 ring (R substituent) seems to favor slightly IN vs RNase H inhibition with a majority of
233 compounds bearing this substitution distributed in the two left quarters of the correlation
234 graphs (green dots, Figure 1B). In fact, from a comparison of the inhibition data of
235 diketobutanoic/diketohexenoic 7a/7l and 7c/7k, which are characterized respectively by the
236 presence of the phenyl ring and by its absence, a correspondence in the orders of magnitude
237 of their IC₅₀ values was observed. Moreover, 7a and 7c linked the diketobutanoic chain in
238 position 3 of the pyrrole ring, while 7k and 7l linked it in 2-position of the same ring. All these
239 compounds were characterized by similar potencies (IC₅₀ values of 22, 24, 57, and 19 nM,
240 respectively). The 4-fluorobenzoyl substitution at position 4 of the pyrrole ring (R substituent)
241 does not seem to favor the inhibition of either enzymes (orange dots, Figure 1B).

242

243 The other three modifications involved exclusively the diketobutanoic chain: (1) the
244 substitution of the enolic OH with a NH₂ (7d), (2) the introduction of a methyl group in 3-
245 position (7b), and (3) the substitution of the carboxylic acid function with its bioisoster triazole
246 ring (7m). The NH₂ (7d) and triazole (7m) derivatives were 2-fold less active with respect to
247 their OH (7a) and COOH (7h) counterparts, respectively (IC₅₀ values were 43, 110, 22, and
248 57 nM, respectively). On the contrary, the 3-methylbutenoate derivative (7b) completely lost
249 its ability to inhibit IN enzyme (IC₅₀ values of >111 μM). It appears that IN seems to tolerate
250 a wide variety of substitutions at the R1 position, including the absence of substituent, with
251 a majority of compounds with such substitutions distributed in the two left quarters of the
252 correlation graphs (Figure 1C). No preferential inhibition pattern can be observed for
253 compounds with substitutions at the R2 position on the correlation plot presented in Figure
254 1 D.

255

256 When the two series of diketobutanoic and the diketohex-enoic esters and acids were tested
257 on the HIV-1 RT-associated RNase H function in biochemical assays, the most active
258 derivatives were the diketohexenoic ester 6f and the diketobutanoic acid 7d, with IC₅₀ value
259 of 1.8, and 2 μM, respectively. Interestingly, both compounds were more active than the
260 references 3 and 4. Noteworthy, the structure of 6f is related to the reference 3, since both
261 are pyrrolyl diketohex-enoic derivatives, but 6f bring a phenyl ring on nitrogen replacing the
262 4-F-benzyl group of 3 and a 4-F-benzoyl moiety linked in 4 position of the pyrrole ring. In the
263 matter of compound 7d, it is a 3-pyrrolyl diketobutanoic acid derivative characterized by the
264 presence of an amine function that replaces the enol OH in position 3 of the diketobutanoic
265 chain. Since 7d is a carboxylic acid, its stronger inhibition toward IN than RNase H function
266 was expected; conversely, the ester 6f is the best dual inhibitor IN/RNase H of this series.
267 From a first analysis of the results we can state that (i) as known, the acid function confers

268 a better inhibitory activity on IN, (ii) the ester function is amenable for inhibition of RNase H
269 function of RT, confirming the results recently reported by the means of docking studies and
270 mutagenesis experiments,⁴⁰ and (iii) contemporaneously, the ester function is necessary
271 for a dual inhibition IN/RNase H.

272

273 Among the ester derivatives (6a–l), the diketohexenoic compounds (6e–j, IC₅₀ values in the
274 range of 1.8–55 μM) were able to inhibit the HIV-1 RNase H function with a slightly higher
275 potency than the diketobutanoic counterpart (6a–d,k,l, IC₅₀ values in the range of 6–72 μM).
276 Differently, when the acid derivatives (7a–m) were tested, this difference was not observed.

277

278 In both the diketohexenoic and the diketobutanoic ester and acid derivatives, the presence
279 of a phenyl substituent in position 4 of the pyrrole ring influenced the inhibitory activity. In
280 fact, the 4-phenyl substituted diketobutanoic ester and acid 6a/7a (IC₅₀ values of 10 μM,
281 7.5 μM, respectively) and diketohex-enoic 6j/7j (IC₅₀ values of 3.0 μM, 7.0 μM, respectively)
282 were consistently more active than the unsubstituted counterparts 6c/7c and 6i/7i (IC₅₀
283 values of >100, 41, 55, and 69 μM, respectively).

284 In general, when the DKA chain was shifted from 2- to 3-position, a 2- to 4-fold increase in
285 potency of inhibition was observed (compare 6h, 7h, and 7l with 6j, 7j, and 7a: IC₅₀ values
286 of 13.4, 23, 14, 3, 7, and 7.5 μM, respectively), with the sole exception of 6a and 6l, which
287 showed comparable potency (IC₅₀ values of 10 and 6 μM, respectively).

288 The substitution of the enolic OH on the diketobutanoic chain (6a/7a) with a NH₂ group
289 (6d/7d) in the ester series led to a 7-fold decrease of the potency of RNase H inhibition (6d,
290 IC₅₀ values of 72 μM) with respect to the unmodified counterpart (6a, IC₅₀ value of 10 μM),
291 while within the acid series, the NH₂ derivative improved its potency of inhibition

292

293 (7d, IC₅₀ value of 2 μM; 7a, IC₅₀ value of 7.5 μM). The introduction of a methyl group in
294 position 3 of the diketobutanoic chain (7b) reduced the inhibition (7b and 7a IC₅₀ values of
295 64 and 7.5 μM, respectively) and likewise the substitution of the carboxylic acid function with
296 its bioisoster triazole (7a and 7m, IC₅₀ values of 7.5 μM and 26 μM, respectively).

297 Among all the newly synthesized derivatives, the 2-pyrrolyl diketohexenoic ester 6f, although
298 not very potent, emerged as dual inhibitor showing similar IC₅₀ values against both IN
299 enzyme and RNase H function of RT (1.2 and 1.8 μM, respectively). This compound retains
300 the ester function that is demonstrated to be necessary for the dual inhibition³¹ and is
301 characterized by a phenyl ring on nitrogen replacing the 4-F-benzyl group of 3 and by a 4-
302 F-benzoyl moiety linked in 4 position of the pyrrole ring.

303

304 **Cell-Based Assays.**

305 Among the newly synthesized pyrrolyl DKA derivatives 6a–l and 7a–m, seven derivatives
306 (6a,c,l and 7a,c,d,l) were characterized by a good anti-HIV activity showing a EC₅₀ values
307 in the submicromolar concentration (EC₅₀ in the range of 0.56–0.9 μM) and a good
308 selectivity index (SI). Three compounds, the 6k,j and 7k derivatives, showed an anti-HIV
309 activity in the range of the micromolar concentration (EC₅₀ in the range of 1–4.3 μM), while
310 10 compounds were less active or inactive (6d–i and 7b,e,f,i,m, EC₅₀ in the range of 17.2
311 to >50 μM). In general, all the compounds had a low cytotoxicity index (CC₅₀ > 50 μM).

312 Compound 7c, characterized by the diketobutanoic chain in 3-position of the pyrrole ring,
313 showed the best antiviral efficacy on HIV-1 infected cell (EC₅₀ = 0.58 μM) and a low
314 cytotoxicity (CC₅₀ > 50 μM, SI > 86). The best dual inhibitor derivative 6f that showed activity

315 at micromolar concentration in bio-chemical assays was 20 times less potent in cell-based
316 assays.

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335 **CONCLUSION**

336

337 The development of HIV-1 dual inhibitors is a highly innovative approach aimed at reducing
338 drug toxic side effects and therapeutic costs. Since HIV-1 IN and RNase H are both selective
339 targets for HIV-1 chemotherapy, the identification of dual IN/RNase H inhibitors is an
340 attractive strategy for new drug development. The newly synthesized pyrrolyl derivatives
341 6a–l and 7a–m exhibit good potency against IN and a moderate inhibition of the RNase H
342 function of RT. In general, by comparison of the inhibition data among the ester and the acid
343 derivatives, a different behavior was observed. As expected, the acid derivatives showed a
344 higher potency of IN inhibition with respect to the corresponding esters, while the latter
345 compounds have been often found more potent than the corresponding acids in inhibiting
346 the RNase H function of RT enzyme. Notably, compound 6f, although not very potent on
347 HIV-infected cell, showed a good correlation between HIV-1 IN and RNase H inhibition. It is
348 characterized by a diketoester function, a phenyl ring on nitrogen, and a 4-F-benzoyl moiety
349 linked in 4 position of the pyrrole ring. We can state that although the acid function confers
350 a better inhibitory activity on IN, an ester function is amenable for inhibition of RNase H
351 function of RT, and as a consequence, the ester function is necessary for a dual inhibition
352 IN/RNase H. These basic chemical features should be considered for development of more
353 potent dual inhibitors. Overall, the data reported in this work confirm the possibility of
354 developing dual HIV-1 IN/ RNase H inhibitors and give new information for the further
355 development of effective dual HIV-1 inhibitors.

356

357

358

359 **EXPERIMENTAL SECTION**

360

361 **Chemistry. General.**

362 Melting points were determined on a Bobby Stuart Scientific SMP1 melting point apparatus
363 and are uncorrected. Compound purities were always >95% determined by high pressure
364 liquid chromatography (HPLC). HPLC analyses were carried out with a Shimadzu LC-10AD
365 VP CTO-10AC VP. Column used was generally Discovery Bio Wide Pore C18 (10 cm × 4.6
366 mm, 3 μm). Infrared (IR) spectra were recorded on a PerkinElmer Spectrum-One
367 spectrophotometer. ¹H NMR spectra were recorded at 400 MHz on a Bruker AC 400
368 Ultrashield 10 spectrophotometer (400 MHz). Dimethyl sulfoxide-d₆ 99.9% (code 44,139-2)
369 and deuteriochloroform 98.8% (code 41,675-4) of isotopic purity (Aldrich) were used. Column
370 chromatographies were performed on silica gel (Merck; 70–230 mesh) column or aluminum
371 oxide (Sigma-Aldrich; 150 mesh) column. All compounds were routinely checked on TLC by
372 using aluminum-baked silica gel plates (Fluka DC-Alufolien Kieselgel 60 F254) or aluminum
373 oxide (Fluka DC-Alufolien). Developed plates were visualized by UV light. Solvents were
374 reagent grade and, when necessary, were purified and dried by standard methods. Concen-
375 tration of solutions after reactions and extractions involved the use of rotary evaporator
376 (Büchi) operating at a reduced pressure (~20 Torr). Organic solutions were dried over
377 anhydrous sodium sulfate (Merck). All reactions were carried out under nitrogen. All solvents
378 were freshly distilled under nitrogen and stored over molecular sieves for at least 3 h prior
379 to use.

380

381 **Microwave Irradiation Experiments.**

382 Microwave reactions were conducted using a CEM Discover system unit (CEM. Corp.,
383 Matthews, NC). The machine consists of a continuous focused microwave-power delivery
384 system with operator selectable power output from 0 to 300 W. The temperature of the
385 contents of the vessel was monitored using a calibrated infrared temperature control
386 mounted under the reaction vessel. All experiments were performed using a stirring option
387 whereby the contents of the vessel are stirred by means of a rotating magnetic plate located
388 below the floor of the microwave cavity and a Teflon-coated magnetic stir bar in the vessel.

389

390 **General Procedure A (GP-A):** Synthesis of Pyrrole Nucleus. A solution of α,β -unsaturated
391 ketone (5.42 mmol) and toluene-4-sulfonylmethyl isocyanide (1.16 g, 5.96 mmol, 1.1 equiv)
392 dissolved in a mixture of anhydrous dimethyl sulfoxide/ethyl ether (14:30 mL) was added
393 dropwise into a well-stirred suspension of sodium hydride (60% in paraffine oil; 0.48 g, 11.93
394 mmol, 2.2 equiv) in dry ethyl ether (30 mL) under argon atmosphere. After the addition the
395 mixture was stirred at room temperature for 1 h. The reaction was treated with water and
396 extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous
397 sodium sulfate, and concentrated under vacuum. The crude product was purified by
398 chromatography on aluminum oxide (chloroform as eluent) to afford the pure product. Yield
399 (%), melting point ($^{\circ}\text{C}$), recrystallization solvent, IR, and ^1H NMR are reported for each
400 compound.

401

402 **General Procedure B (GP-B):** Alkylation of the Pyrrole Nitrogen. A mixture of the
403 appropriate pyrrole (1.1 mmol), alkylating agent (3.3 mmol), and anhydrous K_2CO_3 (210
404 mg, 1.5 mmol) in dry DMF (10 mL) was stirred at $100\text{ }^{\circ}\text{C}$ for 2 h. Then the mixture was
405 cooled, treated with water (40 mL), and extracted with ethyl acetate. The organic layer was
406 washed with brine, dried over anhydrous sodium sulfate, and concentrated under vacuum.

407 The crude product was purified by chromatography on silica gel to afford the pure product.
408 Chromatography eluent, yield (%), melting point (°C), recrystallization solvent, IR, and ¹H
409 NMR are reported for each compound.

410 **General Procedure C (GP-C):** Condensation of Pyrrole Carboxaldehyde with Acetone. The
411 proper pyrrole carboxaldehyde (0.075 mol) was dissolved in 250 mL of acetone. To this
412 mixture was added 4 N NaOH (110 mL), and the mixture was stirred at room temperature
413 for 24 h. After this period water (300 mL) and ethyl acetate (250 mL) were added. The
414 organic layer was separated, washed with water (2 × 100 mL), dried over sodium sulfate,
415 filtered, and evaporated under reduced pressure. The crude product was purified by column
416 chromatography on silica gel to obtain pure products. Chromatography eluent, yield (%),
417 melting point (°C), recrystallization solvent, IR, and ¹H NMR are reported for each
418 compound.

419

420 **General Procedure (GP-D):** Acetyl Transposition. A mixture of opportune α acetyl-
421 substituted pyrrole (1.23 mmol) in trifluoroacetic acid (5 mL) was heated at 80 °C for 20 h.
422 After this period the reaction was quenched with water (30 mL) and extracted with ethyl
423 acetate (2 × 50 mL). The organic layers were collected, dried over sodium sulfate, filtered,
424 and evaporated under vacuum. The crude product was purified by chromatography on silica
425 gel (chloroform as eluent) to afford pure product as a brown oil. Yield (%), melting point (°C),
426 recrystallization solvent, IR, and ¹H NMR are reported for each compound.

427

428 **General Procedure E (GP-E):** Suzuki Reaction. Pd₂(dba)₃ (0.1 g, 1.7 mmol) was added
429 into a well stirred mixture of appropriate 4-iodopyrrole (1.7 mmol), phenylboronic acid (0.85
430 g, 7.0 mmol), Cs₂CO₃ (0.665 g, 2.0 mmol), and P(t-But)₃ in dioxane (20 mL). The mixture

431 was stirred at 80 °C for 24 h under argon atmosphere. Then the mixture was cooled to room
432 temperature, filtered, and washed with dioxane. The organic layer was evaporated under
433 vacuum. The raw material was extracted with water (50 mL) and ethyl acetate (50 mL). The
434 organic phase was separated, dried over sodium sulfate, filtered, and evaporated under
435 vacuum. The raw material was purified by silica gel chromatography. Chromatography
436 eluent, yield (%), melting point (°C), recrystallization solvent, IR, and ¹H NMR are reported
437 for each compound.

438 **General Procedure F (GP-F):** Synthesis of Diketo Esters. Freshly prepared sodium
439 ethoxide (390 mg, 5.5 mmol) was added into a well-stirred mixture of the appropriate acetyl
440 derivative (2.7 mmol) and diethyl oxalate (790 mg, 5.4 mmol) in anhydrous THF (2.7 mL)
441 under nitrogen atmosphere. The mixture was stirred at room temperature for 2 h and then
442 was poured into n-hexane (50 mL). The collected precipitate was vigorously stirred for 30
443 min in 1 N HCl (50 mL). The yellow solid that formed was filtered, washed with water, and
444 dried under IR lamp to afford the pure diketo esters. Yield (%), melting point (°C), IR, and
445 ¹H NMR are reported for each compound.

446

447 **General Procedure G (GP-G):** Synthesis of Diketo Acids. A mixture of 1 N NaOH (6.5 mL)
448 and the appropriate ester (1.3 mmol) in 1:1 THF/methanol (12 mL) was stirred at room
449 temperature for 40 min and then poured onto crushed ice. The aqueous layer was treated
450 with 1 N HCl until pH 3 (pH 7 for 1d) was obtained, and the solid that formed was collected
451 by filtration, then washed with water and dried under warming lamp to afford pure acids.
452 Yield (%), melting point (°C), IR, and ¹H NMR are reported for each compound.

453

454 **1-(4-Phenyl-1H-pyrrol-3-yl)propan-1-one (8)**.⁴¹ Compound 8 was prepared from (E)-1-
455 phenylpent-1-en-3-one³⁵ by means of GP-A. 79% as a yellow solid; 169–170 °C; toluene.
456 Anal. (C₁₃H₁₃NO) C, H, N.

457

458 **1-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)ethanone (9a)**. Compound 9a was
459 prepared from 1-(4-phenyl-1H-pyrrol-3-yl)ethanone³² by means of GP-B, using 4-
460 fluorobenzyl bromide as alkylating agent. Chloroform; 100% as brown oil; IR ν 1705 (C O
461 ketone) cm⁻¹; ¹H NMR (DMSO d₆) δ 2.26 (s, 3H, CH₃), 5.04 (s, 2H, CH₂), 6.63 (s, 1H,
462 pyrrole α -proton), 7.06 (m, 2H, benzyl H), 7.19 (m, 2H, benzyl H), 7.25–7.41 (m, 6H,
463 benzene H and pyrrole α -proton). Anal. (C₁₉H₁₆FNO) C, H, N, F.

464 **1-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)propan-1-one (9b)**. Compound 9b was
465 prepared from 8 by means of GP-B, using 4-fluorobenzyl bromide as alkylating agent.
466 Chloroform/acetate 50:1; 66% as brown oil; IR ν 1656 (C O ketone) cm⁻¹. ¹H NMR (DMSO
467 d₆) δ 1.04 (t, 3H, J = 8 Hz, CH₂CH₃), 2.75 (q, 2H, J = 8 Hz, CH₂CH₃), 5.18 (s, 2H, CH₂),
468 7.07 (d, 1H, J = 2.2 Hz, pyrrole C5-H), 7.2–7.3 (m, 3H, benzene H), 7.32 (t, 2H, benzyl H),
469 7.4 (m, 2H, benzene H), 7.47 (m, 2H, benzyl H), 7.87 (d, 1H, J = 2 Hz, pyrrole C2-H). Anal.
470 (C₂₀H₁₈FNO) C, H, N, F.

471 **1-(1-(4-Fluorobenzyl)-1H-pyrrol-3-yl)ethanone (9c)**. Compound 9c was prepared from
472 1-(1-(4-fluorobenzyl)-1H-pyrrol-2-yl)ethanone³⁶ by means of GP-D. 50% as brown oil; IR ν
473 1655 (C O ketone) cm⁻¹. ¹H NMR (CDCl₃) δ 2.37 (s, 3H, CH₃), 5.03 (s, 2H, CH₂),
474 6.60–6.63 (m, 2H, pyrrole C4-H and C5-H), 7.03 (t, 2H, benzene H), 7.12 (m, 2H, benzene
475 H), 7.28 (t, 1H, J = 2.0 Hz, pyrrole C2-H). Anal. (C₁₃H₁₄FNO) C, H, N, F.

476

477 **1-(4-Fluorobenzyl)-4-iodo-1H-pyrrole-2-carboxaldehyde (10).** Compound 10 was
478 prepared from commercially available 4-iodopyrrole-2-carboxaldehyde by means of GP-B,
479 using 4-fluorobenzyl bromide as alkylating agent. Chloroform/n-hexane 4:1; 39% as brown
480 oil; IR ν 1651 (C O) cm^{-1} . ^1H NMR (CDCl_3) δ 5.47 (s, 2H), 6.9– 7.0 (m, 4H, pyrrole α -
481 proton, pyrrole β -proton, and benzene H), 7.02 (m, 2H, benzene H), 9.48 (s, 1H, CHO). Anal.
482 ($\text{C}_{12}\text{H}_9\text{FINO}$) C, H, N, F, I.

483

484 **4-(4-Fluorobenzoyl)-1H-pyrrole-2-carboxaldehyde (11).** To a well stirred solution of DMF
485 (3.9 mL, 50 mmol) in 1,2-dichloroethane (100 mL) refrigerated in an ice bath was added
486 dropwise a solution of oxalyl chloride (6.35 g, 50 mmol) in 1,2 dichloroethane (100 mL) in a
487 period of 15 min. After addition, the suspension was stirred at room temperature for 15 min.
488 After this time the reaction mixture was refrigerated in ice bath and treated with a solution of
489 pyrrole (49.9 mmol) in 1,2-dichloroethane (100 mL). The mixture was stirred for 15 min at
490 room temperature and then treated with AlCl_3 (14.6 g, 109 mmol) and 4-fluorobenzoyl
491 chloride (50 mmol). The reaction was maintained at room temperature for 4 h. After this
492 period the reaction was quenched with ice and water and extracted with ethyl acetate, dried
493 over sodium sulfate, filtered, and evaporated under vacuum. The crude product was purified
494 with chromatography on aluminum oxide (1:1 ethyl acetate–chloroform as eluent) to afford
495 5 as yellow solid. 85%; 110–111 $^\circ\text{C}$; benzene/cyclohexane; IR ν 2900 (enol), 1660 (C O
496 ketone) 1640 (C O) cm^{-1} . ^1H NMR (CDCl_3) δ 7.1–7.4 (m, 3H, benzene H and pyrrole β -
497 proton), 7.45 (s, 1H, pyrrole α -proton), 7.8–7.9 (m, 2H, benzene H), 9.60 (s, 1H, CHO), 12
498 (sb, 1H, NH). Anal. ($\text{C}_{12}\text{H}_8\text{FNO}_2$) C, H, N, F.

499 **1-Benzyl-4-(4-fluorobenzoyl)-1H-pyrrole-2-carbaldehyde (12e).** Compound 12e was
500 prepared from 11 by means of GP-B, using benzyl bromide as alkylating agent. Acetate/n-
501 hexane 1:2; 34% as brown oil; IR ν 1672 (C O aldehyde), 1638 (C O ketone) cm^{-1} . ^1H NMR

502 (CDCl₃) δ 5.62 (s, 2H, CH₂), 7.1–7.2 (m, 4H, benzyl H and pyrrole β-proton), 7.3–7.4 (m,
503 4H, benzyl H and benzoyl H), 7.58 (s, 1H, pyrrole α-proton), 7.8–7.9 (m, 2H, benzoyl H),
504 9.62 (s, 1H, CHO). Anal. (C₁₉H₁₄FNO₂) C, H, N, F.

505 **4-(4-Fluorobenzoyl)-1-phenyl-1H-pyrrole-2-carbaldehyde (12f)**. Compound 11,
506 phenylboronic acid, pyridine, and copper(II) acetate anhydrous were dissolved in NMP (2.4
507 mL) in a microwave reactor tube and left to react at 60 W, 120 °C for 6 min. After this period
508 the reaction was quenched with water and extracted with ethyl acetate (5 × 20 mL), washed
509 with water (5 × 20 mL), dried over sodium sulfate, filtered, and evaporated under vacuum.
510 The crude product was purified with chromatography on silica gel (1:1 ethyl acetate–hexane
511 as eluent) to afford 12f as brown solid (63% yield). 130–131 °C. Benzene/cyclohexane; IR
512 ν 1680 (C O aldehyde), 1632 (C O ketone) cm⁻¹. ¹H NMR (CDCl₃) δ 7.21 (t, 2H, benzoyl
513 H), 7.4–7.5 (m, 2H, benzene H), 7.5–7.6 (m, 4H, benzene H and pyrrole β-proton), 7.67 (d,
514 1H, J = 2 Hz, pyrrole α-proton), 7.9–8.0 (m, 2H, benzoyl H), 9.68 (s, 1H, CHO). Anal.
515 (C₁₈H₁₂FNO₂) C, H, N, F.

516

517 **1-Methyl-4-(4-fluorobenzoyl)-1H-pyrrole-2-carboxaldehyde (12g)**. Compound 12g was
518 prepared from 11 by means of GP-B, using iodomethane as alkylating agent. Chloroform;
519 90% as gray solid; 115–116 °C; benzene/cyclohexane; IR ν 1660 (C O aldehyde), 1640 (C
520 O ketone) cm⁻¹. ¹H NMR (CDCl₃) δ 4.03 (s, 3H, N-CH₃), 7.2 (m, 2H, benzoyl H), 7.4 (d,
521 1H, pyrrole β-proton), 7.5 (s, 1H, pyrrole α-proton), 7.8–7.9 (m, 2H, benzoyl H), 9.60 (s, 1H,
522 CHO). Anal. (C₁₃H₁₀FNO₂) C, H, N, F.

523 **1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrole-2-carbaldehyde (12h)**. Compound 12h was
524 prepared from 10 by means of GP-E. 1:1 ethyl acetate–hexane as eluent; 95% as brown oil;
525 IR ν 1642 (C O aldehyde) cm⁻¹. ¹H NMR (CDCl₃) δ 5.54 (s, 2H, CH₂), 6.82 (d, 2H, J = 7.0
526 Hz, benzene H), 6.91 (t, 1H, J = 7.0 Hz, benzene H), 6.99 (t, 2H, benzyl H), 7.16–7.24 (m,

527 4H, pyrrole β -proton, pyrrole α -proton, benzyl H), 7.48 (d, 2H, $J = 7$ Hz, benzene H), 9.58
528 (s, 1H, CHO). Anal. (C₁₈H₁₄FNO) C, H, N, F.

529 **4-(1-Benzyl-4-(4-fluorobenzoyl)-1H-pyrrol-2-yl)but-3-en-2-one (13e)**. Compound 13e
530 was prepared from 12e by means of GP-C. 71% as yellow solid; 94–95 °C;
531 benzene/cyclohexane; IR ν 1675 (C O ketone), 1637 (C O ketone) cm⁻¹. ¹H NMR (CDCl₃)
532 δ 2.26 (s, 3H, CH₃), 5.29 (s, 2H, CH₂), 6.60 (d, 1H, $J = 16$ Hz, butenone C₃-H), 7.1–7.2 (m,
533 5H, butenone C₄-H, benzyl H and pyrrole β -proton), 7.3–7.4 (m, 4H, benzyl H and benzoyl
534 H), 7.46 (d, 1H, $J = 2$ Hz, pyrrole α -proton), 7.90 (m, 2H, benzoyl H). Anal. (C₂₂H₁₈FNO₂)
535 C, H, N, F.

536 **4-(1-Phenyl-4-(4-fluorobenzoyl)-1H-pyrrol-2-yl)but-3-en-2-one (13f)**. Compound 13f was
537 prepared from 12f by means of GP-C. 30% as yellow solid; 145–146 °C;
538 benzene/cyclohexane; IR ν 1680 (C O ketone), 1634 (C O ketone) cm⁻¹. ¹H NMR (CDCl₃)
539 δ 2.24 (s, 3H, CH₃), 6.56 (d, 1H, $J = 16$ Hz, butenone C₃-H), 7.19 (t, 2H, benzoyl H), 7.24
540 (d, 1H, $J = 16$ Hz, butenone C₄-H), 7.35 (d, 1H, $J = 2$ Hz, pyrrole β -proton), 7.38 (d, 1H, $J =$
541 2 Hz, pyrrole α -proton), 7.5–7.6 (m, 5H, benzene H), 7.94 (m, 2H, benzoyl H). Anal.
542 (C₂₁H₁₆FNO₂) C, H, N, F.

543 **4-(1-Methyl-4-(4-fluorobenzoyl)-1H-pyrrol-2-yl)but-3-en-2-one (13g)**. Compound 13g
544 was prepared from 12g by means of GP-C. 70% as yellow solid; 117–118 °C;
545 benzene/cyclohexane; IR ν 1660 (C O ketone), 1640 (C O ketone) cm⁻¹. ¹H NMR (CDCl₃)
546 δ 2.32 (s, 3H, CH₃), 3.78 (s, 3H, N-CH₃), 6.62 (d, 1H, $J = 16$ Hz, butenone C₃-H), 7.1–7.2
547 (m, 3H, benzene H and pyrrole β -proton), 7.35 (d, 1H, $J = 3.7$ Hz, pyrrole α -proton), 7.43 (d,
548 1H, $J = 16$ Hz, butenone C₄-H), 7.8–7.9 (m, 2H, benzene H). Anal. (C₁₆H₁₄FNO₂) C, H,
549 N, F.

550

551 **4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-2-yl)but-3-en-2-one (13h)**. Compound 13h
552 was prepared from 12h by means of GP-C. 15% as yellow solid; 160–161 °C;
553 benzene/cyclohexane; IR v 1604 (C O ketone) cm⁻¹. 1H NMR (CDCl₃) δ 2.28 (s, 3H, CH₃),
554 5.24 (s, 2H, CH₂), 6.57 (d, 1H, J = 16 Hz, butenone C3-H), 7.0–7.1 (m, 5H, pyrrole β-proton
555 and benzyl H), 7.18 (d, 1H, J = 2 Hz, pyrrole α-proton,), 7.24 (t, 1H, J = 7 Hz, benzene H),
556 7.3–7.4 (m, 3H, butenone C4-H and benzene H), 7.5–7.6 (m, 2H, benzene H). Anal.
557 (C₂₁H₁₈FNO) C, H, N, F.

558 **1-(4-Fluorobenzyl)-1H-pyrrole-2-carboxaldehyde (14)**. Compound 14 was prepared
559 from commercially available pyrrole-2-carboxaldehyde by means of GP-B, using 4-
560 fluorobenzyl bromide as alkylating agent. Chloroform; 80% as brown oil; IR v 1640 (C O)
561 cm⁻¹. 1H NMR (CDCl₃) δ 5.54 (s, 2H, CH₂), 6.30 (t, 1H, J = 4 Hz, pyrrole C4-H), 6.9–7.0
562 (m, 4H, benzene H), 7.15 (d, 1H, J = 4 Hz, pyrrole C3-H), 7.17 (d, 1H, J = 4 Hz, pyrrole C5-
563 H), 9.57 (s, 1H, CHO). Anal. (C₁₂H₁₀FNO) C, H, N, F.

564 **1-(4-Fluorobenzyl)-1H-pyrrole-3-carboxaldehyde (10)**. Compound 10 was prepared from
565 9 by means of GP-D. 55% as brown oil; IR v 1640 (C O aldehyde) cm⁻¹. 1H NMR (CDCl₃)
566 δ 5.08 (s, 2H, CH₂), 6.6 (s, 1H, pyrrole C4-H), 6.7 (s, 1H, pyrrole C-2H), 7.0–7.2 (m, 4H,
567 benzene H), 7.31 (s, 1H, pyrrole C2-H), 9.75 (s, 1H, CHO). Anal. (C₁₂H₁₀FNO) C, H, N, F.

568 **4-(4-Phenyl-1H-pyrrol-3-yl)but-3-en-2-one (16)**.⁴² Compound 16 was prepared from
569 (3E,5E)-6-phenylhexa-3,5-dien-2-one³⁸ by means of GP-A. 56% as brown solid; toluene;
570 IR v 1640 (C O ketone) cm⁻¹. Anal. (C₁₄H₁₃NO) C, H, N.

571 **4-(1-(4-Fluorobenzyl)-1H-pyrrol-3-yl)but-3-en-2-one (17i)**. Compound 17i was prepared
572 from 15 by means of GP-C. 70% as brown oil; IR v 1655 (C O ketone) cm⁻¹. 1H NMR
573 (CDCl₃) δ 2.34 (s, 3H, CH₃), 5.07 (s, 2H, CH₂), 6.43–6.48 (m, 2H, pyrrole C4-H and
574 butenone C3-H), 6.71 (s, 1H, pyrrole C2-H), 7.0 (s, 1H, pyrrole C5-H), 7.06–7.12 (m, 2H,

575 benzene H), 7.15–7.18 (m, 2H, benzene H), 7.49 (d, 1H, butenone C4-H, J = 16 Hz). Anal.
576 (C₁₅H₁₄FNO) C, H, N, F.

577 **4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)but-3-en-2-one (17j)**. Compound 17j was
578 prepared from 16 by means of GP-B, using 4-fluorobenzyl bromide as alkylating agent.
579 Chloroform; 63% as brown oil; IR ν 1640 (C O) cm^{-1} . ¹H NMR (CDCl₃) δ 2.44 (s, 3H, CH₃),
580 5.07 (s, 2H, CH₂), 6.82 (s, 1H, J = 16.3 Hz, butenone C3-H), 6.90 (s, 1H,), 6.99–7.54 (m,
581 11H, pyrrole C2-H, pyrrole C5-H, benzene H and benzyl H), 7.80 (s, 1H, J = 16.3 Hz,
582 butenone C4-H). Anal. (C₂₁H₁₈FNO) C, H, N, F.

583 **1-(1-(4-Fluorobenzyl)-4-iodo-1H-pyrrol-2-yl)ethanone (18)**. A mixture of 1-(1-(4-
584 fluorobenzyl)-1H-pyrrol-2-yl)ethanone³⁶ (4 g, 18.4 mmol) in dry acetone (100 mL) was
585 cooled at –78 °C. NIS (4.98 g, 22.1 mmol) was added. The mixture was stirred, and the
586 temperature was slowly increased to 25 °C in a period of 96 h. After this period the mixture
587 was evaporated, and ethyl acetate (50 mL) and NaHCO₃(aq) (50 mL) were added. The
588 organic phase was separated, dried over sodium sulfate, filtered, and evaporated under
589 vacuum. The raw material was purified with a column chromatography on silica gel (1:10
590 ethyl acetate–hexane as eluent) to afford 18 as white solid with a yield of 40%. 81–82 °C;
591 n-hexane; IR ν 1640 (C O ketone) cm^{-1} . ¹H NMR (CDCl₃) δ 2.40 (s, 3H, CH₃), 5.50 (s, 2H,
592 CH₂), 6.92 (d, 1H, J = 2.0 Hz, pyrrole C3-H), 7.00 (t, 2H, benzene H), 7.07 (d, 1H, J = 2.0
593 Hz, pyrrole C5-H), 7.1–7.2 (m, 2H, benzene H). Anal. (C₁₃H₁₁FINO) C, H, N, F, I.

594 **1-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-2-yl)ethanone (19)**. Compound 19 was
595 prepared from 18 by means of GP-E. 1:7 ethyl acetate–hexane as eluent. 43% as colorless
596 oil; IR ν 1650 (C O ketone) cm^{-1} . ¹H NMR (CDCl₃) δ 2.48 (s, 3H, CH₃), 5.58 (s, 2H, CH₂),
597 6.97–7.02 (t, 2H, benzyl H), 7.14–7.27 (m, 5H, benzene H, pyrrole C3-H and pyrrole C5-H),
598 7.35–7.39 (t, 2H, benzyl H), 7.51 (d, 2H, benzene H). Anal. (C₁₈H₂₁FNO) C, H, N, F.

599 **1-(1-(4-Fluorobenzyl)-1H-pyrrol-2-yl)-3-hydroxy-3-(1-trityl-1H-1,2,4-triazol-3-yl)prop-2-**
600 **en-1-one (20).** A solution of 1-[1-(4-fluorobenzyl)-1H-pyrrol-2-yl]ethanone³⁶ (1 g, 4.6 mmol)
601 in anhydrous THF (5 mL) was thermostated at $-32\text{ }^{\circ}\text{C}$. LHMDS (9.2 mL) was added, and
602 the mixture was stirred at the same temperature for 2 h. A solution of 1-trityl-1H-
603 [1,2,4]triazole-3-carboxylic acid ethyl ester³⁹ (2 g, 5.3 mmol) in anhydrous THF (18 mL) was
604 added dropwise to the solution thermostated at $-32\text{ }^{\circ}\text{C}$. After the addition, the reaction
605 mixture was stirred for 1.5 h at room temperature. The reaction was poured into 1 N HCl
606 (100 mL) and extracted with ethyl acetate. The organic phase was separated, washed with
607 water, dried over sodium sulfate, filtered, and evaporated under vacuum obtaining 2.6 g of
608 crude product as light yellow solid. The raw material was purified by recrystallization from
609 benzene/cyclohexane, obtaining 1.84 g of pure 20. 54%; $110\text{--}112\text{ }^{\circ}\text{C}$;
610 benzene/cyclohexane; IR ν 2954 (OH enol), 1626 (C O ketone) cm^{-1} . ^1H NMR (CDCl_3) δ
611 5.62 (s, 2H, CH_2), 6.3 (t, 1H, pyrrole C4-H), 6.9–7.0 (m, 4H, butenoate C3-H, pyrrole C3-H,
612 benzyl H), 7.1–7.2 (m, 9H, benzyl H, pyrrole C5-H and benzene H), 7.3–7.4 (m, 9H,
613 benzene H), 8.01 (s, 1H, triazole H), 15 (br s, 1H, OH enol). Anal. ($\text{C}_{35}\text{H}_{27}\text{FN}_4\text{O}_2$) C, H, N,
614 F.

615 **Ethyl 4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-2-hydroxy-4-oxobut-2-enoate**
616 **(6a).** Compound 6a was prepared from 9a by means of GP-F. 88%; $111\text{--}112\text{ }^{\circ}\text{C}$; benzene;
617 IR ν 2900 (OH enol), 1720 (C O ester), 1620 (C O ketone) cm^{-1} . ^1H NMR (CDCl_3) δ 1.29
618 (t, 3H, CH_3CH_2), 4.25 (q, 2H, CH_2CH_3), 5.06 (s, 2H, CH_2 benzyl), 6.48 (s, 1H, butenoate
619 C3-H), 6.67 (d, 1H, $J = 1.5\text{ Hz}$, pyrrole C5-H), 7.02–7.46 (m, 10H, pyrrole C2-H, benzene H
620 and benzyl H), 15 (br s, 1H, OH enol). Anal. ($\text{C}_{23}\text{H}_{20}\text{FNO}_4$) C, H, N, F.

621 **Ethyl 4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-2-hydroxy-3-methyl-4-oxobut-2-**
622 **enoate (6b).** Compound 6b was prepared from 9b by means of GP-F. 6b was extracted with
623 ethyl acetate. The organic phase was separated, washed with brine, dried over sodium

624 sulfate, filtered, and evaporated under vacuum obtaining a crude product that was purified
625 with column chromatography on silica gel (ethyl acetate/n-hexane 1:2). 30% as yellow oil;
626 IR ν 1730 (C O ester), 1650 (C O ketone) cm^{-1} . ^1H NMR (DMSO d_6) δ 1.14 (d, 3H, J = 7
627 Hz, CH₃), 1.34 (t, 3H, J = 7.5 Hz, CH₃CH₂), 4.33 (q, 2H, J = 7.5 Hz, CH₂CH₃), 5.25 (s, 2H,
628 CH₂), 7.14 (d, 1H, J = 1.9 Hz, pyrrole C2-H), 7.22–7.51 (m, 9H, benzene H and benzyl H),
629 8.08 (s, 1H, J = 1.9 Hz, pyrrole C5-H), 14 (br s, 1H, enol). Anal. (C₂₄H₂₂FNO₄) C, H, N, F.

630 **Ethyl 4-(1-(4-Fluorobenzyl)-1H-pyrrol-3-yl)-2-hydroxy-4-oxobut-2-enoate (6c).**

631 Compound 6c was prepared from 9c by means of GP-F. 93% as yellow solid; 63–65 °C;
632 ligroin; IR ν 3500–2500 (OH enol), 1726 (C O ester), 1633 (C O ketone) cm^{-1} . ^1H NMR
633 (DMSO d_6) δ 1.28 (t, 3H, J = 7 Hz, CH₃CH₂), 4.27 (q, 2H, J = 7 Hz, CH₂CH₃), 5.17 (s, 2H,
634 CH₂), 6.62 (t, 1H, pyrrole C4-H), 6.73 (s, 1H, butenoate C3-H), 7.02 (t, 1H, pyrrole C5-H),
635 7.18 (t, 2H, benzene H), 7.36 (dd, 2H, benzene H), 8.07 (s, 1H, pyrrole C2-H), 15 (br s, 1H,
636 enol). Anal. (C₁₇H₁₈FNO₄) C, H, N, F.

637 **Ethyl 2-Amino-4-(1-(4-fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-4-oxobut-2-enoate (6d).**

638 To a stirred mixture of 6a (1 g, 2.5 mmol) and ammonium acetate (0.22 g, 2.9 mmol) in
639 benzene (30 mL) was added acetic acid glacial (0.2 mL, 3.9 mmol). The mixture was stirred
640 at reflux for 20 h with a Dean–Stark apparatus. After this period the mixture was cooled to
641 room temp and washed with a saturated solution of NaHCO₃ (50 mL). The organic layer
642 was separated, dried over sodium sulfate, filtered, and evaporated under vacuum. The raw
643 material was purified with chromatography on aluminum oxide (chloroform as eluent) to
644 afford 6d as yellow solid with a yield of 50%. 135–136 °C. benzene; IR ν 3500 (NH), 1720
645 (C O ester), 1620 (C O ketone) cm^{-1} . ^1H NMR (CDCl₃) δ 1.27 (t, 3H, CH₃CH₂), 4.25 (q,
646 2H, CH₃CH₂), 5.08 (s, 2H, CH₂), 6.13 (s, 1H, butenoate C3-H), 6.67 (d, 1H, J = 1.5 Hz,
647 pyrrole C5-H), 7.0–7.6 (m, 10H, benzene H, benzyl H, and pyrrole C2-H), 9 (br s, 2H, NH₂).
648 Anal. (C₂₃H₂₁FN₂O₂) C, H, N, F.

649 **Ethyl 6-(1-Benzyl-4-(4-fluorobenzoyl)-1H-pyrrol-2-yl)-2-hydroxy-4-oxohexa-2,5-**
650 **dienoate (6e).** Compound 6e was prepared from 13e by means of GP-F. 62% as yellow
651 solid; 154–155 °C; benzene/cyclohexane; IR ν 1730 (C O ester), 1636 (C O ketone) cm^{-1} .
652 ^1H NMR (acetone- d_6) δ 1.34 (t, 3H, CH_3CH_2), 4.32 (q, 2H, CH_2CH_3), 5.60 (s, 2H, CH_2),
653 6.86 (d, 1H, $J = 15.6$ Hz, hexanoate C5-H), 7.2–7.5 (m, 9H, $J = 1.6$ Hz, benzene H, pyrrole
654 β -proton, benzoyl H, and hexanoate C3-H), 7.72 (d, 1H, $J = 15.6$ Hz, hexanoate C6-H), 7.90
655 (d, 1H, $J = 1.6$ Hz, pyrrole α -proton), 7.9–8.0 (m, 2H, benzoyl H), 14 (bs, 1H, enol). Anal.
656 ($\text{C}_{26}\text{H}_{22}\text{FNO}_5$) C, H, N, F.

657 **Ethyl 6-(4-(4-Fluorobenzoyl)-1-phenyl-1H-pyrrol-2-yl)-2-hydroxy-4-oxohexa-2,5-**
658 **dienoate (6f).** Compound 6f was prepared from 13f by means of GP-F. 80% as yellow solid;
659 129–130 °C; benzene/cyclohexane; IR ν 3500 (OH enol), 1720 (C O ester), 159 (C O
660 ketone) cm^{-1} . ^1H NMR (CD_3OD) δ 1.36 (t, 3H, $J = 7.5$ Hz, CH_3CH_2), 4.31 (q, 2H, $J = 7.5$
661 Hz, CH_2CH_3), 7.26–7.48 (m, 3H, benzoyl H and hexanoate C5-H), 7.39 (d, 1H, $J = 15.6$ Hz,
662 hexanoate C6-H), 7.43–7.48 (m, 3H, benzene H and hexanoate C3-H), 7.58–7.62 (m, 4H,
663 benzene H and pyrrole β -proton), 7.72 (s, 1H, pyrrole α -proton), 7.97–8.01 (m, 2H, benzoyl
664 H), 14 (bs, 1H, OH enol). Anal. ($\text{C}_{25}\text{H}_{20}\text{FNO}_5$) C, H, N, F.

665 **Ethyl 6-(4-(4-Fluorobenzoyl)-1-methyl-1H-pyrrol-2-yl)-2-hydroxy-4-oxohexa-2,5-**
666 **dienoate (6g).** Compound 6g was prepared from 13g by means of GP-F. 83% as yellow
667 solid; 166–167 °C; benzene/cyclohexane; IR ν 1 2900 (OH enol), 1720 (C O ester), 1650 (C
668 O ketone) cm^{-1} . ^1H NMR (CDCl_3) δ 1.39 (t, 3H, $J = 7.5$ Hz, CH_3CH_2), 3.81 (s, 3H, N- CH_3),
669 4.39 (q, 2H, $J = 7.5$ Hz, CH_2CH_3), 6.45 (s, 1H, hexanoate C3-H), 6.50 (d, 1H, $J = 15.4$ Hz,
670 hexanoate C5-H), 7.12–7.20 (m, 3H, benzene H and pyrrole β -proton), 7.36–39 (m, 1H,
671 pyrrole α -proton), 7.63 (d, 1H, $J = 15.4$ Hz, hexanoate C6-H), 7.83–7.91 (m, 2H, benzene
672 H), 15 (bs, 1H, enol). Anal. ($\text{C}_{20}\text{H}_{18}\text{FNO}_5$) C, H, N, F.

673 **Ethyl 6-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-2-yl)-2-hydroxy-4-oxohexa-2,5-**
674 **dienoate (6h).** Compound 6h was prepared from 13h by means of GP-F. 92% as red solid;
675 169–170 °C; benzene; IR v 2900 (OH enol), 1723 (C O ester), 1602 (C O ketone) cm⁻¹. ¹H
676 NMR (acetone-d₆) δ 1.35 (t, 3H, J = 7.5 Hz, CH₃CH₂), 4.32 (q, 2H, J = 7.5 Hz, CH₂CH₃),
677 5.53 (s, 2H, CH₂), 6.46 (s, 1H, hexanoate C3-H), 6.74 (d, 1H, J = 16 Hz, hexanoate C5-H),
678 7.13–7.24 (m, 3H, benzyl H and benzene H), 7.30–39 (m, 4H, benzyl H, hexanoate C6-H,
679 and pyrrole β-proton), 7.43 (s, 1H, pyrrole α-proton), 7.65 (d, 2H, benzene H), 7.7–7.8 (m,
680 2H, benzene H), 15 (bs, 1H, enol). Anal. (C₂₅H₂₂FNO₄) C, H, N, F.

681 **Ester 6-(1-(4-Fluorobenzyl)-1H-pyrrol-3-yl)-2-hydroxy-4-oxohexa-2,5-dienoate (6i).**
682 Compound 6i was prepared from 17i by means of GP-F. 41% as yellow solid; 88–90 °C;
683 ligroin; IR v 3400 (OH enol), 1725 (C O ester), 1625 (C O ketone) cm⁻¹. ¹H NMR (CDCl₃)
684 δ 1.44 (t, 3H, J = 7 Hz, CH₃CH₂), 4.41 (q, 2H, J = 7 Hz, CH₂CH₃), 5.08 (s, 2H, CH₂), 6.39
685 (s, 1H, J = 16 Hz, hexanoate C5-H), 6.5–6.6 (m, 2H, pyrrole C4-H and hexanoate C3-H),
686 6.7 (t, 1H, pyrrole C5-H), 7.0 (t, 1H, pyrrole C2-H), 7.08–7.20 (m, 4H, benzene H), 7.75 (s,
687 1H, J = 16 Hz, hexanoate C6-H), 14 (br s, 1H, enol). Anal. (C₁₉H₁₈FNO₄) C, H, N, F.

688 **Ethyl 6-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-2-hydroxy-4-oxohexa-2,5-**
689 **dienoate (6j).** Compound 6j was prepared from 17j by means of GP-F. 56% as yellow solid;
690 121–122 °C; cyclohexane; IR v 2900 (OH enol), 1720 (C O ester), 1620 (C O ketone) cm⁻¹.
691 ¹H NMR (CDCl₃) δ 1.43 (t, 3H, CH₃CH₂), 4.40 (q, 2H, CH₂CH₃), 5.09 (s, 2H, CH₂), 6.73
692 (s, 1H, hexanoate C3-H), 6.87 (d, 1H, J = 16.6 Hz, hexanoate C5-H), 7.01 (s, 1H, pyrrole
693 C5-H), 7.08–7.54 (m, 10H, pyrrole C2-H, benzene H and benzyl H), 7.74 (d, 1H, hexanoate
694 C6-H), 15 (bs, 1H, enol). Anal. (C₂₅H₂₁FNO₄) C, H, N, F.

695 **Ethyl 4-(1-(4-Fluorobenzyl)-1H-pyrrol-2-yl)-2-hydroxy-4-oxobut-2-enoate (6k).**
696 Compound 6k was prepared from 1-(1-(4-fluorobenzyl)-1H-pyrrol-2-yl)ethanone³⁶ by
697 means of GP-F. 41% as yellow solid; 88–89 °C; ligroin; IR v 2900 (OH enol), 1720 (C O

698 ester), 1620 (C O ketone) cm^{-1} . $^1\text{H NMR}$ (DMSO d_6) δ 1.26 (t, 3H, $J = 7.5$ Hz, CH_3CH_2),
699 4.25 (q, 2H, $J = 7.5$ Hz, CH_2CH_3), 5.59 (s, 2H, CH₂), 6.32 (dd, 1H, $J = 2.5$ Hz, $J = 3.5$ Hz,
700 pyrrole C4-H), 6.84 (s, 1H, butenoate C3-H), 7.0–7.2 (m, 4H, benzene H), 7.43 (d, 1H, $J =$
701 3.5 Hz, pyrrole C3-H), 7.53 (s, 1H, pyrrole C5-H), 14 (br s, 1H, enol). Anal. ($\text{C}_{17}\text{H}_{16}\text{FNO}_4$)
702 C, H, N, F.

703 **Ethyl 4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-2-yl)-2-hydroxy-4-oxobut-2-enoate**
704 **(6l)**. Compound 6l was prepared from 19 by means of GP-F. 95% as yellow solid; 102–103
705 °C; benzene/ cyclohexane; IR ν 3400 (OH enol), 1720 (C O ester), 1620 (C O ketone) cm^{-1} .
706 $^1\text{H NMR}$ (DMSO d_6) δ 1.34 (t, 3H, $J = 7.5$ Hz, CH_3CH_2), 4.33 (q, 2H, $J = 7.5$ Hz, CH_2CH_3),
707 5.68 (s, 2H, CH₂), 7.06 (s, 1H, butenoate C3-H), 7.17–7.31 (m, 5H, benzene H and benzyl
708 H), 7.40–7.44 (m, 2H, benzene H), 7.75 (d, 2H, benzene H), 8.01 (s, 1H, pyrrole C5-H), 8.13
709 (s, 1H, pyrrole C3-H), 14 (bs, 1H, enol). Anal. ($\text{C}_{22}\text{H}_{20}\text{FNO}_4$) C, H, N, F.

710 **4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-2-hydroxy-4-oxobut-2-enoic Acid (7a)**.
711 Compound 7a was prepared from 6a by means of GP-G. 98% as brown solid; 109–110 °C;
712 toluene; IR ν 3500–2000 (OH acid and enol), 1740 (C O acid), 1620 (C O ketone) cm^{-1} . ^1H
713 NMR (CDCl_3) δ 5.06 (s, 2H, CH₂), 6.48 (s, 1H, butenoate C3-H), 6.67 (d, 1H, $J = 1.5$ Hz,
714 pyrrole C5-H), 7.06–7.10 (t, 2H, benzyl H), 7.18–7.25 (m, 2H, benzyl H), 7.30–7.39 (m, 5H,
715 benzene H), 7.49 (s, 1H, pyrrole C2-H), 14 (br s, 2H, OH acid and enol). Anal.
716 ($\text{C}_{21}\text{H}_{16}\text{FNO}_4$) C, H, N, F.

717 **4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-2-hydroxy-3-methyl-4-oxobut-2-enoic**
718 **Acid (7b)**. Compound 7b was prepared from 6b by means of GP-G. 38% as white solid;
719 233–234 °C; benzene; IR ν 3500–2500 (OH acid and enol), 1700 (C O acid), 1640 (C O
720 ketone) cm^{-1} . $^1\text{H NMR}$ (DMSO d_6) δ 1.14 (d, 3H, CH₃, $J = 7$ Hz), 5.13 (s, 2H, CH₂), 7.1–7.2
721 (m, 3H, benzyl H and pyrrole C5-H), 7.34 (t, 2H, benzyl H), 7.47 (s, 1H, pyrrole C2-H), 7.51

722 (m, 3H, benzene H), 7.9–8.0 (m, 2H, benzene H), 14 (br s, 2H, enol and acid). Anal.
723 (C₂₂H₁₈FNO₄) C, H, N, F.

724 **4-(1-(4-Fluorobenzyl)-1H-pyrrol-3-yl)-2-hydroxy-4-oxobut-2-enoic Acid (7c).**

725 Compound 7c was prepared from 6c by means of GP-G. 57% as yellow solid; 146–147 °C;
726 toluene; IR ν 3500–2500 (OH acid and enol), 1727 (C O acid), 1621 (C O ketone) cm⁻¹. ¹H
727 NMR (DMSO-d₆) δ 5.16 (s, 2H, CH₂), 6.60 (dd, 1H, J₁ = 1.5 Hz, J₂ = 3.0 Hz, pyrrole C4-
728 H), 6.72 (s, 1H, butenoate C3-H), 7.00 (dd, 1H, J₁ = 1.5 Hz, J₂ = 3 Hz, pyrrole C5-H), 7.18
729 (t, 2H, benzene H), 7.36 (dd, 2H, benzene H), 8.04 (s, 1H, pyrrole C2-H), 14 (br s, 1H, enol),
730 15 (br s, 1H, COOH). Anal. (C₁₅H₁₂FNO₄) C, H, N, F.

731 **6-(1-Benzyl-4-(4-fluorobenzoyl)-1H-pyrrol-2-yl)-2-hydroxy-4-oxohexa-2,5-dienoic Acid**

732 **(7e).** Compound 7e was prepared from 6e by means of GP-G. 87% as orange solid;
733 165–166 °C; benzene; IR ν 3500–2500 (OH acid and enol), 1727 (C O acid), 1630 (C O
734 ketone) cm⁻¹. ¹H NMR (CD₃OD) δ 5.22 (s, 2H, CH₂), 6.69 (s, 1H, J = 15.2 Hz, hexanoate
735 C5-H), 7.15–7.44 (m, 9H, hexanoate C3-H, benzyl H, benzoyl H, and pyrrole β -proton), 7.65
736 (s, 1H, J = 15.2 Hz, hexanoate C6-H), 7.79 (s, 1H, pyrrole α -proton), 7.92–7.97 (m, 2H,
737 benzoyl H). Anal. (C₂₄H₁₈FNO₅) C, H, N, F.

738 **6-(4-(4-Fluorobenzoyl)-1-phenyl-1H-pyrrol-2-yl)-2-hydroxy-4-oxohexa-2,5-dienoic**

739 **Acid (7f).** Compound 7f was prepared from 6f by means of GP-G. 85% as yellow solid;
740 183–184 °C; benzene; IR ν 3500–2500 (OH acid and enol), 1724 (C O acid), 1596 (C O
741 ketone) cm⁻¹. ¹H NMR (CD₃OD) δ 6.69 (s, 1H, J = 16 Hz, hexanoate C5-H), 7.26 (t, 2H,
742 benzoyl H), 7.41 (d, 1H, hexanoate C6-H), 7.47– 7.49 (m, 2H, pyrrole β -proton and
743 hexanoate C3-H), 7.73 (s, 1H, pyrrole α -proton), 7.98–8.01 (m, 2H, benzoyl H). Anal.
744 (C₂₃H₁₆FNO₅) C, H, N, F.

745 **6-(4-(4-Fluorobenzoyl)-1-methyl-1H-pyrrol-2-yl)-2-hydroxy- 4-oxohexa-2,5-dienoic**

746 **Acid (7g).** Compound 7g was prepared from 6g by means of GP-G. 95% as orange solid;

747 162–163 °C; ethanol; IR ν 3500–2500 (OH acid and enol), 1700 (C O acid), 1600 (C O
748 ketone) cm^{-1} . ^1H NMR (DMSO d_6) δ 3.32 (s, 3H, N-CH₃), 6.52 (s, 1H, hexanoate C3-H),
749 6.93 (d, 1H, hexanoate C5-H), 7.32–7.86 (m, 7H, benzene H, pyrrole β -proton, pyrrole α -
750 proton, and hexanoate C6-H), 14 (br s, 2H, OH enol and acid). Anal. (C₁₈H₁₄FNO₅) C, H,
751 N, F.

752 **6-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-2-yl)-2-hydroxy-4-oxohexa-2,5-dienoic Acid**
753 **(7h)**. Compound 7h was prepared from 6h by means of GP-G. 73% as red solid; decompose;
754 benzene; IR ν 3500–2500 (OH acid and enol), 1707 (C O acid), 1571 (C O ketone) cm^{-1} .
755 ^1H NMR (acetone- d_6) δ 5.51 (s, 2H, CH₂), 6.49 (s, 1H, hexanoate C3-H), 6.7, (d, 1H,
756 hexanoate C5-H), 7.07–7.42 (m, 7H, benzyl H, pyrrole β -proton, pyrrole α -proton, and
757 hexanoate C6-H), 7.6–7.8 (m, 5H, benzene H), 14 (br s, 2H, enol and acid). Anal.
758 (C₂₃H₁₈FNO₄) C, H, N, F.

759 **6-(1-(4-Fluorobenzyl)-1H-pyrrol-3-yl)-2-hydroxy-4-oxohexa-2,5-dienoic Acid (7i)**.
760 Compound 7i was prepared from 6i by means of GP-G. 92% as yellow solid; >300 °C;
761 DMF/H₂O; IR ν 3500–2500 cm^{-1} (OH acid and enol), 1720 (C O acid), 1630 (C O ketone)
762 cm^{-1} . ^1H NMR (DMF- d_7) δ 5.34 (s, 2H, benzyl), 6.39 (s, 1H, hexanoate C3-H), 6.54 (d, 1H,
763 hexanoate C5-H), 6.7 (bs, 1H, pyrrole C4-H), 7.38–7.67 (m, 7H, pyrrole C2-H, pyrrole C5-
764 H, benzene H, and hexanoate C6-H), 14 (br s, 2H, OH enol and acid). Anal. (C₁₇H₁₆FNO₄)
765 C, H, N, F.

766 **6-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-2-hydroxy-4-oxohexa-2,5-dienoic Acid**
767 **(7j)**. Compound 7j was prepared from 6j by means of GP-G. 75% as yellow solid; 128–129
768 °C; benzene; IR ν 3500–2500 (OH acid and enol), 1700 (C O acid), 1600 (C O ketone)
769 cm^{-1} . ^1H NMR (CDCl₃) δ 5.20 (s, 2H, CH₂), 6.75 (bs, 1H, hexanoate C3-H), 6.80 (d, 1H,
770 hexanoate C5-H), 7.1–7.6 (m, 12H, pyrrole C2-H, pyrrole C5-H, benzene H, benzyl H, and
771 hexanoate C6-H), 14 (br s, 2H, enol and acid). Anal. (C₂₃H₁₇FNO₄) C, H, N, F.

772 **4-(1-(4-Fluorobenzyl)-1H-pyrrol-2-yl)-2-hydroxy-4-oxobut-2-enoic Acid (7k).**

773 Compound 7k was prepared from 6k by means of GP-G 80% as yellow solid; 156–157 °C;
774 benzene; IR ν 3500–2500 (OH acid and enol), 1700 (C=O acid), 1620 (C=O ketone) cm^{-1} .

775 ^1H NMR (DMSO- d_6) δ 5.59 (s, 2H, CH₂), 6.30 (dd, 1H, J = 2.5 Hz, J = 3.5 Hz, pyrrole C4-
776 H), 6.81 (s, 1H, butenoate C3-H), 7.0–7.09– 7.16 (m, 4H, benzene H), 7.39 (d, 1H, J = 3.5
777 Hz, pyrrole C3-H), 7.51 (s, 1H, pyrrole C5-H) 14 (br s, 2H, OH enol and acid). Anal.
778 (C₁₅H₁₂FNO₄) C, H, N, F.

779 **4-(1-(4-Fluorobenzyl)-4-phenyl-1H-pyrrol-2-yl)-2-hydroxy-4-oxobut-2-enoic Acid (7l).**

780 Compound 7l was prepared from 6l by means of GP-G. 82% as yellow solid; 195–196 °C;
781 toluene; IR ν 3500–2000 (OH acid and enol), 1740 (C=O acid), 1620 (C=O ketone) cm^{-1} .

782 ^1H NMR (DMSO d_6) δ 5.68 (s, 2H, CH₂ benzyl), 7.05 (s, 1H, butenoate C3-H), 7.1–7.3 (m,
783 5H, benzyl H and benzene H), 7.41 (t, 2H, benzene H), 7.76 (d, 2H, benzene H), 8.01 (s,
784 1H, pyrrole C3-H), 8.13 (s, 1H, pyrrole C5-H) 14 (br s, 1H, OH enol), 15 (bs, 2H, OH acid).
785 Anal. (C₂₀H₁₆FNO₄) C, H, N, F.

786 **(E/Z)-2-Amino-4-(1-(4-fluorobenzyl)-4-phenyl-1H-pyrrol-3-yl)-4-oxobut-2-enoic Acid**

787 **(7d).** Compound 6d (340 mg, 0.9 mmol) was dissolved in anhydrous THF (4.5 mL) under
788 argon atmosphere and cooled to 0 °C. To this was added dropwise a solution of 0.5 N KOH
789 (2 mL, 1.0 mmol), and the mixture was allowed to stir at room temperature overnight. The
790 reaction mixture was concentrated in vacuo and the residue partitioned between water and
791 ethyl acetate. The aqueous layer was cooled on ice and acidified with 1 N HCl. After chilling
792 at 4 °C for 2 h, the resulting precipitate was extracted with ethyl acetate. The organic layer
793 was washed with brine, dried over Na₂SO₄, and concentrated under vacuum to yield the
794 acid 7d (250 mg). 76% yellow solid; 86–88 °C; toluene; IR ν 3000–2500 (OH acid and enol),
795 1700 (C=O acid) cm^{-1} . ^1H NMR (CDCl₃) δ 4.95 (m, 2H, CH₂ E/Z form), 6.70 (s, 1H,

796 butenoate C3-H), 6.8–7.43 (m, 11H, pyrrole C2-H, pyrrole C5-H, benzene H, and benzyl H),
797 14 (br s, 2H, acid and enol). Anal. (C₂₁H₁₇FN₂O₃) C, H, N, F.

798 **1-(1-(4-Fluorobenzyl)-1H-pyrrol-2-yl)-3-hydroxy-3-(1H-1,2,4-triazol-3-yl)prop-2-en-1-**
799 **one (7m)**. 20 (1.84 g, 1.8 mmol) was suspended in 12 mL of dioxane and treated with 4.4
800 mL of 1 N HCl. The reaction mixture was stirred at 70 °C for 4 h. After cooling, the mixture
801 was poured into 4.4 mL of 1.5 N NaOH. The formed precipitate was filtered and portioned
802 between ethyl acetate and 1 N NaOH. The water phase was separated, and acidification
803 until pH 4 was obtained was done with concentrated HCl. The formed solid was filtered,
804 washed with water, and recrystallized from absolute ethanol, obtaining 430 mg of pure 7m.
805 64%; 188–189 °C; ethanol; IR ν 3200–2400 (NH, OH enol, and acid), 1712 (C=O ketone)
806 cm⁻¹. ¹H NMR (DMSO-d₆) δ 4.50 (s, 1H, butanoate C3-H), 5.47 and 5.62 (s, 2H, CH₂ keto
807 and enol form), 6.21 and 6.29 (t, 1H, pyrrole C4-H), 6.91 (s, 1H, butenoate C3-H), 7.06–7.16
808 (m, 6H, benzene H keto and enol form), 7.22 and 7.29 (m, 1H, pyrrole C3-H), 7.36 and 7.45
809 (s, 1H, pyrrole C2-H keto and enol form), 8.59 (bs, 1H, NH), 14 (br s, 1H, OH enol and acid).
810 Anal. (C₁₆H₁₃FN₄O₂) C, H, N, F.

811 **Biological Methods.** *RT Expression and Purification.* The recombinant HIV-1 RT protein,
812 whose coding gene was subcloned in the p6HRT_prot plasmid, was expressed in E. coli
813 strain M15.43,44 The bacteria cells were grown up to an OD₆₀₀ of 0.8 and induced with 1.7
814 mM IPTG for 5 h. HIV-1 RT purification was carried out as described. Briefly, cell pellets
815 were resuspended in lyses buffer (20 mM Hepes, pH 7.5, 0.5 M NaCl, 5 mM β -
816 mercaptoethanol, 5 mM imidazole, 0.4 mg/mL lysozyme), incubated on ice for 20 min,
817 sonicated, and centrifuged at 30 000g for 1 h. The supernatant was applied to a His-binding
818 resin column and washed thoroughly with wash buffer (20 mM Hepes, pH 7.5, 0.3 M NaCl,
819 5 mM β -mercaptoethanol, 60 mM imidazole, 10% glycerol). The RT protein was eluted with

820 elute buffer. The enzyme-containing fractions were pooled, dialyzed, and aliquots were
821 stored at $-80\text{ }^{\circ}\text{C}$.

822 *HIV-1 RT RNase H Inhibition.* The RT-associated RNase H function was measured in a
823 polymerase-independent cleavage assay, in which the poly(dC)-[^3H]poly(rG) hybrid was
824 used as reaction substrate as previously described.⁴³

825 *HIV-1 IN Inhibition.* HIV-1 IN gel-based assays were carried out as previously published.⁴⁵

826 *HIV-1 Replication Inhibition.* The antiviral activity of compounds was determined in a cell-
827 based assay according to the procedure described previously⁴⁶ and modified as follows.
828 HeLa-CD4-LTR- β -gal cells were maintained in DMEM with 10% serum and 0.5 mg/mL
829 G418. The day prior to experimentation, 96-well plates were prepared to contain 10 000
830 cells per well in 100 μL of Dulbecco's modified Eagle medium (DMEM) complemented with
831 10% serum. On day 1, each drug was serial diluted directly on cells following a 3-fold dilution
832 over 6 points, and each well was then filled to 200 μL with either fresh medium or
833 concentrated viral supernatant (HIV-1(IIIB), Advanced Biotechnologies Inc.). The highest
834 compound concentration tested was 50 μM . On day 2, cells were washed three times with
835 PBS before adding 200 μL of a solution containing 50 mM Tris-HCl, pH 7.5, 100 mM β -
836 mercaptoethanol, 0.05% Triton X100, and 5 mM 4-methyl-umbelliferyl- β -D-
837 galactopyranoside (4-MUG, Sigma). On day 3, sealed plates were read in a SpectraMax
838 GEMINI-XS (Molecular Devices) with $\lambda_{\text{ex}} = 360\text{ nm}$ and $\lambda_{\text{em}} = 460\text{ nm}$.

839 *Cellular Toxicity.* Similar to the antiviral assays, plates were prepared with 10 000 HeLa-
840 CD4-LTR- β -gal cells per well and a serial dilution of compounds in 100 μL . After 24 h of
841 culture, 100 μL of ATPlite reagent (PerkinElmer) was added to each well. After 5 min at room
842 temperature, the plates' luminescence was quantified using an EnVision multilabel reader
843 (PerkinElmer) according to the manufacturer's instructions.

844 **ASSOCIATED CONTENT**

845 *S Supporting Information

846 Elemental analysis results of compounds 6a–l and 7a–m. This material is available free of
847 charge via the Internet at <http://pubs.acs.org>.

848

849 **ACKNOWLEDGMENTS**

850 The authors thank the Italian MIUR for financial support, ISS 40H4, PRIN 2010–2011 (Grant
851 2010W2KM5L_002). R.D.S. and R.C. thank the FP7 CHAARM project for support. This work
852 was also supported by the NIH Intramural Research Program, Center for Cancer Research,
853 National Cancer Institute.

854

855 **ACKNOWLEDGMENTS**

856 The authors thank the Italian MIUR for financial support, ISS 40H4, PRIN 2010–2011 (Grant
857 2010W2KM5L_002). R.D.S. and R.C. thank the FP7 CHAARM project for support. This work
858 was also supported by the NIH Intramural Research Program, Center for Cancer Research,
859 National Cancer Institute.

860

861

862

863

864

865

866 **REFERENCES**

- 867 (1) Patyar, S.; Prakash, A.; Medhi, B. Dual inhibition: a novel promising pharmacological
868 approach for different disease conditions. *J. Pharm. Pharmacol.* 2011, 63, 459–471.
- 869 (2) Mellors, J. W.; Dutschman, G. E.; Im, G. J.; Tramontano, E.; Winkler, S. R.; Cheng,
870 Y.-C. In vitro selection and molecular characterization of human immunodeficiency virus-1
871 resistant to non-nucleoside inhibitors of reverse transcriptase. *Mol. Pharmacol.* 1992, 41,
872 446–451.
- 873 (3) Mellors, J.; Im, G. J.; Tramontano, E.; Winkler, S. R.; Medina, D. J.; Dutschman, G.
874 E.; Bazmi, H. Z.; Piras, G.; Gonzales, C. J.; Cheng, Y.-C. A single conservative amino acid
875 substitution in the reverse transcriptase of human immunodeficiency virus type 1 confers
876 resistance to (+)-(5S)-4,5,6,7-tetrahydro-5-methyl-6-(3-methyl-
877 2butenyl)imidazo[4,5,1,jk][1,4]benzodiazepin-2(1H)-thione (TIBO R82150). *Mol.*
878 *Pharmacol.* 1993, 43, 11–16.
- 879 (4) Mehellou, Y.; De Clercq, E. Twenty-six years of anti-HIV drug discovery: Where do
880 we stand and where do we go? *J. Med. Chem.* 2010, 53, 521–538.
- 881 (5) Brühlmann, C.; Ooms, F.; Carrupt, P. A.; Testa, B.; Catto, M.; Leonetti, F.; Altomare,
882 C.; Carotti, A. Coumarins derivatives as dual inhibitors of acetylcholinesterase and
883 monoamine oxidase. *J. Med. Chem.* 2001, 44, 3195–3198.
- 884 (6) Pretorius, J.; Malan, S. F.; Castagnoli, N., Jr.; Bergh, J. J.; Petzer, J. P. Dual inhibition
885 of monoamine oxidase B and antagonism of the adenosine A(2A) receptor by (E,E)-8-(4-
886 phenylbutadien-1-yl)caffeine analogues. *Bioorg. Med. Chem.* 2008, 16, 8676–8684.
- 887 (7) Altavilla, D.; Squadrito, F.; Bitto, A.; Polito, F.; Burnett, B. P.; Di Stefano, V.; Minutoli,
888 L. Flavocoxid, a dual inhibitor of cyclooxygenase and 5-lipoxygenase, blunts pro-

889 inflammatory phenotype activation in endotoxin-stimulated macrophages. *Br. J. Pharmacol.*
890 2009, 157, 1410– 1418.

891 (8) Park, S.; Chapuis, N.; Bardet, V.; Tamburini, J.; Gallay, N.; Willems, L.; Knight, Z. A.;
892 Shokat, K. M.; Azar, N.; Viguie, F.; Ifrah, N.; Dreyfus, F.; Mayeux, P.; Lacombe, C.; Bouscary,
893 D. PI-103, a dual inhibitor of class IA phosphatidylinositide 3-kinase and mTOR, has
894 antileukemic activity in AML. *Leukemia* 2008, 22, 1698–1706.

895 (9) Li, T.; Wang, J.; Wang, X.; Yang, N.; Chen, S. M.; Tong, L. J.; Yang, C. H.; Meng, L.
896 H.; Ding, J. WJD008, a dual PI3K/mTOR inhibitor, prevents PI3K signaling and inhibits the
897 proliferation of transformed cells with oncogenic PI3K mutant. *J. Pharmacol. Exp. Ther.*
898 2010, 334, 830–838.

899 (10) Wang, Q. M.; Johnson, R. B.; Jungheim, L. N.; Cohen, J. D.; Villarreal, E. C. Dual
900 inhibition of human rhinovirus 2A and 3C proteases by homophthalimides. *Antimicrob.*
901 *Agents Chemother.* 1998, 42, 916–920.

902 (11) Didierjean, J.; Isel, C.; Querre, F.; Mouscadet, J. F.; Aubertin, A. M.; Valnot, J. Y.;
903 Piettre, S. R.; Marquet, R. Inhibition of human immunodeficiency virus type 1 reverse
904 transcriptase, RNase H, and integrase activities by hydroxytropolones. *Antimicrob. Agents*
905 *Chemo-ther.* 2005, 49, 4884–4894.

906 (12) Budihas, S.; Gorshkova, I.; Gaidamakov, S.; Wamiru, A.; Bona, M.; Parniak, K.;
907 Crouch, R.; McMahon, J.; Beutler, J.; Le Grice, S. Selective inhibition of HIV-1 reverse
908 transcriptase-associated ribonuclease H activity by hydroxylated tropolones. *Nucleic Acids*
909 *Res.* 2005, 33, 1249–1256.

910 (13) Semenova, E. A.; Johnson, A. A.; Marchand, C.; Davis, D. A.; Yarchoan, R.;
911 Pommier, Y. Preferential inhibition of the magnesium dependent strand transfer reaction of
912 HIV-1 integrase by α -hydroxytropolones. *Mol. Pharmacol.* 2006, 69, 1454–1460.

- 913 (14) Marchand, C.; Beutler, J. A.; Wamiru, A.; Budihis, S.; Möllmann, U.; Heinisch, L.;
914 Mellors, J. W.; Le Grice, S. F.; Pommier, Y. Madurahydroxylactone derivatives as dual
915 inhibitors of human immunodeficiency virus type 1 integrase and RNase H. *Antimicrob.*
916 *Agents Chemother.* 2008, 52, 361–364.
- 917 (15) Billamboz, M.; Bailly, F.; Barreca, M. L.; De Luca, L.; Mouscadet, J. F.; Calmels, C.;
918 Andreola, M. L.; Witvrouw, M.; Christ, F.; Debyser, Z.; Cotelle, P. Design, synthesis, and
919 biological evaluation of a series of 2-hydroxyquinoline-1,3(2H,4H)-diones as dual inhibitors
920 of human immunodeficiency virus type 1 integrase and reverse transcriptase RNase H
921 domain. *J. Med. Chem.* 2008, 51, 7717–7730.
- 922 (16) Billamboz, M.; Bailly, F.; Lion, C.; Calmels, C.; Andreola, M. L.; Witvrouw, M.; Christ,
923 F.; Debyser, Z.; De Luca, L.; Chimirri, A.; Cotelle, P. 2-Hydroxyisoquinoline-1,3(2H,4H)-
924 diones as inhibitors of HIV-1 integrase and reverse transcriptase RNase H domain: influence
925 of the alkylation of position 4. *Eur. J. Med. Chem.* 2011, 46, 535–546.
- 926 (17) Cherepanov, P.; Maertens, G. N.; Hare, S. Structural insights into the retroviral DNA
927 integration apparatus. *Curr. Opin. Struct. Biol.* 2011, 21, 249–256.
- 928 (18) Beare, K. D.; Coster, M. J.; Rutledge, P. J. Diketoacid inhibitors of HIV-1 integrase:
929 from L-708,906 to raltegravir and beyond. *Curr. Med. Chem.* 2012, 19, 1177–11792.
- 930 (19) Cocohoba, J.; Dong, B. J. Raltegravir: the first HIV integrase inhibitor. *Clin. Ther.*
931 2008, 30, 1747–1765.
- 932 (20) (a) Kirschberg, T.; Parrish, J. Metal chelators as antiviral agents. *Curr. Opin. Drug*
933 *Discovery Dev.* 2007, 10, 460–472. (b) Rogolino, D.; Carcelli, M.; Sechi, M.; Neamati, N.
934 Viral enzymes containing magnesium: metal binding as a successful strategy in drug design.
935 *Coord. Chem. Rev.* 2012, 256, 3063–3086. (c) Rouffet, M.; Cohen, S. M. Emerging trends
936 in metalloprotein inhibition. *Dalton Trans.* 2011, 40, 3445–3454.

- 937 (21) Shaw-Reid, C. A.; Munshi, V.; Graham, P.; Wolfe, A.; Witmer, M.; Danzeisen, R.;
938 Olsen, D. B.; Carroll, S. S.; Embrey, M.; Wai, J. S.; Miller, M. D.; Cole, J. L.; Hazuda, D. J.
939 Inhibition of HIV-1 ribonuclease H by a novel diketo acid, 4-[5-(benzoylamino)thien-2-yl]-2,4-
940 dioxobutanoic acid. *J. Biol. Chem.* 2003, 278, 2777–2780.
- 941 (22) Tramontano, E.; Esposito, F.; Badas, R.; Di Santo, R.; Costi, R.; La Colla, P. 6-[1-(4-
942 Fluorophenyl)methyl-1H-pyrrol-2-yl]-2, 4-dioxo-5-hexenoic acid ethyl ester a novel diketo
943 acid derivative which selectively inhibits the HIV-1 viral replication in cell culture and the
944 ribonuclease H activity in vitro. *Antiviral Res.* 2005, 65, 117–124.
- 945 (23) Esposito, F.; Corona, A.; Tramontano, E. HIV-1 reverse transcriptase still remains a
946 new drug target: structure, function, classical inhibitors and new inhibitors with innovative
947 mechanisms of actions. *Mol. Biol. Int.* 2012, 2012, 586401.
- 948 (24) Tramontano, E. HIV-1 RNase H: Recent progress in an exciting, yet little explored,
949 drug target. *Mini-Rev. Med. Chem.* 2006, 6, 727– 737.
- 950 (25) Tramontano, E.; Di Santo, R. HIV-1 RT-associated RNase H function inhibitors:
951 recent advances in drug development. *Curr. Med. Chem.* 2010, 17, 2837–2853.
- 952 (26) Kharlamova, T.; Esposito, F.; Zinzula, L.; Floris, G.; Cheng, Y.-C.; E. Dutschman, G.;
953 Tramontano, E. Inhibition of HIV-1 ribonuclease H activity by novel frangula-emodine
954 derivatives. *Med. Chem.* 2009, 5, 398–410.
- 955 (27) Tramontano, E.; Kharlamova, T.; Zinzula, L.; Esposito, F. Effects of new quinizarin
956 derivatives on both HCV NS5B RNA polymerase and HIV-1 reverse transcriptase
957 associated ribonuclease H activities. *J. Chemother.* 2011, 23, 273–276.
- 958 (28) Esposito, F.; Corona, A.; Zinzula, L.; Kharlamova, T.; Tramontano, E. New
959 anthraquinone derivatives as inhibitors of the HIV-1 reverse transcriptase-associated
960 ribonuclease H function. *Chemotherapy* 2012, 58, 299–307 DOI: 10.1159/000343101.

- 961 (29) Suchaud, V.; Bailly, F.; Lion, C.; Tramontano, E.; Esposito, F.; Corona, A.; Christ, F.;
962 Debyser, Z.; Cotellet, P. Development of a series of 3-hydroxyquinolin-2(1H)-ones as
963 selective inhibitors of HIV-1 reverse transcriptase associated RNase H activity. *Bioorg. Med.*
964 *Chem. Lett.* 2012, 22, 3988–3992.
- 965 (30) Distinto, S.; Esposito, F.; Kirchmair, J.; Cardia, M. C.; Gaspari, M.; Maccioni, E.;
966 Alcaro, S.; Markt, P.; Wolber, G.; Zinzula, L.; Tramontano, E. Identification of HIV-1 reverse
967 transcriptase dual inhibitors by a combined shape-, 2D-fingerprint- and pharmacophore-
968 based virtual screening approach. *Eur. J. Med. Chem.* 2012, 50, 1–14.
- 969 (31) Costi, R.; Metifiot, M.; Esposito, F.; Cuzzucoli Crucitti, G.; Pescatori, L.; Messori, A.;
970 Scipione, L.; Tortorella, S.; Zinzula, L.; Novellino, E.; Pommier, Y.; Tramontano, E.;
971 Marchand, C.; Di Santo, R. 6-(1-Benzyl-1H-pyrrol-2-yl)-2,4-dioxo-5-hexenoic acids as dual
972 inhibitors of recombinant HIV-1 integrase and ribonuclease H, synthesized by a parallel
973 synthesis approach. *J. Med. Chem.* 2013, 56, 8588–8598.
- 974 (32) Costi, R.; Metifiot, M.; Suhman, C.; Cuzzucoli Crucitti, G.; Maddali, K.; Pescatori, L.;
975 Messori, A.; Madia, V. N.; Pupo, G.; Scipione, L.; Tortorella, S.; Di Leva, F. S.; Cosconati,
976 S.; Marinelli, L.; Novellino, E.; Le Grice, S. F. J.; Corona, A.; Pommier, Y.; Marchand, C.; Di
977 Santo, R. Basic quinolinonyl diketo acid derivatives as inhibitors of HIV integrase and their
978 activity against RNase H function of reverse transcriptase. *J. Med. Chem.* 2014, 57,
979 3223–3234.
- 980 (33) Hazuda, D. J.; Felock, P.; Witmer, M.; Wolfe, A.; Stillmock, K.; Grobler, J. A.;
981 Espeseth, A.; Gabryelski, L.; Schleif, W.; Blau, C.; Miller, M. D. Inhibitors of strand transfer
982 that prevent integration and inhibit HIV-1 replication in cells. *Science* 2000, 287, 646–650.

- 983 (34) Van Leusen, A. M.; Siderius, H.; Hoogenboom, B. E.; Van Leusen, D. A new simple
984 synthesis of the pyrrole ring system from Michael acceptors and tosylmethylisocyanides.
985 Tetrahedron Lett. 1972, 52, 5337–5340.
- 986 (35) Anguillera, A.; Alcantara, A. R.; Marinas, M. J.; Sinistera, V. Ba(OH)₂ as the catalyst
987 in organic reaction. Part XIV. Mechanism of Claisen–Schmidt condensation in solid liquid
988 conditions. Can. J. Chem. 1987, 65, 1165–1171.
- 989 (36) Sechi, M.; Bacchi, A.; Carcelli, M.; Sechi, M.; Bacchi, A.; Carcelli, M.; Compari, C.;
990 Duce, E.; Fisicaro, E.; Rogolino, D.; Gates, P.; Derudas, M.; Al-Mawsawi, L. Q.; Neamati,
991 N. From ligand to complexes. Inhibition of human immunodeficiency virus tipe 1 by β-diketo
992 acid metal complexes. J. Med. Chem. 2006, 49, 4248–4260.
- 993 (37) Drysdale, M. J.; Hind, S. L.; Jansen, M.; Reinhard, J. F., Jr. Synthesis and SAR of 4-
994 aryl-2-hydroxy-4-oxobut-2-enoic acids and esters and 2-amino-4-aryl-4-oxobut-2-enoic
995 acids and esters: potent inhibitors of kynurenine-3-hydroxylase as potential neuroprotective
996 agents. J. Med. Chem. 2000, 43, 123–127.
- 997 (38) Nongkhilaw, R. L.; Nongrum, R.; Myrboh, B. Synthesis of substituted hexa-3,5-dienoic
998 acid methyl esters from conjugated dienones. J. Chem. Soc., Perkin Trans. 1 2001,
999 1300–1303.
- 1000 (39) Fujishita, T.; Yoshinaga, T.; Sato, A. PCT Int. Appl. WO-00/ 39086, 2000.
- 1001 (40) Corona, A.; Di Leva, F. S.; Thierry, S.; Pescatori, L.; Cuzzucoli Crucitti, G.; Subra, F.;
1002 Delelis, O.; Esposito, F.; Rigogliuso, G.; Costi, R.; Cosconati, S.; Novellino, E.; Di Santo, R.;
1003 Tramontano, E. Identification of highly conserved residues involved in inhibition of HIV-1
1004 RNase H function by diketo acid derivatives. Antimicrob. Agents Chemother. 2014, 58,
1005 6101–6110.

- 1006 (41) Alberola, A.; Jose, M. A.; Gonzalez, A.; Pedrosa, R.; Vicente, M. Differential reactivity
1007 of β -amino enones and 3-(dimethylamino)-acrylaldehyde towards α -amino derivatives. *J.*
1008 *Chem. Soc., Perkin Trans. 1* 1990, 10, 2681–2685.
- 1009 (42) Di Santo, R.; Costi, R.; Artico, M.; Ragno, R.; Greco, G.; Novellino, E.; Marchand, C.;
1010 Pommier, Y. Design, synthesis and biological evaluation of heteroaryl diketohexenoic and
1011 diketobutanoic acids as HIV-1 integrase inhibitors endowed with antiretroviral activity.
1012 *Farmaco* 2005, 60, 409–417.
- 1013 (43) Tramontano, E.; Cheng, Y.-C. HIV-1 reverse transcriptase inhibition by a
1014 dipyridodiazepinone derivative: BI-RG-587. *Biochem. Pharmacol.* 1992, 43, 1371–1376.
- 1015 (44) Esposito, F.; Kharlamova, T.; Distinto, S.; Zinzula, L.; Cheng, Y.-C.; Dutschman, G.;
1016 Floris, G.; Markt, P.; Corona, A.; Tramontano, E. Alizarine derivatives as new dual inhibitors
1017 of the HIV-1 reverse transcriptase(RT)-associated DNA polymerase and ribonuclease H
1018 (RNase H) activities effective also on the RNase H activity of non-nucleoside resistant RTs.
1019 *FEBS J.* 2011, 278, 1444–1457.
- 1020 (45) Metifiot, M.; Maddali, K.; Johnson, B. C.; Hare, S.; Smith, S. J.; Zhao, X. Z.; Marchand,
1021 C.; Burke, T. R., Jr.; Hughes, S. H.; Cherepanov, P.; Pommier, Y. Activities, crystal
1022 structures, and molecular dynamics of dihydro-1H-isoindole derivatives, inhibitors of HIV-1
1023 integrase. *ACS Chem. Biol.* 2013, 8, 209–217.
- 1024 (46) Chung, S.; Wendeler, M.; Rausch, J. W.; Beilhartz, G.; Gotte, M.; O’Keefe, B. R.;
1025 Bermingham, A.; Beutler, J. A.; Liu, S.; Zhuang, X.; Le Grice, S. F. Structure–activity
1026 analysis of vinylogous urea inhibitors of human immunodeficiency virus-encoded
1027 ribonuclease H. *Anti-microb. Agents Chemother.* 2010, 54, 3913–3921.
- 1028
- 1029

Table 1. Cytotoxicity, Enzymatic, and Antiviral Activities of Compounds 6a–l and 7a–m

| compd | R | R ₁ | R ₂ | X | activity in enzyme assay, | | antiviral activity | | SI ^f |
|-----------------|--------------------|--------------------|-----------------|-----------------|-------------------------------|-----------------|-------------------------------|-------------------------------|-----------------|
| | | | | | IC ₅₀ ^a | | EC ₅₀ ^d | CC ₅₀ ^e | |
| | | | | | RH ^b | ST ^c | | | |
| 6a | Ph | H | OH | Et | 10 | 0.42 | 0.56 | 11 | 19.6 |
| 6b | Ph | Me | OH | Et | | | | | |
| 6c | H | H | OH | Et | >100 | 1.6 | 0.90 | >50 | >55 |
| 6d | Ph | H | NH ₂ | Et | 72 | NT | 19 | >50 | |
| 6e | 4-FBz ^g | Bn ^h | | Et | >100 | 4.3 | 19 | >50 | |
| 6f | 4-FBz ^g | Ph | | Et | 1.8 | 1.2 | 20 | >50 | |
| 6g | 4-FBz ^g | Me | | Et | 28 | >333 | 48 | >50 | |
| 6h | Ph | 4-FBn ^h | | Et | 13.4 | 2.5 | | | |
| 6i | H | | | Et | 55 | 90 | 50 | >50 | |
| 6j | Ph | | | Et | 3.0 | >21 | 4.3 | 26.9 | 6.3 |
| 6k | H | | | COOEt | 21 | 0.51 | 1.2 | 33 | 27 |
| 6l | Ph | | | COOEt | 6.0 | 0.79 | 0.70 | 3.9 | 6 |
| 7a | Ph | H | OH | H | 7.5 | 0.022 | 0.66 | >50 | >75 |
| 7b | Ph | CH ₃ | OH | H | 64 | >111 | >50 | >50 | |
| 7c | H | H | OH | H | 41 | 0.024 | 0.58 | >50 | >86 |
| 7d | Ph | H | NH ₂ | H | 2.0 | 0.043 | 0.63 | >50 | >79 |
| 7e | 4-FBz ^g | Bn ^h | | H | 7.5 | 0.063 | >50 | | |
| 7f | 4-FBz ^g | Ph | | H | 100 | 0.59 | >50 | | |
| 7g | 4-FBz ^g | Me | | H | 20 | | | | |
| 7h | Ph | 4-FBn ^h | | H | 23 | 0.066 | | | |
| 7i | H | | | H | 69 | 26 | >50 | >50 | |
| 7j | Ph | | | H | 7.0 | 0.73 | 17.2 | >50 | >2.9 |
| 7k ⁱ | H | | | COOH | 54 | 0.057 | 1.0 | 28 | 28 |
| 7l | Ph | | | COOH | 14 | 0.019 | 0.7 | >50 | >72 |
| 7m | H | | | Tr ^j | 26 | 0.11 | 20.4 | >50 | |
| 1 | | | | | | 0.007 | 0.016 | >250 | >15000 |
| 2 | | | | | 3.2 | 1.9 | >50 | | |
| 3 | | | | | 8 | 98 | <0.2 | >50 | >250 |
| 4 | | | | | 3 | 0.60 | 2 | >50 | >25 |
| 5 | | | | | 26.2 | 2.4 | 3.6 | >50 | >13.8 |

1030

1032 ^aInhibitory concentration 50% (μM) determined from dose–response curves. ^bExperiments performed against1033 HIV-1 RT-associated RNase H activity. ^cExperiments performed against HIV-1 IN ST activity. ^dEffective1034 concentration 50% (μM). ^eCytotoxic concentration 50% (μM). ^fSI = CC₅₀/EC₅₀. ^gBz, benzoyl. ^hBn, benzyl. ⁱSee1035 also ref 33. ^jTr, triazolyl.

1036

1037


1038

1039

1040

1041

1042



1044


1045

1046 **Figure 1.** Scatter plot for the inhibition of RNase H and IN enzymes. (A) Compounds are categorized according
 1047 to their acidic or ester function. (B) Compounds are categorized according to the nature of their R substitution.
 1048 (C) Compounds are categorized according to the nature of their R1 substitution. (D) Compounds are
 1049 categorized according to the nature of their R2 substitution. Compounds with one IC_{50} value missing such as
 1050 6d have been left out of the plot, and compounds with IC_{50} values above 111 μ M have been arbitrary
 1051 positioned at the 100 μ M value.

1052

1053

1054



1056


1057 a Reagents and conditions: (i) TosMIC, NaH, Et₂O/DMSO, room temp, 1 h; (ii) tri uoroacetic acid, 80
 1058 °C, 24 h; (iii) 4-F-benzyl bromide, K₂CO₃, DMF, 100 °C, 24 h; (iv) diethyl oxalate, C₂H₅ONa, THF, room temp,
 1059 2 h; (v) CH₃COONH₄, benzene, glacial acetic acid, reflux, 20 h; (vi) 1 N NaOH, THF/CH₃OH, room temp, 1
 1060 h.

1061

1062

1063

1064 **Scheme 2.** Synthetic Route to Pyrrolyl DKAs 6e-h and 7e-h^a



1075 ^aReagents and conditions: (i) alkylating agent, K₂CO₃, DMF, 100 °C, 24 h; (ii) phenylboronic acid, Cs₂CO₃,
 1076 P(t-But)₃, Pd₂(dba)₃, dioxane, 80 °C, 24 h; (iii) (1) DMF, 1,2-dichloroethane dry, oxalyl chloride, 0 °C, 15 min,
 1077 room temp, 15 min; (2) 4-F-benzoyl chloride, AlCl₃, room temp, 4 h; (iv) phenylboronic acid, copper(II) acetate
 1078 anhydrous, pyridine/NMP 1:1, microwave at 60 W, 120 °C, 6 min; (v) acetone, 4 N NaOH, room temp, 24 h;
 1079 (vi) diethyl oxalate, C₂H₅ONa, THF, room temp, 2 h; (vii) 1 N NaOH, THF/CH₃OH, room temp, 1 h.

1080

1081

1082

1083


1084

1085

1086

1087

1088 **Scheme 3.** Synthetic Route to Pyrrolyl DKAs 6i,j and 7i,j



1089

1090 aReagents and conditions: (i) 4-F-benzyl bromide, K₂CO₃, DMF, 100 °C, 24 h; (ii) trifluoroacetic acid, 80 °C,
 1091 24 h; (iii) acetone, 4 N NaOH, room temp, 24 h; (iv) Et₂O/DMSO, NaH, TosMIC, room temp, 1 h; (v) 4-F-benzyl
 1092 bromide, K₂CO₃, DMF, 100 °C, 24 h; (vi) diethyl oxalate, C₂H₅ONa, THF, room temp, 2 h; (vii) 1 N NaOH,
 1093 THF/CH₃OH, room temp, 1 h.


1094

1095

1096

1097

1098 **Scheme 4.** Synthetic Route to Pyrrolyl DKAs 6k,l and 7k,l



1106 aReagents and conditions: (i) NIS, acetone, $-78\text{ }^{\circ}\text{C}$, 96 h; (ii) phenylboronic acid, Cs_2CO_3 , $\text{P}(\text{t-Bu})_3$,
1107 $\text{Pd}_2(\text{dba})_3$, dioxane, $80\text{ }^{\circ}\text{C}$, 24 h; (iii) diethyl oxalate, $\text{C}_2\text{H}_5\text{ONa}$, THF, room temp, 2 h; (iv) 1-trityl-1H-
1108 [1,2,4]triazole-3-carboxylic acid ethyl ester, 36 $n\text{-BuLi}$, THF, from -78 to $0\text{ }^{\circ}\text{C}$, 3.5 h; (v) 3 M HCl sol, 1,4-
1109 dioxane, $60\text{ }^{\circ}\text{C}$, 30 min; (vi) 1 N NaOH, THF/ CH_3OH , room temp, 1 h.

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122 **Chart 1.** Selective Inhibitor of IN Enzyme (1), First Described Dual IN/RNase H Inhibitor (2), and Recently
 1123 Discovered Dual IN/RNase H Inhibitors 3–5

1124

1125

1126

1127

1128

1129

1130

1131

1132 **Chart 2.** Newly Designed Pyrrolyl DKA Derivatives 6a–l and 7a–m

1133

